

WEAVE SPRINT POWERED BY TRIJOG
PAPER III · VOLUME ONE

MD and DNB psychiatry, theory revision

Speciality Psychiatry

Milestones, Disability and Autism

How a child normally develops, and what counts as a deviation worth acting on. Then intellectual disability and autism spectrum disorder.

Dr. Wilfred D'souza · Dr. Niharika Reddy

WEAVE POWERED BY TRIJOG
CENTRE FOR INTEGRATIVE PSYCHIATRY

When therapy is part of the plan, there is somewhere to refer.

For clinicians who assess, prescribe or manage care but do not provide psychotherapy themselves. Trijog gives patients a route to online therapy and in-person sessions in Mumbai.



A CLINICIAN REFERRAL ROUTE



Refer to Trijog for therapy.

Your patient can contact Trijog directly. Our team can help them choose online or in-person care and identify a therapist based on their needs and preferences. The patient decides whether to book and whether the fit is right.

booking.trijog.com • 1800-2027-180 • the Trijog app

Online therapy

For patients who prefer to meet remotely.

In person in Mumbai

Powai, Bandra and Prabhadevi.

Across ages

Adults, children and teenagers.

Across relationships

Individuals, couples and families.

What the referral route keeps clear



01

Your role

You remain the referring clinician. A therapist takes on the psychotherapy work.



02

The fit

Concern, age, language, format and preferred way of working can all shape the match.



03

The setting

Your patient can meet online, or in person at one of Trijog's Mumbai clinics.



04

The coordination

If your patient wants it, their clinicians can share what is needed to keep care aligned.

Weave, powered by Trijog

PSYCHIATRY VERTICAL

Weave is the psychiatry vertical of Trijog. Weave Sprint is part of its Psychiatry Outreach Programme, supporting psychiatry residents and clinicians with practical learning and a clear route into Trijog's wider care network.

The notes in your hands are part of that programme. Therapy through Trijog can sit alongside psychiatric care when the patient and clinicians choose.

You do not have to provide therapy yourself to make it part of good care.

Share this route with a patient

Online therapy or in-person care in Mumbai.

booking.trijog.com • 1800-2027-180

Keep consent visible

Any coordination between clinicians happens with the patient's permission. The patient remains free to choose whether to book.

Routine referrals only. This is not an emergency route. For urgent risk, follow your local emergency and safeguarding pathway.

About this insert. Weave Sprint is educational outreach from Weave, the psychiatry vertical of Trijog. Therapy is provided through Trijog.

WEAVE SPRINT · PAPER III

Speciality Psychiatry



THE WEAVE SPRINT SERIES

TITLE

Weave Sprint, Paper III, Volume One: Milestones, Disability and Autism
Part of the Weave Sprint series, MD and DNB psychiatry theory revision.

WRITTEN BY

Dr. Wilfred D'souza and Dr. Niharika Reddy

PUBLISHED BY

Weave, Centre for Integrative Psychiatry · weave.clinic

EDITION

First compiled 2026. Free to read, share and print for study.

The clinical content is written for doctors and doctors in training. It is a study and revision aid, not a substitute for current guidelines or clinical judgement. Verify doses and thresholds against the current references and your local protocols before any clinical use.

*For every resident who has ever sat up the night before,
afraid it will not be enough.*

A word of thanks

I want to begin by thanking something I cannot quite name. Call it whatever you like. I have never been certain what to call it, and I have stopped needing to be. There is a part of this work that does not feel like mine. The line that arrives before I have thought of it, the connection I did not plan, the steadiness on a morning I had no reason to be steady. I only hold the pen. Something moves through the holding, and I have learned to thank it without asking it for a name.

Someone once asked me why I give all of this away. Why hand out the notes I sat up making, to the very people sitting the same exam beside me. It is a fair question. If I had kept them, I would probably have scored a few marks more than the next person, and I might have been thought of as the better student. I have turned that over honestly, and I have decided it is a small thing to want. What I get instead, by putting the work in your hands, is the chance to take a little of the fear out of a great many nights. That trade is not close. Easing that fear is not a detour on the way to becoming a good psychiatrist. It is the thing itself. I did not come here to be the best student in the room. I came here to be a good psychiatrist.

And to Niharika, who reads everything first and knows when to be honest and when to wait; and to the residents who study from these pages and write back with the corrections and the questions: thank you. The book is better for you, and so am I.

Dr. Wilfred D'souza

Foreword

This first volume begins with what normal development looks like, because every judgement that follows is a judgement of deviation from it. The neurodevelopmental disorders come next, intellectual disability and autism, where the diagnosis is made against the timetable set out in the chapter before. You cannot call something a delay until you know the schedule it is late against.

The notes were written to be understood, not just revised. Each chapter is a full account of its subject, from the mechanism up to the point the exam tests, so that you can read to learn the first time and read to recall every time after. Where a fact is worth holding exactly, it is marked. Where a mistake is easy to make, it is named before you make it. The colour code on the next pages tells your eye what kind of fact it is looking at before your mind has to decide, and after a few pages it stops being a key you consult and becomes a sense you carry.

Read it in order if you are meeting the material, or open it anywhere if you are returning to it. The contents at the front and the index at the back both jump to the page you want. However you use it, the hope held in these pages is plain: that one ordered, trustworthy place to study from makes the work a little lighter, and that you walk into the hall steadier than you would have otherwise.

The colour memory code

Every note in this volume speaks one visual language, so once you learn it in the first chapter you read the rest without thinking about it.

 Concept and rule	The core idea and the principle that follows from it. When a line is petrol, it is the thing to understand, and the thing worth being able to state.
 Key number	A figure to hold exactly: a dose, a threshold, a prevalence, a cut-off. Amber means memorise this number, not just the shape of it.
 Trap	The mistake the exam is waiting for. When a line is coral, it is warning you off an error that looks right until it costs you the mark.

Hold this is the single line from a topic you must carry out of it. **Good to remember** gathers, for each named thing, what it is, when it matters, and the one number or formula worth keeping. Both help a tired brain find the centre of a dense page fast.

The contents at the front is clickable, and so is the index at the back: every entry jumps to its page. Read to learn once, and to recall as often as you need.

Contents

FRONT MATTER	
Foreword	7
How to use this book	8
PART I Normal Child Development and Milestones	
The developmental frameworks	11
The domains and the milestones	84
Development in context and clinical use	135
Apply and consolidate	154
Previously asked	162
Quick review	164
References	167
PART II Intellectual Disability and Autism Spectrum Disorder	
Intellectual disability: the concept and its measurement	171
The causes of intellectual disability	212
Living with intellectual disability	261
Autism: concept, causes and criteria	288
The clinical picture of autism and how it is recognised	315
Managing autism and its outcome	343
Discriminate, apply and consolidate	382
BACK MATTER	
Subject index	400
About the authors	408

PART I



Normal Child Development and Milestones

How a child normally develops: the domains, the milestones and the ages, and what counts as a deviation worth acting on.

01 The developmental frameworks

02 The domains and the milestones

03 Development in context and clinical use

04 Apply and consolidate

05 Previously asked

06 Quick review

07 References

A decorative background graphic consisting of concentric dashed circles in orange and grey, with radial lines extending from the center. At the bottom, there is a wavy line pattern in grey.

PAPER III

Normal child development & milestones

Three bands . one complete notes document

The developmental frameworks

1 . Piaget stages of cognitive development

Jean Piaget described cognitive development as an active construction of knowledge, not the passive accumulation of facts. The child acts on the world, builds organised patterns of understanding, finds that some patterns no longer explain experience, and modifies them. The theory therefore asks a more useful question than "What does the child know?": *what form of reasoning is available to the child, and what kind of problem can that reasoning solve?* Piaget proposed an invariant sequence of four broad organisations of thought: sensorimotor, preoperational, concrete operational and formal operational. The sequence is the examinable backbone. The age bands attached to it are textbook descriptors with source variation, not developmental screening cut-offs and never referral thresholds.

How to use this page. First learn the direction of change: action becomes mental representation; representation becomes logical but remains tied to concrete situations; logic then extends to possibilities, propositions and hypotheses. Next learn the characteristic achievement and characteristic error of each stage. Finally, separate three questions that are often collapsed in examinations and clinics: Piagetian stage theory describes the *organisation of cognition*; milestones describe observable developmental attainments across several domains; developmental surveillance and screening decide whether a child's course needs closer evaluation. A child does not "fail Piaget" because one conservation task is failed, and a Piagetian task is not a developmental screening instrument.

Must memorise: sensorimotor action becomes preoperational representation, concrete operations make representation logical for real and tangible problems, and formal operations extend logic to abstract and hypothetical possibilities.

MNEMONIC

"Sensorimotor - Preoperational - Concrete operational - Formal operational"

Sensorimotor action becomes preoperational representation, concrete operations make representation logical for real and tangible problems, and formal operations extend logic to abstract and hypothetical possibilities.

4 BROAD ORGANISATIONS OF THOUGHT	birth to 2 years SENSORIMOTOR (BERK)	2 to 7 years PREOPERATIONAL (BERK)	7 to 11 years CONCRETE OPERATIONAL (BERK)
---	--	--	--




Fig 20.1 The developmental theories aligned on a shared age axis. Freud, Erikson and Piaget after Berk; Mahler after Wolberg; Kohlberg's levels as descriptive bands, since no numeric ages are sourced. Vygotsky carries no row because his theory is not stage-based. Boundaries are approximate and differ between texts.




Fig 20.2 Piaget's four stages on an age axis, with the central operation each stage permits and the error it explains. The saffron spine follows Berk, Development Through the Lifespan; the coral bars are Tasman Psychiatry's differing bands for the same stages, shown rather than reconciled. No source states any band as a median, a normal limit or a referral threshold, and no source in the corpus gives Tasman a sensorimotor band.

1.1 The engine of Piaget's theory: schemes, adaptation and equilibration

A **scheme** is an organised pattern for acting on, representing or thinking about experience. An infant may have a grasping scheme; an older child may have a scheme for classifying animals or balancing an equation. Development occurs through **adaptation**, which has two complementary processes. In **assimilation**, the child interprets a new event through an existing scheme, as when every four-legged animal is initially treated as a dog. In **accommodation**, the scheme itself changes because the new event does not fit, so dogs and cows become distinct categories. Assimilation preserves continuity; accommodation permits novelty.

Organisation integrates separate schemes into larger cognitive systems. **Equilibration** is the regulatory process that moves thought from a less adequate balance to a more adequate one: a contradiction produces disequilibrium, and reorganisation restores coherence at a higher level. The theory is therefore *constructivist*. The child is neither a miniature adult nor an empty vessel, but an active reasoner whose errors disclose the structure of thought. In an examination, do not define a stage merely by its age band. Define it by the new operation it permits and the systematic limitation that remains.

■ **RULE** Stage is inferred from the structure of reasoning, not from the answer alone. Ask the child to explain, compare, predict and justify. A correct answer obtained by guessing or rote learning does not demonstrate an operation; an incorrect answer produced by misunderstanding the language of the task does not prove that the operation is absent.

1.2 Sensorimotor stage: knowledge through action

Berk's *Development Through the Lifespan* lists the **sensorimotor stage** as **birth to 2 years**. This wording is a textbook stage band, not a threshold. Cognition is initially organised through sensory experience and motor action. Repeated actions become circular reactions; schemes are coordinated into intentional means-end behaviour; experimentation becomes increasingly active; and, eventually, action can be represented mentally before it is performed. Piaget divided this progression into six substages, but the clinically useful principle is the movement from reflex-linked activity to deliberate action and internal representation.

The central achievement is **object permanence**, the understanding that an object continues to exist when it is no longer perceived. Before that representation is secure, searching is governed heavily by the immediate perceptual and motor situation. The classic **A-not-B error** occurs when an infant repeatedly retrieves an object from one location and then searches there again after seeing it hidden elsewhere. Later achievements include intentional problem-solving, imitation of an absent model, and the capacity to anticipate an outcome mentally. These are not isolated tricks. Together they mark the beginning of representation, which makes language, pretend play and later symbolic reasoning possible.

Clinical use. Observe how the child explores, searches, imitates and coordinates means with ends rather than relying on a verbal question. Interpretation must remain modest: visual access, motor ability, attention, fatigue, anxiety and familiarity with the task can all alter performance. Object-permanence tasks are demonstrations of a cognitive construct, not stand-alone diagnostic tests.

1.3 Preoperational stage: symbolic thought without reversible operations

Berk lists the **preoperational stage** as **2 to 7 years**. Tasman Psychiatry summaries use 2 to 6 years and 3 to 6 years in separate presentations. These variants should be retained as source conventions, not averaged into a new boundary. The decisive gain is the ability to use one thing to stand for another. Language expands, drawings carry meaning, and pretend play allows a block to become a car or an empty cup to contain imaginary tea. Thought has become representational, but it is not yet governed consistently by logical operations.

The characteristic limitations are linked. **Egocentrism** is difficulty coordinating one's own perspective with another person's perspective; it is not selfishness. The three-mountains task illustrates this failure of perspective-taking. **Centration** is attention to one striking dimension while other relevant dimensions are neglected. **Irreversibility** is difficulty mentally undoing a transformation. Together they explain failure of conservation: after the same quantity is rearranged into a taller or more spread-out appearance, the child judges by the perceptually dominant feature rather than by identity, compensation and reversibility.

Other characteristic forms include animism, in which life or intention is attributed to inanimate objects; artificialism, in which natural events are understood as made for human purposes; and transductive reasoning, in which one particular event is linked directly to another without a valid general principle. In clinical work, such reasoning can make illness, separation or family conflict feel caused by the child's thought or behaviour. The psychiatrist should explore the child's meaning and gently correct self-blame rather than treat developmentally shaped magical explanations as delusions.

1.4 Concrete operational stage: logical operations tied to tangible content

Berk lists the concrete operational stage as **7 to 11 years**; Tasman Psychiatry summaries use 6 to 12 years and 6 to 11 years. Again, these are alternate textbook mappings, not clinical cut-offs. The child can now perform an operation, an internalised mental action that is reversible and coordinated with other actions. Perceptual appearance no longer dominates every judgement, provided the problem concerns familiar or concretely imaginable material.

OPERATION	WHAT IT PERMITS
Conservation	Becomes possible because the child can use identity, compensation and reversibility.
Decentration	Permits simultaneous attention to more than one dimension.
Classification and class inclusion	Permit an item to belong to nested categories.
Seriation and transitive inference	Seriation orders objects along a dimension, and transitive inference coordinates relations to infer an unobserved relation.

Perspective-taking becomes more flexible, rules can be understood as coordinated systems, and explanations become less dependent on a single salient feature.

The limitation is contained in the name: the operations are most reliable when they can be anchored in actual objects, familiar events or imaginable transformations. Purely hypothetical premises, propositions contrary to fact, and systematic testing of all possible combinations remain difficult. For interviewing, use concrete language, timelines, drawings, demonstrations and examples from the child's experience. Abstract labels such as "mood," "risk" or "consequences" may need to be unpacked into observable situations.

1.5 Formal operational stage: reasoning about possibilities

Berk lists the formal operational stage as **11 years onward**, whereas a Tasman Psychiatry summary lists 13 years to adulthood. The two descriptions must not be collapsed into a falsely precise onset. The qualitative advance is that reasoning can operate on propositions and possibilities, not only on directly represented objects. The person can accept a hypothetical premise, derive its implications, and judge whether the conclusion follows even when the premise is not factually true.

- Hypothetico-deductive reasoning begins with possible explanations, derives predictions and tests variables systematically.
- Combinatorial reasoning considers the possible combinations within a problem rather than changing factors haphazardly.
- Propositional thought evaluates the logical relation between statements.
- Metacognition permits reflection on one's own thinking.

These abilities support scientific reasoning, long-range planning, moral debate, identity exploration and consideration of alternative futures.

Formal reasoning is not an all-or-none possession applied uniformly across every subject. Education, cultural tools, practice, interest, emotional arousal and domain knowledge shape whether it appears in a particular task. An adolescent or adult may reason formally in a familiar academic domain and concretely in an unfamiliar or emotionally charged one. The stage should therefore guide the form of questioning, not substitute for an assessment of actual comprehension, decision-making or capacity.

■ **TRAP** **Do not turn Piaget's age bands into milestone norms.** The cited texts do not label these bands as medians, normal limits or referral ages. They describe theoretical stages. Source variation, uneven performance and task demands make it incorrect to diagnose delay from a stage label or from one failed Piagetian task.

1.6 The classic tasks: what each task actually tests

A Piagetian task is valuable because the pattern of explanation reveals an underlying operation.

TASK	WHAT IT TESTS
Object-permanence task	The issue is whether the hidden object is represented despite absence from perception.
Perspective-taking task	The issue is coordination of viewpoints.
Conservation task	The examiner changes appearance while preserving quantity, then asks for both judgement and justification.
Class-inclusion task	The child must coordinate a subclass with the larger class.
Seriation and transitivity tasks	Relations must be ordered and mentally coordinated.
Formal-reasoning task	The examiner looks for generation of hypotheses and systematic control of variables.

Administration matters. Establish that the child understands the words, can see and manipulate the materials, and has not interpreted repetition of the question as a signal that the first answer was wrong. Ask "How do you know?" and record the reasoning rather than scoring only correct or incorrect. Repeating the same transformation with familiar objects can distinguish a conceptual limitation from confusion about the task. Do not coach the target operation and then claim to have measured spontaneous competence.

The best viva formulation is: *achievement, error, mechanism, task, limitation*. For preoperational thought, for example, the achievement is symbolic representation; the error is failure of conservation; the mechanism includes centration and irreversibility; the task is transformation of equal quantities; and the limitation is reliance on perceptual appearance. This structure prevents a recital of ages without psychology.

1.7 Strengths, limitations and contemporary reading

Piaget's enduring achievement was to show that children's errors are organised rather than random. He supplied testable constructs, linked knowledge to the child's own activity, and shifted developmental psychology from a catalogue of deficits to a study of changing cognitive structures. His tasks remain powerful teaching devices because they make abstract constructs visible in behaviour.

The limitations are equally examinable. Performance can improve when language is simplified, memory load is reduced or the situation is made familiar, suggesting that some classic tasks underestimate competence. Development is often gradual, overlapping and domain-specific rather than a clean step from one global structure to another. Social interaction, instruction, culture and the tools valued by a community receive less weight than in sociocultural theories. Formal reasoning is not consistently demonstrated across every person and every domain. The useful modern reading is therefore neither to discard Piaget nor to literalise him: retain the sequence as a model of increasingly coordinated representation and logic, while assessing the child's actual strategy in context.

1.8 Age alignment with Erikson, Freud, Kohlberg, Mahler and Vygotsky

The chapter's theory-alignment plate should be read horizontally: theories may discuss the same period of life while explaining different developmental work. Piaget explains the structure of knowing. Erikson's psychosocial stages explain successive tensions between the self and the social world. Berk places autonomy versus shame and doubt at 1 to 3 years, initiative versus guilt at 3 to 6 years, and industry versus inferiority at 6 to 11 years. Thus symbolic expansion, initiative and later school-age mastery overlap in time but are not interchangeable constructs.

Freud's psychosexual stages organise development around changing instinctual zones and conflicts. Berk lists the anal stage at 1 to 3 years, the phallic stage at 3 to 6 years, and latency at 6 to 11 years. Piaget's account does not explain libido or defence, and Freud's account does not test conservation or operations. Alignment helps recall; it does not imply that one theory validates another.

Kohlberg extended a structural approach to moral development, classifying the reasoning used to justify moral decisions rather than merely whether an answer was socially approved. Kaplan and Sadock describes preconventional morality before adolescence, conventional response levels during and later in adolescence, and a postconventional response level in late adolescence or early adulthood. Moral-stage reasoning and Piagetian cognition are related conceptually, but moral judgement also depends on social experience, values and context.

Mahler's separation-individuation theory concerns the child's emerging psychological separateness in relation to the caregiver. Kaplan and Sadock and Wolberg place the hatching period from about 4 or 5 months to 10 months; Wolberg places rapprochement onset between 15 and 18 months, describes the rapprochement crisis around 18 to 24 months, and places libidinal object constancy during the third year of life. These are the sources' own stage descriptions, not screening thresholds. Clinically, Mahler supplies relational language; Piaget supplies cognitive-structural language.

Vygotsky places greater weight on language, shared activity and culturally transmitted tools. The zone of proximal development is the gap between what a learner can do independently and what the learner can do with skilled support. Guided participation and scaffolding reveal emerging capacity that unsupported performance may conceal. Piaget asks what structure the child has constructed; Vygotsky asks what becomes possible in collaboration. The practical synthesis is dynamic: assess independent reasoning, then observe how performance changes with graduated help.

1.9 Piaget within the domains of child development

Piaget principally addresses cognition, although cognition is expressed through movement, language, play and social interaction. A child with a motor or language limitation may therefore perform poorly on a cognitive task for reasons other than the cognitive operation being tested.

1.10 Clinical use: developmental history, surveillance, red flags and Indian screening

Piaget calibrates the cognitive level of the interview; surveillance, red flags and screening are taught under the red-flags topic.

2 . Erikson psychosocial stages

Erik Erikson described personality development as a lifespan sequence of eight **psychosocial stages**. Each stage is organised around a **psychosocial crisis**, a tension between two necessary tendencies that arises from the interaction of biological maturation, the developing ego and the social world. A crisis is not a psychiatric emergency and it is not a choice in which one pole must be eliminated. It is a developmental turning point. A predominantly adaptive resolution strengthens the ego and yields a characteristic virtue; an unfavourable or markedly one-sided resolution leaves greater vulnerability when later tasks are encountered.

How to use this page. Learn the sequence as a chain of questions: *Can I rely on care? Can I act for myself? May I initiate? Can I master valued work? Who am I? Can I share myself without losing myself? What do I contribute? Can I accept the life I have lived?* Then attach the conflict, virtue, relational setting and common clinical distortion to each question. The age bands are textbook mappings whose normative construct was not specified by the sources. They are not medians, outer limits, screening cut-offs or referral thresholds. The theory-alignment plate may help compare Erikson with Piaget, Freud, Kohlberg, Mahler and Vygotsky, but temporal overlap never makes their constructs interchangeable.

Must memorise: trust gives hope, autonomy gives will, initiative gives purpose, industry gives competence, identity gives fidelity, intimacy gives love, generativity gives care, and integrity gives wisdom.

MNEMONIC

"Trust, autonomy, initiative, industry, identity, intimacy, generativity, integrity"

Trust gives hope, autonomy gives will, initiative gives purpose, industry gives competence, identity gives fidelity, intimacy gives love, generativity gives care, and integrity gives wisdom.




Fig 20.3 Erikson's eight psychosocial stages as conflict pairs. The first four sit on a sourced age axis, Berk's mapping above Tasman Psychiatry's; where the two disagree both are shown rather than averaged. The axis stops at 11 years because neither text gives an Erikson age past it, so the last four stages carry the sources' own life-period words instead of numbers. No boundary here is a referral threshold.

2.1 Architecture of the theory: epigenesis, crisis and ego strength

Erikson extended psychoanalytic development beyond childhood and beyond intrapsychic instinct. His **epigenetic principle** proposes that each psychosocial capacity has a period of relative ascendancy within an ordered sequence, while earlier themes remain active and can be revisited. Trust does not stop when autonomy begins; identity does not become irrelevant after intimacy. Later relationships, illness, migration, parenthood, loss and social change can reactivate earlier tensions and permit partial reworking. The model is therefore stage-like but not mechanically closed.

The two poles are sometimes called syntonic and dystonic tendencies. Healthy development does not mean total trust, autonomy or generativity. Some mistrust protects against exploitation, some shame supports social awareness, and some isolation preserves reflection and boundaries. The task is a workable ratio in which the adaptive tendency predominates without erasing the protective information carried by the other pole. The resulting **ego strength** is not a symptom score. It is a durable capacity, such as hope or fidelity, that supports later developmental work.

Erikson's distinctive contribution is the social location of the crisis.

- Caregiving relationships organise trust.
- Limits and bodily control organise autonomy.
- Family and play organise initiative.

- School and culturally valued work organise industry.
- Peers, ideals and social roles become increasingly important for identity.

Culture determines what counts as competence, legitimate independence, adulthood, contribution and a good life. The theory must therefore be used as a formulation scaffold, not as a culturally neutral measuring instrument.

RULE Name four elements for every stage: the conflict, the psychosocial task, the central relational or social field, and the virtue associated with a sufficiently adaptive resolution. Age alone is never an adequate answer.

2.2 Age alignment and how to state it safely

STAGE	BERK, DEVELOPMENT THROUGH THE LIFESPAN	TASMAN PSYCHIATRY
basic trust versus mistrust	birth to 1 year	trust versus mistrust, birth to 12 to 18 months
autonomy versus shame and doubt	1 to 3 years	autonomy versus shame, 12 to 18 months to 36 months
initiative versus guilt	3 to 6 years	initiative versus guilt, 3 to 5 years
industry versus inferiority	6 to 11 years	
identity versus role confusion	adolescence	11 years
intimacy versus isolation	early adulthood	
generativity versus stagnation	middle adulthood	
integrity versus despair	old age	

These are retained as each source states them. They must not be averaged into a new age boundary, treated as competing referral ages or presented as proof that a person has entered a stage on a birthday.

8 stages ONE LIFESPAN SEQUENCE	Birth to 1 year BERK TRUST MAPPING	1 to 3 years BERK AUTONOMY MAPPING	3 to 6 years BERK INITIATIVE MAPPING
-----------------------------------	---------------------------------------	---------------------------------------	---

2.3 Trust and autonomy: hope, will and the beginning of agency

Basic trust versus mistrust asks whether the world is sufficiently reliable for hope. The infant depends on caregivers for regulation, protection and the repeated relief of distress. Sensitive and reasonably predictable care supports a felt expectation that needs can be communicated and that another person may respond. The virtue is hope, the capacity to sustain expectation despite frustration. Mistrust is not produced by every delay or mismatch, and perfect gratification is neither possible nor desirable. The clinically important pattern is whether care is sufficiently coherent for the infant to tolerate absence and uncertainty without the world becoming wholly unsafe.

Trust is relational but not reducible to attachment classification. Attachment theory studies proximity, secure-base behaviour and reunion patterns in particular caregiver relationships; Erikson describes a broader ego expectation about self, others and the social world. Severe inconsistency, neglect or frightening care may compromise both, but a psychiatrist should not infer a specific attachment-pattern from the Eriksonian conflict or diagnose a caregiver from one observed interaction.

Autonomy versus shame and doubt asks whether emerging self-control can coexist with connectedness. Growing motor ability, language, choice, bodily regulation and the repeated use of "no" expand the child's sense of agency. Supportive adults offer bounded choices, tolerate safe attempts, set comprehensible limits and help after failure. The virtue is will, the capacity to choose and persist. Chronic overcontrol, humiliation, ridicule of accidents or demands far beyond capacity can make self-assertion feel dangerous and foster excessive shame or doubt. Permissiveness is not the opposite of shaming control: reliable limits help the child experience autonomy as organised and safe.

In a developmental history, ask how caregivers respond to distress, exploration, refusal, feeding, dressing and bodily-control struggles. Formulate the fit between the child's capacities, temperament and the caregiving environment. Do not attribute current anxiety, dependency or defiance to a single early-stage failure. Erikson supplies a hypothesis about meaning; history, observation and longitudinal evidence must test it.

2.4 Initiative and industry: purpose, competence and the social valuation of action

Initiative versus guilt asks whether the child may imagine, plan and act. Symbolic play, questions, fantasy, imitation of adult roles and self-started activity enlarge the field of action. Initiative is more than independence: it adds direction and purpose. Adults support it by taking ideas seriously, permitting safe experimentation and setting limits without treating curiosity or ambition as badness. The virtue is purpose. Guilt becomes excessive when wishes, questions, rivalry or ordinary attempts at action are repeatedly interpreted as harmful or unacceptable. Some guilt is developmentally useful because it permits concern for the effects of one's actions; incapacitating guilt suppresses initiative itself.

Industry versus inferiority asks whether effort can produce socially valued competence. Schooling, peer comparison, tasks, tools, rules and feedback move the child into a wider field of evaluation. The child learns to sustain attention, practise, cooperate, complete work and take justified pride in mastery. The virtue is competence. Repeated failure, unrecognised learning needs, exclusion, harsh comparison or praise detached from actual effort can undermine industry. Inferiority is not simply low self-esteem. It is the expectation that one's effort will not produce valued mastery.

In Indian clinical practice, marks and rank may dominate the family's account of competence, but industry is wider than examination performance. Practical skill, sport, art, caregiving, language, social contribution and persistence may all organise mastery. A child with ADHD, a specific learning disorder, chronic illness or sensory impairment may be industrious while producing

weak conventional outputs; those conditions themselves are developed in the ADHD and specific learning/communication disorders notes. Formulation should separate ability, opportunity, effort, accommodations and the social meaning attached to performance.

2.5 Identity versus role confusion: the central crisis of adolescence

Identity versus role confusion asks whether the adolescent can experience continuity while exploring possible selves. Identity formation integrates past identifications, bodily change, values, relationships, vocation, culture and imagined futures into a sufficiently coherent answer to "Who am I, and where am I going?" The virtue is fidelity, the capacity to commit to people and ideals while tolerating difference and remaining recognisably oneself. Role confusion may appear as unstable commitments, incompatible self-definitions, chronic diffusion or an identity chosen mainly to meet others' demands.

Erikson's **psychosocial moratorium** is a socially permitted interval for exploration before durable commitments are expected. It is not mere idleness. Marcia operationalised the identity theme along two dimensions, exploration and **commitment**, producing four identity statuses: diffusion, foreclosure, moratorium and achievement. These are statuses rather than an invariant stage sequence.

IDENTITY STATUS	DESCRIPTION
Foreclosure	describes commitment without meaningful exploration.
Moratorium	describes active exploration without settled commitment.
Achievement	combines exploration with commitment.
Diffusion	lacks both.

A person's status may differ across vocation, values and relationships.

Gender identity is a person's experienced sense of gender. It is distinct from gender expression, sexual orientation and sexual behaviour, and none of these should be treated as evidence of role confusion. Identity exploration can be developmentally normal. The clinical task is to assess distress, coercion, stigma, safety, family response, comorbidity and functioning without converting cultural nonconformity into psychopathology. Erikson's language can illuminate coherence and commitment, but it does not determine what a person's identity ought to be.

Psychosexual development is also broader than Freud's sequence of instinctual zones. In adolescence it includes bodily maturation, sexual feelings, relational boundaries, consent, values and the integration of sexuality into identity. The psychiatrist should distinguish normative exploration from exploitation, risky behaviour, compulsivity, trauma-related phenomena, mood-related disinhibition and distress driven by stigma.

The puberty reference plate should be read separately from Erikson's theory-alignment plate. The former describes bodily maturation; the latter aligns theoretical accounts of psychosocial, cognitive, psychosexual, moral and relational development. Physical maturation may alter the

social pressures surrounding identity, but no Tanner stage proves a level of identity formation, moral development or judgement.

2.6 Intimacy, generativity and integrity across adult life

Intimacy versus isolation asks whether identity can enter mutual commitment without dissolution. Berk places it in early adulthood. **Intimacy** includes emotional reciprocity, vulnerability, sexuality, friendship and durable commitment; it is not synonymous with marriage. Erikson's logic is that a sufficiently coherent identity makes it possible to share oneself without either fusion or defensive withdrawal. The virtue is love. Isolation may protect temporarily after loss or threat, but chronic avoidance of mutuality narrows the person's relational world.

Generativity versus stagnation asks what the adult creates, sustains and hands onward. Berk places it in middle adulthood. **Generativity** includes parenting but is not limited to biological parenthood. Teaching, mentoring, clinical service, productive work, institution-building, creativity and care for future generations can all express it. The virtue is care. Stagnation is an excessive narrowing toward self-absorption, repetition or the sense that one's activity does not nourish anything beyond the self.

Integrity versus despair asks whether the life course can be owned as a whole. Berk places it in old age. **Ego integrity** is not the belief that every decision was correct. It is the capacity to accept the singular life one has lived, integrate success with regret and face finitude without repudiating the self. The virtue is wisdom. Despair arises when time feels too short for repair and the life appears wasted, fragmented or intolerably unlike the life that should have been.

Adult developmental crises can organise the meaning of depression, caregiving burden, infertility, retirement, bereavement, migration or occupational disruption, but they do not replace diagnosis. A person without a partner is not necessarily isolated; a person without children is not necessarily stagnant; regret does not by itself mean despair. Ask what the person values, what social opportunities were available and how the present illness has altered the developmental task.

2.7 Alignment with Piaget, Freud, Kohlberg, Mahler and Vygotsky

The chapter's theory-alignment plate should be read horizontally, with one question attached to each theorist. Piaget asks how the structure of cognition changes. His sensorimotor account concerns knowledge through action and the emergence of representation; it does not explain trust. Freud asks how instinctual conflict is organised across the psychosexual stages; temporal overlap with an Eriksonian stage does not make autonomy equivalent to the anal stage or identity equivalent to the genital stage.

Kohlberg asks how reasons for moral decisions are structured. Kaplan and Sadock describe preconventional morality before adolescence, conventional response levels during and later in adolescence, and a postconventional response level in late adolescence or early adulthood. Moral development concerns the justification of right action, whereas Eriksonian identity concerns continuity, values and commitment. A sophisticated moral argument does not prove identity achievement, and social conformity does not by itself establish conventional moral reasoning.

Mahler asks how psychological separateness emerges within the caregiver relationship. Her separation-individuation theory includes a hatching period described by Kaplan and Sadock and Wolberg from about 4 or 5 months to 10 months. Wolberg places rapprochement onset between 15 and 18 months, the rapprochement crisis around 18 to 24 months, and libidinal object constancy during the third year of life. These are source descriptions, not screening thresholds. Mahler's individuation is not the same construct as Erikson's autonomy, although both attend to emerging separateness.

Vygotsky asks what development becomes possible through language, cultural tools and skilled collaboration. The zone of proximal development is the difference between independent performance and performance achieved with appropriate support. Erikson identifies the psychosocial meaning of mastery; Vygotsky helps the clinician observe how guided participation changes actual performance. A child who appears inferior in an unsupported task may reveal strong learning potential when the task is scaffolded.

The safest comparison formula in a viva is *same period, different explanatory target*. The theories can be placed beside one another for recall, but they cannot be added into a single universal clock. Their stage ages are descriptive conventions, their units of analysis differ, and none of the theoretical bands is a developmental referral threshold.

■ **TRAP** **Do not diagnose an "unresolved Erikson stage."** The conflicts are formulation constructs, not ICD diagnoses, psychometric scores or developmental-screening results. Avoid retrospective certainty, such as attributing adult symptoms to one caregiving event, and avoid reading age bands as deadlines.

2.8 Clinical formulation: strengths, limitations and cultural use

Erikson is clinically useful because he joins symptom, biography and social context. It prompts the psychiatrist to ask what developmental task the illness interrupts, what virtue remains available as a strength, and what social response is amplifying the conflict. Depression during vocational transition may be experienced through identity and competence; mania may destabilise commitments and intimacy; chronic psychosis may obstruct industry or generativity; caregiving and retirement may transform the meaning of contribution. These formulations enrich management when they remain hypotheses grounded in the person's narrative.

The model also has limits. Its broad concepts are difficult to operationalise, stage boundaries are not sharply defined, and development is more variable and recursive than a strict sequence suggests. The classical account can overprivilege individual autonomy, stable occupational identity and a linear adult life course. Collective identities, caste, class, gendered opportunity, disability, migration, chosen family and economic constraint may shape which resolutions are possible and valued. Indian family interdependence should not be mislabelled dependency merely because it differs from an individualist ideal.

Use **goodness-of-fit** to prevent moralising. **Temperament** refers to constitutionally influenced differences in reactivity and regulation; goodness-of-fit refers to the compatibility between those characteristics and environmental expectations. A cautious child in a patient environment may

develop competence, while the same child under ridicule and forced performance may experience inferiority. The temperament is not the Eriksonian crisis, but the fit can alter how the crisis is negotiated.

A sound formulation states the evidence and its limits: "The current transition appears to reactivate concerns about competence and identity" is defensible; "industry was not resolved, therefore this disorder developed" is not. Erikson should direct questions, identify strengths and organise meaning. It should not be used to blame parents, prescribe a culturally narrow life path or explain away neurodevelopmental, psychiatric and social causes.

2.9 Five worked cases: from recall to application

Worked case 1 - milestone-dating and red-flag reasoning. A child at 24 months of age is described as energetic, insistent on self-feeding and easily frustrated by adult help. The child does not use spontaneous two-word phrases. The Eriksonian reading is autonomy versus shame and doubt: the self-directed behaviour is developmentally meaningful and should be met with bounded choices rather than ridicule. The language finding is a separate red-flag: absence of spontaneous two-word phrases at 24 months of age warrants further developmental evaluation. Do not use "the autonomy stage" to normalise the language concern, and do not infer global delay from a single finding. Assess all developmental domains, hearing, regression and context.

Worked case 2 - attachment-pattern identification. In Ainsworth's Strange Situation, a young child explores while the caregiver is available, becomes distressed on separation, seeks contact on reunion and settles sufficiently to resume exploration. This pattern is most consistent with secure attachment because the caregiver functions as a secure base and haven. The Eriksonian link is supportive, not diagnostic: reliable care may contribute to basic trust, but secure attachment and trust are different constructs. One observation cannot establish the whole caregiving history or prove that an Eriksonian crisis is resolved.

Worked case 3 - temperament and goodness-of-fit. A child within Berk's 6 to 11 years industry-versus-inferiority mapping is slow to approach unfamiliar classroom tasks but works carefully after a demonstration. Repeated public comparison has led the child to say, "I am bad at everything." The theory-application is industry versus inferiority, but the mechanism is not laziness. A cautious temperament is meeting a poor goodness-of-fit. Assess learning, attention, sensory and emotional contributors; provide graduated demonstration, private feedback and tasks that permit observable mastery. The goal is evidence-based competence, not empty reassurance.

Worked case 4 - adolescent identity and risk-taking. An adolescent alternates between career plans, adopts different styles with different peers and takes a hazardous risk in a highly charged group setting, while reasoning thoughtfully about the same risk in clinic. Exploration without settled commitment may fit moratorium rather than pathology. The context shift is compatible with a dual-systems or maturational imbalance account of risk-taking, but it does not excuse harm or prove immaturity. Assess mood elevation, substance use, coercion, self-harm, sleep,

consequences and capacity. Explore identity formation and values without treating gender identity, appearance or nonconformity as role confusion.

Worked case 5 - adult theory-application. An older adult after bereavement reviews a life containing valued work, estrangement and choices now regretted, yet can hold gratitude and sorrow together and remains engaged in mentoring. The dominant Eriksonian formulation is integrity rather than despair, with generativity still active through mentoring. Integrity does not require a flawless or regret-free biography. If pervasive hopelessness, anhedonia, guilt or suicidal thinking is present, assess depressive illness directly; a developmental formulation must never replace risk assessment or diagnosis.

Exam synthesis. In any vignette, first identify the actual phenomenon: a developmental attainment, warning sign, relationship pattern, temperament-environment fit, physical pubertal stage, form of reasoning or psychosocial conflict. Then name the correct framework and state what it can and cannot infer. That separation prevents the commonest error in developmental psychiatry: using an elegant theory as if it were a measurement instrument.

3 . Freud psychosexual stages

Freud's psychosexual theory proposes that personality develops through a sequence in which **libido**, understood as psychic energy associated with the sexual drives, is organised predominantly around a different bodily zone and a different developmental conflict. In examination answers, "sexual" must be used in Freud's broad drive-theory sense rather than reduced to adult genital intercourse. The sequence is **oral, anal, phallic, latency and genital**. Each phase leaves residues in later personality, while excessive frustration or gratification may produce **fixation**; under stress, functioning may move back toward an earlier mode through **regression**. This is a historical psychoanalytic model of personality development, not a contemporary milestone schedule and not a method for diagnosing developmental delay.

How to use this page. First memorise the order, age brackets, principal zone and central conflict. Then add the characteristic mode of gratification, the proposed resolution and the classical fixation themes. Finally, separate three things that candidates commonly collapse: a psychosexual phase is not a developmental milestone, Freud's genital phase is not a Tanner stage, and a psychodynamic formulation is not proof that an early event caused the patient's current disorder.

Must memorise: oral, anal, phallic, latency, genital; the drive focus moves from incorporation, through control and oedipal identification, to relative quiescence and then mature genital organisation.

3.1 The organising model and the five-stage sequence

A changing drive focus, a phase-specific conflict and a possible residue. Freud did not describe five watertight compartments. Earlier modes of pleasure and relating remain available after a later organisation becomes dominant. A stage is therefore best learnt through four linked questions:

- 1. *Where is gratification centred?*
- 2. *What conflict must be negotiated?*
- 3. *How is it resolved?*
- 4. *What pattern did Freud associate with fixation?*

The theory assumes that constitutional drives interact with caregiving and prohibition. The child's ego gradually develops ways of regulating these pressures, and identification contributes to internal structure, especially during the phallic phase.

Source-reported teaching brackets, not referral thresholds:

STAGE	BERK'S DEVELOPMENT THROUGH THE LIFESPAN	TASMAN PSYCHIATRY
Oral	Birth to 1 year	birth to 12-18 months
Anal	1-3 years	12-18 to 36 months
Phallic	3-6 years	3-5 years
Latency	6-11 years	
Genital	Adolescence	11 years plus for the adolescent genital phase

Birth to 1 year ORAL - BERK	1-3 years ANAL - BERK	3-6 years PHALLIC - BERK	6-11 years LATENCY - BERK
--------------------------------	--------------------------	-----------------------------	------------------------------

The sources state these ages without defining their normative construct. They should therefore be reproduced as the sources' approximate theoretical brackets, never converted into a median, an upper limit of normal or a when to refer threshold. Their overlap also warns against false precision. A child does not become abnormal because a psychodynamic theme appears outside its conventional bracket.

3.2 Oral stage: incorporation, dependence and the first relation to an object

The mouth is the principal zone, and feeding is both gratification and relationship. In the oral stage, sucking, swallowing and biting organise pleasure. The breast or feeding caregiver is not merely a source of nutrition in the theory; it is the first major object, meaning the person or representation toward whom a drive is directed. Satisfaction involves taking in, while delay and absence introduce frustration. The central developmental themes are receiving, incorporation, dependence and the beginnings of tolerating separation from immediate gratification.

Classical accounts distinguish an earlier oral receptive or incorporative mode from a later biting, aggressive mode. Freud and later psychoanalytic writers linked oral fixation with dependency, passivity, demandingness, gullibility or incorporative habits, and linked the biting mode with verbal aggression and hostility. These are theoretical associations, not validated one-to-one predictions. Feeding difficulty, smoking, overeating or dependency in an adult cannot by itself

demonstrate oral fixation, and a resident should never retrofit an infancy narrative merely because an adult symptom looks symbolically oral.

3.3 Anal stage: control, autonomy and the social demand for regulation

Pleasure in retention and expulsion meets the caregiver's demand for bowel control. In the **anal stage**, elimination becomes the principal bodily focus. Toilet training gives the child an arena in which bodily pleasure, autonomy, obedience, defiance, praise and shame can become linked. The theoretically important issue is not simply continence; it is the negotiation between the child's wish to control what is given or withheld and the environment's demand that discharge occur at an acceptable place and time.

Classical descriptions contrast anal-retentive themes, such as excessive orderliness, obstinacy, parsimony and control, with anal-expulsive themes, such as messiness, defiance and poorly regulated discharge. The familiar triad of orderliness, obstinacy and parsimony is a high-yield psychoanalytic association, but it is not a diagnostic triad. Contemporary formulation should ask how control, shame, autonomy and caregiver responses were experienced, while avoiding blame directed at toilet training or at the caregiver.

3.4 Phallic stage: oedipal conflict, identification and superego development

Genital awareness is organised around triangular attachment, rivalry and identification. The **phallic stage** centres on the phallus in Freud's theory and introduces the **Oedipus complex**: desire for the opposite-sex parent, rivalry with the same-sex parent and anxiety about the consequences of that rivalry. In Freud's account of the boy, castration anxiety promotes renunciation of the oedipal wish and identification with the father. This identification contributes to the consolidation of the superego and to the internalisation of prohibitions and ideals.

Freud's account of female development invoked penis envy and a different route through oedipal attachment. It has been criticised as androcentric, anatomically reductionist and based on male development as the norm. The term Electra complex is associated with Jung, not Freud's preferred terminology. In an examination, state the classical proposition accurately, then state the criticism. Resolution is described through relinquishment of the incestuous wish, identification and movement toward relationships outside the original triangle. Fixation has been associated with vanity, exhibitionism, rivalry, sexual inhibition or reckless assertion, but again these are hypotheses within a theory, not diagnostic findings.

■ **TRAP** **Phallic is not genital, and genital is not Tanner.** The phallic phase concerns oedipal organisation in early childhood. The later genital phase concerns the mature organisation of drives. Tanner staging describes observable pubertal anatomy and is a different construct altogether. Also remember that Jung, not Freud, popularised "Electra complex" as a label.

3.5 Latency: relative drive quiescence, sublimation and social learning

Latency is a relative reduction in overt psychosexual preoccupation, not the disappearance of libido. During the **latency stage**, oedipal wishes are held by repression and other defences, while energy is increasingly channelled into learning, peer relationships, play, competence and

culturally valued activity. **Sublimation** is the key mechanism: drive energy is redirected toward aims that are socially acceptable and psychologically productive.

Latency is sometimes described as an interval rather than a new erogenous-zone organisation, but it remains part of the five-part examination sequence. Its clinical themes include the capacity to learn, work with peers, follow rules and invest in interests outside the family triangle. It should not be mistaken for a period without sexuality, fantasy or conflict. The word *relative* protects the answer from that overstatement.

3.6 Genital stage: integration, reciprocity and mature object love

Adolescent drive reactivation is reorganised toward reciprocal relationships. In the **genital stage**, earlier partial drives are theoretically integrated under genital primacy. The developmental aim is not mere sexual activity. It is the capacity for reciprocal intimacy, concern for the other person, productive work and a balance between love and responsibility. Mature gratification now depends on recognising the object as a separate person rather than only as a source of need satisfaction.

Unresolved earlier conflicts may reappear when adolescent drives intensify, so the final phase includes both reworking and integration. The phrase *mature genital character* belongs to Freud's normative ideal and should not be treated as a contemporary measure of mental health. Sexual orientation, celibacy, disability, cultural restraint or absence of partnered sexual activity does not indicate failed genital development. This is another reason to keep the historical theory separate from moral judgement and from anatomical puberty.

3.7 Fixation and regression: the mechanisms that connect stages to adult formulation

Fixation is the persistence of a disproportionate libidinal investment in an earlier mode of gratification or conflict. Classical theory attributes it to marked frustration, excessive gratification or difficulty resolving the phase-specific conflict. Regression is a return, especially under stress, to a developmentally earlier organisation of wishes, defences, relationships or behaviour. Fixation describes a relatively enduring vulnerability; regression describes movement toward that point of vulnerability. They are related but not synonyms.

The clinically useful move is thematic and probabilistic. Dependency and taking-in may suggest an oral theme; control, withholding and shame may suggest an anal theme; rivalry, display and triangular relationships may suggest a phallic theme. The clinician then tests the hypothesis against the patient's history, current relationships, transference, defences and culture. The clinically unsafe move is deterministic: inferring a specific childhood event from an adult trait, or explaining a disorder solely by naming a stage.

■ **RULE** A psychosexual formulation generates hypotheses; it does not establish causation. Use stage language to organise recurring themes in gratification, control, rivalry, identification and intimacy. Do not use it as a developmental test, a diagnosis or proof of what happened in childhood.

3.8 Alignment with other developmental theories: same age axis, different constructs

The companion synthesis map aligns the major theories by age, but alignment means temporal overlap, not conceptual equivalence. Freud maps changing drive organisation through the psychosexual stages. Piaget maps cognitive structure, beginning with the sensorimotor period and moving through preoperational, concrete operational and formal operational thought. Erikson maps relational and social crises through the psychosocial stages. Thus an early-childhood conflict may simultaneously be described as anal in Freud, autonomy versus shame and doubt in Erikson, and sensorimotor or early preoperational in Piaget, without the terms meaning the same thing.

Kohlberg maps moral development from preconventional through conventional to postconventional reasoning. His sourced age relationships are descriptive bands such as before adolescence, during adolescence and later in adolescence, not numeric age ranges; adding numbers would manufacture precision. Mahler maps separation-individuation, including hatching, practising, rapprochement and movement toward object constancy. Vygotsky offers no stage sequence: the zone of proximal development, scaffolding, private speech and socially mediated learning are cross-cutting mechanisms. Vygotsky therefore belongs beside the age plate, not inside it as invented stages.

The clinical use of the alignment is to ask several different questions about the same child: how the child thinks, what social conflict is being negotiated, how drives and relationships are organised, how morality is reasoned, how separation is managed and what learning becomes possible with guidance. The plate prevents tunnel vision; it does not turn six theories into one grand developmental schedule.

3.9 Clinical value, cultural limits and contemporary critique

Freud's model remains important because it introduced developmental continuity, unconscious conflict, infantile sexuality, defence, symbolic meaning and the possibility that early relationship patterns recur in adult life. In psychodynamic assessment it can sharpen attention to how a patient receives care, manages dependency, uses control, enters triangles, responds to prohibition and negotiates intimacy. These themes can be observed in narrative, dreams, symptoms, transference and countertransference, then integrated with attachment, temperament, trauma, learning and the current social environment.

Its limitations are equally examinable. The theory was derived largely from retrospective clinical reconstruction rather than prospective developmental research; several central propositions are difficult to falsify; the account of female development is androcentric; the model can be heteronormative and overly centred on sexuality; and stage-to-trait links are vulnerable to circular explanation. Culture changes the meanings of feeding, sleeping, toilet training, bodily privacy, family hierarchy and adolescent intimacy. For Indian practice, joint-family structures and family-mediated caregiving may alter the relational expression of dependence or autonomy without indicating pathology. Use the theory as one formulation lens, not as a universal cultural ruler.

INDIA

The Freud brackets above come from general developmental and psychiatry texts and are not **Indian norms**. No verified India-specific psychosexual-stage age adaptation is supplied here, so none should be invented. Indian developmental instruments answer a different question: the **Trivandrum Development Screening Chart** screens development from **0 to 6 years** using 51 items, with one item delay constituting a positive screen, while **DASII** yields Motor DQ and Mental DQ, and a DQ below **85** is used as a delay marker in Indian follow-up practice. Neither instrument validates oral, anal or phallic fixation.

3.10 Spotting deviation from normal: developmental surveillance is a separate clinical task
Psychosexual stages do not define developmental red flags. Developmental surveillance is longitudinal observation of developmental progress, caregiver concerns and function, supported by milestone-based screening when indicated. It is the correct framework for spotting deviation, warning signs and deciding when to refer. Calling a child "fixated" does none of these jobs and may delay proper assessment.

High-yield developmental red flags include the following:

REFERRAL SIGNAL	AGE
Absent babbling	at 12 months .
Absent single words	at 16 months .
Absent spontaneous two-word phrases	at 24 months .
Loss of language or social skills	at any age.

These are referral signals from *Developmental Psychopathology*, not Freud-derived ages. For gross motor development, the **WHO norms** are achievement windows bounded by the 1st and 99th centiles and are intended to signal the need for appropriate screening when a child appears late; they are screening signals, not diagnostic thresholds. India was one of the WHO study sites. Use the companion milestone plate for the individual windows rather than recreating it here.

The apparent Western-versus-Indian comparison must also be handled honestly. Freud's source brackets are theoretical ages, not **Western norms**, while the Trivandrum chart and DASII are Indian screening or assessment instruments, not psychosexual-stage adaptations. There is therefore no like-for-like set of **Indian norms** to place beside Freud's ages. Record genuine differences when two sources measure the same construct; when they do not, name the mismatch instead of forcing an alignment.

3.11 Consolidation: mnemonic, active recall and reconstruction

MNEMONIC

"Old Analysts Prefer Long Games"

O-A-P-L-G gives Oral, Anal, Phallic, Latency, Genital. Attach one organising word to each: *incorporation, control, identification, sublimation, reciprocity*. The mnemonic fixes order; the five organising words recover the theory.

Redraw from memory. Reconstruct the companion **synthesis map** with **theories by age** on separate horizontal lanes, place Freud's five stages on its lane, keep Kohlberg's bands descriptive, and draw Vygotsky as a cross-cutting mechanism. Then add the companion **milestone anchors** and developmental red flags on a separate clinical lane. Keeping the theory lanes separate from the surveillance lane is the final examination safeguard.

4 . Moral development (Kohlberg)

Lawrence Kohlberg's theory of moral development describes development in the *reasoning used to justify a moral decision*. It is not chiefly a theory of whether a person gives the examiner the approved answer, behaves well, obeys the law, or possesses a particular religious belief. Kohlberg extended Piaget's interest in the development of moral judgement and proposed an invariant sequence of increasingly differentiated ways of answering a central question: *why is an action right or wrong?* The sequence contains **three** levels, each containing **two** stages. The centre of moral authority shifts from externally imposed consequences, to interpersonal and social expectations, and finally to principles that can be used to evaluate the social order itself.

How to use this page. First fix the architecture: **preconventional**, **conventional**, and **postconventional**; then learn the contrast within each pair of stages. Stage 1 avoids punishment, stage 2 pursues exchange; stage 3 seeks interpersonal approval, stage 4 maintains the social system; stage 5 tests rules against rights and social contract, stage 6 reasons from universal ethical principles. In a dilemma, always identify the justification rather than the chosen act. Timing must remain descriptive: Kaplan and Sadock relates the stages to bands such as "school-aged children" and "during adolescence", but supplies no numeric ages for Kohlberg.

Must remember: Kohlberg stages the *reason for the judgement*, not the judgement itself.

MNEMONIC

"Punishment - Exchange - Approval - Law - Contract - Principle"

The sequence is punishment, exchange, approval, law, social contract, universal principle.




Fig 20.4 Kohlberg's three levels and six ordered stages, with what a moral dilemma scores. The timing column quotes Kaplan and Sadock's descriptive bands verbatim; that source gives no numeric age for any level or stage, so the plate has no age axis and nothing on it is drawn to scale. A level-wide band is sourced for the preconventional level only, and stage 6 has no band at all, so its cell is left empty rather than inferred.

4.1 Method and architecture: three levels, six ordered stages

A stage is a structure of reasoning, not a list of moral opinions. Kohlberg used hypothetical moral dilemmas, classically the Heinz dilemma, and questioned the respondent about the decision and its justification. In that dilemma, a person must consider whether a husband should steal an unaffordable medicine to save his wife. Saying "steal" or "do not steal" does not by itself locate a stage. A respondent might oppose theft because of punishment, because it is a poor bargain, because respectable people disapprove, because law preserves society, or because a law may be overridden by a more fundamental right. Conversely, respondents at different stages can endorse the same action for structurally different reasons.

3 LEVELS	6 ORDERED STAGES	2 STAGES PER LEVEL
-------------	---------------------	-----------------------

The model is cognitive-developmental. Stages are proposed to occur in sequence, with later reasoning reorganising rather than merely adding to earlier reasoning. Movement depends on

cognitive development, perspective-taking and encounters with moral conflict. A person may show a dominant mode with adjacent-stage reasoning rather than perfectly pure performance in every situation. Developmental advance also does not guarantee moral conduct: behaviour is additionally shaped by emotion, impulse control, identification, attachment, reward, coercion, group norms and opportunity.

■ **RULE** **Score the justification, not the verdict.** Two people may make the same choice at different stages, and opposite choices may be supported by reasoning at the same stage.

4.2 Preconventional morality: consequences and personal interest

At the preconventional level, rules are experienced as external to the self. The child recognises authority and social rules, but moral judgement is organised around concrete consequences and personal advantage rather than around preserving a shared social system. Kaplan and Sadock describes this level as occurring **before adolescence**. This is a descriptive relationship, not a cutoff, expected range, or referral threshold.

Stage 1: punishment-and-obedience orientation. An act is wrong because authority prohibits it and punishment follows. Consequences are taken as evidence of wrongness, and the more severe the consequence, the worse the act may appear. Obedience is heteronomous: the rule comes from a powerful other, and the perspectives or intentions of those involved are poorly coordinated. Kaplan and Sadock associates this response level with **school-aged children**.

Stage 2: instrumental purpose, individualism and exchange. The person now recognises that different individuals have different interests. What is right serves one's needs and, where useful, the needs of others through a concrete exchange: "I will help you if you help me." Fairness is therefore understood as equal exchange or reciprocity, not yet as loyalty, social duty or impartial principle. Kaplan and Sadock notes this response level in **some young adolescents**. The phrase "some" matters: it must not be converted into a universal age norm.

4.3 Conventional morality: relationship approval and maintenance of society

At the conventional level, the reference point expands from the actor's immediate consequences to membership in relationships and society. The person values expectations, roles, rules and institutions because these sustain interpersonal trust and an ordered social world. "Conventional" here means conformity to a recognised social order, not shallow thinking.

Stage 3: mutual interpersonal expectations, relationships and conformity. The morally good person is kind, loyal, trustworthy and approved by significant others. Intentions become important, and the respondent can take the viewpoint of another person, but judgement remains tied to the expectations of close relationships and the wish to be seen as a "good" person. Kaplan and Sadock places this response level **during adolescence**.

Stage 4: social-system and conscience orientation. The frame widens from close relationships to society as an organised whole. Right conduct means performing one's duty, respecting legitimate authority and maintaining laws and institutions because social cooperation would otherwise break down. This is not simply fear of punishment: stage 1 obeys power to avoid

consequences, whereas stage 4 endorses law as necessary for the system. Kaplan and Sadock relates this response level to **later in adolescence**.

4.4 Postconventional morality: evaluating rules by principles

At the postconventional level, a person can stand back from a particular society's rules and ask whether they are justifiable. Laws remain important, but their legitimacy depends on values and principles that rational persons could endorse. The crucial move is from being governed by the social order to being able to evaluate it. Postconventional reasoning should not be confused with reflexive rebellion, personal preference, or disregard for law.

Stage 5: prior rights and social-contract orientation. Rules are understood as human arrangements designed to protect welfare, liberty and rights. Because arrangements are socially created, they may be changed by agreed procedures when they fail those purposes. Some values and rights retain priority even when a majority prefers otherwise. Kaplan and Sadock associates this response level with **late adolescence or early adulthood**; this wording is descriptive and does not imply that everyone reaches the stage.

Stage 6: universal ethical-principle orientation. Moral judgement is guided by self-chosen, generalisable principles of justice, equality and respect for human dignity. A valid principle applies consistently across persons, including oneself, and conscience follows the principle when ordinary law conflicts with it. Stage 6 differs from stage 5 in its source of obligation: stage 5 emphasises a fair social contract and basic rights, whereas stage 6 emphasises universal principles chosen through autonomous moral reasoning.

4.5 The sequence as a set of contrasts

The fastest way to retrieve the stages in an examination is to ask where moral authority sits. At stage 1 it sits with the punishing authority; at stage 2 with the individual's concrete interest and negotiated return; at stage 3 with valued relationships and approval; at stage 4 with the duties of the social system; at stage 5 with rights and procedures that legitimate the system; and at stage 6 with universal ethical principle. Three contrasts prevent most errors.

CONTRAST	DISTINCTION
Punishment versus law:	both may support obedience, but one avoids a consequence while the other preserves social order.
Approval versus contract:	one asks what a good person in the relationship would do, while the other asks what free and equal persons could justify.
Contract versus principle:	one revises law through shared procedures, while the other tests every law and procedure against a generalisable ethical commitment.

■ **TRAP** **Do not turn descriptive bands into numeric ages.** "School-aged children", "some young adolescents", "during adolescence", "later in adolescence", and "late adolescence or early adulthood" are Kaplan and Sadock's formulations. Kohlberg has no sourced numeric age row for the theory-alignment plate.

4.6 What the theory explains, and what it does not

Kohlberg explains the organisation of moral judgement. It does not provide a complete psychology of morality. Knowing what is right does not ensure doing it, and sophisticated reasoning can be used after the fact to rationalise conduct. Empathy, guilt, shame, temperament, attachment, modelling, cultural learning, power, material constraint and the immediate social context can all influence behaviour without being reducible to a reasoning stage. Nor should a stage label be mistaken for intelligence, diagnosis, legal competence, criminal responsibility or overall maturity.

The approach nevertheless made important contributions. It treated children as active constructors of moral meaning rather than passive recipients of rules; separated the structure of a reason from its surface content; and supplied a developmental account of perspective-taking, authority, reciprocity, obligation and justice. Its structured dilemmas also made moral reasoning open to systematic interview and longitudinal study.

4.7 Criticisms and later refinements

Justice is not the whole of morality. Carol Gilligan argued that an emphasis on rights, rules and abstract justice can undervalue an ethic of care centred on relationship, responsibility and response to need. The useful examination answer is not that one sex reasons by justice and another by care as a fixed biological divide. The stronger criticism is that the scoring framework privileged one moral voice and that care and justice may both be available across persons and contexts.

Culture shapes the meaning of mature reasoning. The proposed order has received support across settings, but the claim that abstract individual rights represent the highest endpoint can favour liberal, individualistic assumptions. Community, sacred obligation, family duty and relational responsibility may be morally elaborated rather than merely conventional. **Dilemma performance is also context-sensitive.** Hypothetical answers may not predict action under fear, loyalty, intoxication, peer pressure or personal loss; education and verbal ability influence how reasons are expressed; and the same person may reason differently across dilemmas. Finally, stage boundaries may be less discrete than a strict staircase suggests, with mixed and adjacent-stage responses occurring in practice.

4.8 Clinical use and relation to the other developmental theories

Use the framework to formulate reasoning, never to stamp a child with a stage. In a child or adolescent assessment, a clinician can ask how the patient understood a conflict involving rules, peers, harm or fairness, then explore intentions, consequences, perspective-taking and the perceived source of obligation. A shift from "I will be punished" to "people will be harmed" or "the rule protects everyone" adds clinically useful information about social cognition and judgement. Interpretation must remain contextual and should be integrated with cognitive ability, language, family values, school experience, neurodevelopment, affect regulation and actual behaviour. It is not a diagnostic test and should not be used alone for a forensic conclusion.

CLINICAL USE

After a recurrent rule-breaking episode, ask the patient to explain why the act was acceptable or unacceptable, what each person believed, and whether the rule would remain fair if roles were reversed. The purpose is to sample perspective-taking and the structure of moral reasoning, not to demand a socially approved answer. A rehearsed answer without corresponding conduct is still clinically informative: it shows the gap between moral cognition and moral action.

Use the accompanying theory-alignment plate as a map of different questions, not as proof that the theories describe the same process.

- Piaget, from the sensorimotor period onward, maps changes in cognitive structure.
- Erikson maps psychosocial stages and their central conflicts.
- Freud maps psychosexual stages.
- Mahler describes separation-individuation.
- Kohlberg maps moral reasoning.

Vygotsky should sit beside, not inside, that stage sequence: sociocultural mediation, private speech and the zone of proximal development are cross-cutting mechanisms, not age-bound stages. Kohlberg likewise appears on the plate only in Kaplan and Sadock's descriptive bands, never on an invented numeric age axis.

5 . Mahler separation-individuation theory

Margaret Mahler's separation-individuation theory is a psychoanalytic account of how the infant moves from an experienced psychological unity with the mother toward a stable sense of being a separate person who can remain emotionally connected to her. Its subject is the *psychological birth* of the individual, not biological birth and not the attainment of a motor milestone. **Two** developmental lines proceed together but are not identical: **separation** is the emergence from symbiotic fusion and the differentiation of self from object, while **individuation** is the acquisition of a distinctive identity, autonomous ego functions and an enduring inner sense of self.



■ Normal symbiosis

■ Differentiation

■ Practising

■ Rapprochement

■ Object constancy

Fig 20.5 Mahler's account of psychological birth. Separation and individuation proceed together through differentiation, practising and rapprochement toward a stable inner representation of the caregiver and a distinct sense of self.

How to use this page. Fix the distinction between separation and individuation, then learn the sequence: the historical precursor phases of normal autism and normal symbiosis, followed by differentiation or hatching, practising, rapprochement, and consolidation toward libidinal object constancy. For each phase, ask what changes in self-object boundaries, exploration, use of the caregiver and mental representation. Finally, retain the limits: this is an object-relations formulation, not a developmental screening schedule, and its age brackets must not be converted into referral thresholds.

Must memorise: separation means "I am not you"; individuation means "I am me"; their sequence is hatching, practising, rapprochement and consolidation toward object constancy.

MNEMONIC

"Hatching - practising - rapprochement - consolidation"

separation means "I am not you"; individuation means "I am me"; their sequence is hatching, practising, rapprochement and consolidation toward object constancy.

5.1 The organising model: psychological birth through two intertwined lines

The child becomes separate without becoming unrelated. Mahler's model belongs to object-relations theory, where an *object* is a significant person as experienced and mentally represented, not an inanimate thing. Development requires a gradual reorganisation of the infant-mother relationship. Perceptual differentiation, locomotion and autonomous ego capacities support separation; the consolidation of identity and internal representations supports individuation. The endpoint is therefore neither fusion nor detachment. It is the capacity to function as a distinct self while maintaining a dependable emotional bond with a whole caregiver.

Mahler developed the theory through observations of young children and their mothers. In the classical vocabulary, *mother* denotes the central maternal object. In contemporary clinical use, *primary caregiver* is often the more accurate functional term, while the original terminology should still be named in an examination answer. The theory focuses on intrapsychic organisation inferred from behaviour within a relationship; it does not claim that visible independence alone proves psychological individuation.

Source-reported theoretical timings, not normative cut-offs:

TIMING	STAGE	SOURCE
The first weeks of the infant's life	normal autistic phase	in Wolberg's <i>Technique of Psychotherapy</i>
Three to four weeks	normal symbiotic phase	as reported by Wolberg
About 4 or 5 months to 10 months	hatching phase	in Wolberg
Between 15 and 18 months	onset of rapprochement	in Wolberg
15-24 months	is the rapprochement phase	in <i>Cognitive Therapy of Personality Disorders</i>
Around 18 to 24 months of age	rapprochement crisis	in Wolberg
During the third year of life	libidinal object constancy	in Wolberg

The first weeks of the infant's life NORMAL AUTISTIC PHASE	About 4 or 5 months to 10 months HATCHING PHASE	15-24 months RAPPROCHEMENT PHASE	During the third year of life LIBIDINAL OBJECT CONSTANCY
---	--	-------------------------------------	---

These sources state theoretical ages without identifying them as medians, expected ranges, upper limits or red-flag thresholds. They must therefore be repeated only as source-reported stage

timings. Overlap or variation between texts is not evidence of abnormal development, and none of these values tells a clinician when to refer a child.

-
- **RULE** **Mahler's ages locate concepts within a theory; they do not define developmental normality.**
Use developmental surveillance and validated screening instruments to detect delay. Do not diagnose a failed phase, fixation or developmental disorder from a child's calendar age.
-

5.2 Precursor phases: normal autism and normal symbiosis

Mahler begins with states in which self and other are not yet represented as clearly separate.

In the historical **normal autistic phase**, the infant was described as predominantly invested in physiological regulation and protected from external stimulation by a relatively closed psychological barrier. The term *autistic* here belongs to Mahler's early psychoanalytic vocabulary. It does not mean autism spectrum disorder, does not predict it and should never be entered into a modern case record as though it were a diagnosis.

In the **normal symbiotic phase**, the infant is increasingly responsive to the mothering environment but is theorised to experience infant and mother as a dual unity within a common boundary. Symbiosis in this model is an intrapsychic state of undifferentiation, not merely closeness, dependence or frequent physical contact. The caregiver regulates tension and supplies gratification, while repeated experiences begin to organise an expectation of relief and relationship. Normal symbiosis is a precursor to differentiation; persistent adult dependency cannot simply be labelled proof that the person is "stuck" in it.

5.3 Differentiation or hatching: the beginning of separation

Attention turns outward and the infant begins to inspect the boundary between familiar self-object unity and the wider world. The **differentiation subphase**, also called hatching, marks the proper beginning of separation-individuation. The metaphor is psychological, not anatomical. The infant becomes more alert to the environment, studies the mother's face and body, compares familiar and unfamiliar people, and begins to move physically away while checking back perceptually. These behaviours are read as signs of emerging differentiation rather than complete selfhood.

The clinical emphasis is on discrimination and boundary formation. Familiarity becomes more organised, the external world acquires greater salience and the mother is increasingly apprehended as a particular object rather than merely part of a need-satisfying unit. Yet separation remains supported by the caregiver's availability. Hatching is therefore not abrupt detachment. It is a gradual opening of a relationship in which difference can be noticed without the bond disappearing.

5.4 Practising: locomotion, exploration and the exhilaration of autonomy

New autonomous capacities widen the child's psychological and physical world. In the **practising subphase**, locomotion and other developing ego functions permit active exploration away from the caregiver. Mahler described an exuberant investment in the wider world, with the child appearing temporarily absorbed in newly acquired competence and relative independence.

The caregiver remains the secure emotional centre. The child returns for brief contact or emotional refuelling, then ventures outward again.

The high-yield paradox is that apparent independence still rests on dependable availability. Exploration does not prove that the object is no longer needed; it shows that connection can support movement away. Clinically, this helps the resident observe how autonomy and dependence coexist rather than forcing behaviour into either "independent" or "dependent". Motor progress may facilitate practising, but Mahler's phase is a relational and intrapsychic construct, not a motor-development category.

5.5 Rapprochement: separateness becomes emotionally costly

The child now recognises separateness more fully and seeks renewed emotional closeness without surrendering autonomy. In the **rapprochement subphase**, the earlier exhilaration of practising gives way to a sharper awareness that the caregiver is a separate person who cannot be controlled by omnipotent wish. The child brings discoveries back to share, follows the caregiver, demands participation and alternates between bids for help and assertions of independence. Mahler described this push-pull as **ambitendency**: a simultaneous wish to move away and to be close.

The rapprochement crisis is the intensification of that conflict. Dependence may feel necessary, while closeness may threaten the new autonomy. Frustration exposes the fact that the loved object also disappoints, so affectionate and angry experiences must increasingly be held together rather than kept apart. A sufficiently available caregiver neither abandons the child nor erases every frustration. She permits approach, tolerates rejection and survives ambivalence, helping the child discover that separateness need not mean loss.

5.6 Consolidation and libidinal object constancy

A stable inner representation permits connection across absence, frustration and mixed feeling. During consolidation, individuation advances and self-other boundaries become more durable. **Libidinal object constancy** means that a predominantly positive, emotionally invested representation of the caregiver can be maintained when she is absent, unavailable or temporarily experienced as frustrating. Love and anger can be directed toward the same whole person without that person having to become entirely good or entirely bad.

Object constancy depends on integrating affectively contrasting representations of self and object. It supports tolerance of delay, separation and ordinary ambivalence, while stabilising identity and relationships. It is an achievement with degrees, not an all-or-none certificate permanently acquired at a single moment. Stress can still strain the capacity to hold a complex view of self and others.

■ **TRAP** **Object constancy is not Piagetian object permanence.** Object permanence is the cognitive understanding that an object continues to exist when not perceived. Mahler's object constancy is affective and representational: the relationship to a whole loved person remains internally available across absence and frustration. Also, Mahler's "normal autistic phase" is not autism spectrum disorder.

5.7 Clinical formulation: useful relational questions, not developmental verdicts

Mahler gives the psychiatrist a vocabulary for recurrent tensions between closeness and autonomy. In assessment, ask:

- whether separation evokes abandonment;
- whether intimacy evokes engulfment;
- whether the patient alternates between clinging and withdrawal;
- whether disappointment rapidly converts an admired person into a hated one;
- whether an absent attachment figure can remain emotionally real.

Similar patterns may appear in relationships, the therapeutic frame, transference and countertransference.

The model has been influential in psychodynamic formulations of borderline personality organisation, particularly accounts of unstable self-image, splitting, fear of abandonment and rapprochement-like oscillation. The safe formulation is analogical and probabilistic: a present pattern resembles a conflict described by the theory. The unsafe formulation is causal and retrospective: the patient has a disorder because a named Mahler phase failed. Adult symptoms cannot reconstruct infancy with that certainty, and no single relational pattern is specific to one developmental history.

Its limitations are examinable. The earliest phases rely on inferences about infant subjectivity; later infant research challenges the idea of an originally closed or wholly undifferentiated infant; the word *autistic* is now misleading; and the original mother-infant dyad may understate fathers, grandparents, siblings, multiple caregivers, temperament and the wider social environment. The theory may also invite caregiver blame if used crudely. Its continuing value lies in organising clinical questions about boundaries, ambivalence and internal representations, not in serving as a literal developmental test.

5.8 Alignment with the major developmental theories

The companion theory-alignment map places the models on a shared age axis, but temporal overlap does not make their constructs interchangeable. Mahler explains separation-individuation and self-object differentiation. Piaget explains changing structures of thought from the sensorimotor stage onward. Erikson organises development through psychosocial stages such as trust versus mistrust and autonomy versus shame and doubt. Freud organises drive and conflict through the psychosexual stages. The map supplies the source-reported age bands; the text must not turn lateral alignment into theoretical equivalence.

Kohlberg describes moral development in the ordered levels preconventional, conventional and postconventional. Kaplan and Sadock relate these to descriptive bands such as before adolescence, during adolescence, later in adolescence, and late adolescence or early adulthood; no sourced numeric ages should be invented. Vygotsky is not a stage theorist. His zone of proximal development, scaffolding, private speech and socially mediated learning are mechanisms that

operate across development, so he belongs beside the age-aligned stage theories rather than inside the plate as a fabricated sequence.

Clinically, each theory asks a different question of the same presentation: Piaget asks how the child represents and reasons; Erikson asks what social conflict is being negotiated; Freud asks how drive and defence are organised; Kohlberg asks how a moral judgement is reasoned; Mahler asks how separateness and connection are managed; Vygotsky asks what learning becomes possible with guided participation. A strong examination answer preserves those levels instead of blending them into a single omnibus theory.

5.9 Nearby relational constructs: theory of mind, language and parenting

Theory of mind is the capacity to attribute mental states to oneself and others and to understand that another person's belief may differ from reality and from one's own belief. *Psychotherapy for Borderline Personality Disorders* reports that passing a false-belief task is expected at **four or five years**. This is not Mahler's object constancy: theory of mind concerns representing another mind, whereas object constancy concerns sustaining an affective representation of a whole attachment object.

The major language acquisition theories also answer a different question.

THEORY	EMPHASIS
Chomsky	emphasises an innate biological preparedness for language and rule formation.
Skinner	emphasises operant learning, reinforcement, shaping and imitation.
Vygotsky	emphasises social mediation, shared activity and the transformation of interpersonal speech into private speech and self-regulation.

Contemporary understanding is not served by making any one of these a subphase of separation-individuation. Language can express emerging self-other distinction, but language development and Mahler's relational sequence are not the same construct.

Baumrind parenting styles classify patterns of warmth and behavioural demand. The classic styles are authoritative, authoritarian and permissive; the later two-dimensional extension adds uninvolved or neglectful parenting. They describe caregiving climate rather than the child's intrapsychic phase. In a clinical interview they help characterise limits, responsiveness and autonomy support, but they neither diagnose a Mahlerian disturbance nor prove how the child experiences the caregiver.

5.10 Indian clinical use and examination reconstruction

INDIA

Mahler's source-reported timings are not Indian developmental norms, and no verified India-specific age adaptation of the theory is supplied here. In Indian families, caregiving may be distributed across parents, grandparents, siblings and other relatives; proximity and interdependence may be valued without implying failed individuation. Formulate the child's actual pattern of boundaries, exploration, availability and emotional representation within that family and culture. Do not mistake a Western ideal of early behavioural independence for psychological health, and do not use Mahler in place of Indian developmental screening or formal assessment.

How to write the long answer. Begin with psychological birth and define separation and individuation separately. Name the precursor phases, then describe hatching, practising, rapprochement and consolidation in order. Attach one organising idea to each: *differentiation, exploration, ambivalence, integration*. Distinguish object constancy from object permanence, give one psychodynamic formulation use, and close with the empirical, cultural and caregiver-blame limitations.

6 . Vygotsky sociocultural theory and zone of proximal development

Lev Vygotsky's sociocultural theory explains cognitive development as a socially mediated process. The child does not develop as an isolated thinker who later enters society. Learning begins in shared activity with more knowledgeable people, is organised by culturally supplied tools and signs, especially language, and is gradually transformed into the child's own psychological functioning. The best-known construct is the **zone of proximal development** (ZPD): the region between what a learner can accomplish independently and what the learner can accomplish through effective guidance or collaboration. The theory is developmental, but it is **not stage-based**. It supplies no Vygotskian sequence of age-linked stages.

How to use this page. Build the answer around one movement: **from shared regulation to self-regulation**. First define social mediation and internalisation, then define the ZPD by its **two** boundaries, explain contingent support and its withdrawal, and connect private speech with the development of self-guidance. End with educational and clinical use, followed by the limits of inferring learning potential from a brief interaction. Schunk's *Learning Theories: An Educational Perspective*, Berk's *Development Through the Lifespan* and Ciccarelli's account support the conceptual treatment; none supplies age stages for Vygotsky.

Must memorise: Development moves from the social plane to the psychological plane: what the learner first does with guidance is gradually internalised and done independently.

6.1 The sociocultural proposition: mind develops through mediated participation

Higher mental functions have social origins and are culturally organised. Vygotsky distinguished elementary capacities with a biological basis from **higher mental functions**, such as deliberate attention, strategic memory, planning and conceptual thought. Higher functions do

not simply unfold from maturation. They are reorganised through participation in culturally structured activity. A caregiver labels objects, directs attention, breaks a task into parts, models a strategy and helps the child remember what to do next. In time, the child takes over these regulatory operations.

This account is dialectical: the social world shapes the child, but the child is not passive. The learner acts, responds, selects tools, alters the interaction and gradually becomes a participant with greater responsibility. Development is therefore neither the direct transmission of adult knowledge nor solitary discovery alone. It is the transformation of activity conducted between people into activity organised within the person.

6.2 Cultural tools, signs and internalisation

Human beings change cognition by using culturally inherited means. **Mediation** occurs when action is organised through a tool or sign rather than by a direct stimulus-response link. Material tools change the external environment; psychological tools and signs change how attention, memory and problem-solving are organised. Language, writing, counting systems, diagrams, mnemonics, maps and culturally taught problem-solving routines are examples. Because cultures provide different tools and practices, the form and priorities of cognition cannot be understood apart from the learner's social setting.

Internalisation is not the copying of an external act into an unchanged inner space. It is a transformation. An instruction first spoken by another person may later be spoken by the child to guide action, then become abbreviated as inner speech. The original interaction is reconstructed as a means of self-regulation. The common summary is that a higher function appears first on the *interpsychological* plane, between people, and later on the *intrapsychological* plane, within the person.

■ **RULE** **Social origin does not mean passive absorption.** Internalisation is an active reorganisation of shared activity into self-directed psychological functioning.

6.3 The zone of proximal development: actual and potential performance

The ZPD is defined by the difference that appropriate assistance makes. The learner's lower boundary is the level of **independent performance**, often called the actual developmental level. The upper boundary is the level of **assisted performance** that becomes possible with adult guidance or collaboration with a more capable peer. The ZPD is the region between these boundaries. It is not everything a child cannot do, and it is not a fixed store of hidden intelligence. A task lies beyond the useful zone when even well-matched assistance does not make performance possible.

Three

INSTRUCTIONAL LOCATIONS

INSTRUCTIONAL LOCATION	DESCRIPTION
Below the ZPD	already mastered independently, so extensive help adds little developmental value.
Within the ZPD	not yet independent, but achievable with responsive guidance, making it the principal teaching zone.
Beyond the ZPD	not achievable even with suitable support at that point, so prompting alone becomes unproductive.

The ZPD shifts attention from a static question, *What can this learner do now?*, to a developmental question, *What can this learner do with help, and how much help is required?* Two learners with the same unaided score may differ in responsiveness to modelling, prompts, feedback and transfer. Conversely, assisted success must not be counted as independent mastery. The quality of the support, the learner's familiarity with the language and task, motivation, anxiety, fatigue and the relationship with the examiner can all affect the observed zone.

6.4 Scaffolding: contingent support, fading and transfer of responsibility

Scaffolding is the instructional process most commonly used to operationalise the ZPD. The more knowledgeable participant recruits attention, simplifies the task, models a step, asks a leading question, supplies a cue, highlights an error or maintains motivation. Support is *contingent*: it increases when the learner is stuck and decreases when competence appears. The aim is not permanent assistance but a transfer of responsibility to the learner through fading.

The term *scaffolding* was introduced later by Wood, Bruner and Ross rather than coined by Vygotsky. It is compatible with his account, but the **two** terms are not synonyms. The ZPD describes a relation between present independent performance and assisted possibility; scaffolding describes one way of supporting activity within that relation. Good scaffolding is calibrated and temporary. Giving the answer, overprompting every step or withdrawing help abruptly can all obscure what the learner understands.

■ **TRAP** **ZPD is not the task itself, and scaffolding is not unlimited help.** The zone is defined by the learner's change in performance under appropriate assistance. Support must be adjusted and then faded until the learner regulates the task independently.

6.5 Language, private speech and the growth of self-regulation

Speech changes from a means of coordinating with others into a means of coordinating oneself. In shared activity, **social speech** allows a caregiver and child to direct attention, label a goal and organise action. The child later uses **private speech**, audible speech addressed to the self, to plan, inhibit an impulsive response, sequence steps and correct errors. With increasing internalisation, this becomes condensed **inner speech**. Inner speech is not merely ordinary conversation moved silently into the head; it becomes abbreviated and shaped for self-guidance.

Private speech is therefore functional rather than simply immature. It is especially likely when a task is difficult, novel or requires control, and overt self-talk may reappear when an older learner is challenged. The classic contrast is with Piaget, who treated egocentric speech mainly as

evidence that the young child had not yet fully accommodated the listener's perspective. Vygotsky instead emphasised its transition from interpersonal communication to intrapersonal regulation. The views address different features of children's self-directed talk and should not be reduced to a slogan that one theorist found it useless and the other useful.

6.6 Learning leads development: teaching within the emerging competence

Vygotsky rejected teaching that merely waits for fully formed capacities. Instruction is most developmentally productive when it engages functions that are emerging and supports performance within the ZPD. In this sense, appropriately organised learning leads development: guided participation creates opportunities for the learner to practise forms of attention, reasoning and self-regulation that later become independent. This does not mean that any teaching can override readiness, neurodevelopmental constraints or prior knowledge. The learner must be able to participate meaningfully in the supported activity.

Applications include modelling with gradual withdrawal, guided questioning, cooperative learning, peer tutoring, reciprocal teaching, cognitive apprenticeship and worked examples that become progressively incomplete. A more capable peer may support learning, but peer status alone does not guarantee effective mediation. The peer must understand the task, communicate accessibly, notice the learner's response and adjust help. Collaboration is valuable when responsibility and explanation are shared, not when one student completes the work for another.

6.7 Dynamic assessment and clinical use

Dynamic assessment extends the ZPD into assessment by embedding teaching within the evaluation. A test-teach-retest format, graduated prompts or mediated learning can show which cues unlock performance, how rapidly a strategy is acquired, whether improvement transfers to a related task and how much assistance remains necessary. This complements a static score by sampling learning processes and modifiability. It does not make standardised assessment redundant, and assisted performance should be reported separately from unaided performance.

CLINICAL USE

When a child cannot organise a multistep task, first observe independent performance. Then offer the least intrusive useful support, such as restating the goal, modelling the first step or giving a visual sequence, and record the response. Improvement with modest, transferable support suggests an accessible learning pathway; persistent difficulty despite well-matched help suggests the need to examine language comprehension, attention, memory, intellectual functioning, anxiety, sensory factors and the suitability of the task. This is formulation evidence, not a diagnosis.

In psychiatry, the same logic can guide cognitive remediation, executive-function support, psychoeducation and rehabilitation. The clinician identifies what the patient can manage alone, co-constructs the next step and withdraws prompts as self-management improves. The goal is agency, not dependence on the therapist. A person's need for scaffolding may vary across tasks and states, so the ZPD should not be treated as a single stable trait or converted into a global intelligence label.

6.8 Alignment with stage theories: a cross-cutting mechanism, not another row

The accompanying theory-alignment plate places different developmental questions beside one another. Piaget describes changes in cognitive structure from the sensorimotor period onward. Erikson organises development through psychosocial stages and their central conflicts. Freud proposes psychosexual stages organised around drive, bodily focus and conflict. Kohlberg describes ordered forms of moral development, but his sourced age relationships are descriptive bands rather than numeric ranges. Mahler describes separation-individuation and the development of psychological separateness in relationship.

Vygotsky differs structurally from all five: sociocultural theory and the zone of proximal development specify mechanisms through which learning and self-regulation can occur across development. There are no Vygotskian stages to align by age. The plate should therefore show Vygotsky beside the stage sequences as a cross-cutting mechanism, never as an invented age row. Temporal overlap between theories also does not make their constructs equivalent: cognitive structure, psychosocial conflict, psychosexual organisation, moral reasoning, separation and supported learning answer different questions.

6.9 Culture, Indian clinical interpretation and limitations

Culture is constitutive in the theory, not decorative context added after cognition develops.

Languages, scripts, number systems, schooling practices, work routines, narratives and patterns of adult-child participation influence which skills are practised and how they are organised. This is a major strength of the framework. It also creates a methodological challenge: performance in an unfamiliar testing interaction may underestimate competence acquired through everyday participation, while familiarity with school-like dialogue may advantage another learner.

INDIA

In Indian practice, mediation may be distributed across parents, grandparents, siblings, teachers and peers, and learning may occur through observation, participation, recitation, demonstration or direct questioning. These forms should be described rather than ranked against a single classroom ideal. Language of testing matters in a multilingual child, as does familiarity with the examiner's style. No verified India-specific numeric ZPD norm or Vygotskian age-stage adaptation is supplied here, and none should be invented. The absence of an age table is expected because the ZPD is relational and task-specific, not an age stage.

The principal criticisms are equally examinable.

- The theory gives powerful concepts but less precise specification of how broad cultural processes produce particular individual cognitive changes.
- The boundaries of a ZPD can be difficult to measure reliably because assistance is interactive and examiner-dependent.
- A more capable partner may transmit error, prejudice or conformity as well as skill.
- Social mediation can be overstated at the expense of biological maturation, temperament, spontaneous exploration and individual construction.

- Successful performance with help does not prove durable learning unless support can be faded and the strategy transfers beyond the teaching episode.

6.10 Examination reconstruction

Long-answer sequence. Define sociocultural theory, state the social origin of higher mental functions, explain cultural tools and internalisation, define both boundaries of the ZPD, distinguish the ZPD from scaffolding, trace social speech to private and inner speech, then add dynamic assessment, one clinical use and the principal criticisms. Close by stating that Vygotsky proposed no age-linked stages.

MNEMONIC

Social activity - mediation - internalisation - ZPD - contingent support - fading - self-regulation.

It traces the movement from shared regulation to self-regulation. If that chain is intact, the rest of the answer can be rebuilt rather than recited.

7 . Bowlby and Ainsworth attachment theory and Strange Situation

Attachment is an enduring, selective emotional bond in which a child seeks proximity to a familiar caregiver, especially when frightened, tired, ill or uncertain. John Bowlby supplied the evolutionary and developmental theory; Mary Ainsworth converted that theory into naturalistic observation and a standardised laboratory assessment, the Strange Situation. The central idea is functional rather than sentimental: the caregiver becomes a **safe haven** under threat and a **secure base** from which the child can explore. The examiner usually wants **three** distinctions. Attachment is not simply feeding or dependency, an attachment pattern is not a fixed personality trait, and insecure attachment is not in itself a psychiatric disorder.

How to use this page. First learn Bowlby's attachment behavioural system and internal working model. Then learn why Ainsworth treated exploration, separation and reunion as one organised system. Finally, identify the **four** patterns from reunion behaviour: secure, avoidant, ambivalent or resistant, and disorganised. The final section connects attachment with normal adolescent development, because adolescence reorganises rather than abolishes the secure-base relationship.




Fig 20.6 The Strange Situation and the four attachment patterns, keyed to behaviour on separation and on reunion. The separation row is drawn grey and undifferentiated because separation distress varies and does not classify; the reunion row carries the classification. The organised A, B and C patterns follow Ainsworth's original system, the disorganised D classification follows Main and Solomon. No age, episode count or pattern distribution appears, because none was verified in the sourced texts.

7.1 Bowlby: attachment as an evolved behavioural system

Proximity to a protective caregiver is the set goal. Bowlby rejected the drive-reduction claim that the infant becomes attached merely because the caregiver supplies food. He drew on psychoanalysis, ethology, control-systems thinking and observations of separation to propose an innate **attachment behavioural system**. Signals such as crying, smiling, calling, clinging and following increase caregiver proximity. The system is activated by threat, pain, fatigue, unfamiliarity or separation; it is relatively quiet when the child feels safe, allowing exploration and play. Attachment and exploration are therefore complementary systems. A securely based child is not continuously close to the caregiver. The child moves away to explore and returns when safety must be restored.

Bowlby's term monotropy described a tendency for one attachment figure to acquire special importance, not a biological rule that a child can form only one bond. Children can develop a hierarchy of attachments to several consistent caregivers. What matters clinically is not whether care matches a narrow family form, but whether available caregivers are familiar, responsive and sufficiently predictable. Bowlby's early writing on maternal deprivation was historically influential, but a modern reading separates the effects of prolonged loss, disrupted continuity,

insensitive care and the wider adversity that often accompanies them. The theory predicts risk, not destiny.

- **RULE** **Attachment is a safety-regulation system, not a measure of how much a child loves a parent.**
Read proximity-seeking, exploration and reunion as organised behaviour serving protection.

7.2 Internal working models and developmental organisation

Repeated interactions become expectations about self, others and help. Through recurrent caregiver responses, the child constructs **internal working models**: expectations about whether distress will be noticed, whether another person will be available, and whether the self is worthy and effective in eliciting care. These models guide attention, affect regulation, memory, help-seeking and later relationships. A history of responsive care favours open signalling and confidence in comfort; recurrent rejection may favour suppression of attachment signals; inconsistent response may amplify them. These are adaptive strategies in a particular relationship, not conscious decisions and not immutable templates.

Bowlby's developmental account is commonly organised as a phase sequence, and the qualitative progression runs as follows:

PHASE	QUALITATIVE PROGRESSION
Pre-attachment	broadly directed social signals.
Attachment-in-the-making	preference for familiar caregivers.
Clear-cut attachment	active proximity regulation.
Goal-corrected partnership	an increasing ability to represent the caregiver's goals and negotiate separation.

Working models are carried forward through continuity of expectations and relationships, but they remain open to revision through later caregiving, reflective capacity, psychotherapy and corrective relationships. Stability is therefore probabilistic, with both continuity and change expected.

7.3 Ainsworth: sensitivity, secure base and observational method

Ainsworth made caregiver-child organisation observable. Her naturalistic work in Uganda and later longitudinal observations in Baltimore refined Bowlby's theory. She described the infant's use of the caregiver as a secure base and emphasised **caregiver sensitivity**: noticing the child's signals, interpreting them accurately, and responding promptly and appropriately. Sensitivity is associated with security, but neither one interaction nor one caregiver characteristic mechanically produces a classification. Child temperament, caregiver state, cumulative experience, culture and the immediate context all contribute to observed behaviour.

Ainsworth's major methodological advance was to judge the organisation of behaviour rather than count isolated acts. Crying alone does not define insecurity, and a child who does not cry is not automatically secure. The question is whether the child can use the caregiver to regulate arousal and then resume exploration. This shift from single behaviours to the pattern linking

separation, proximity-seeking, contact, soothing and renewed play is the conceptual bridge to the Strange Situation.

7.4 Strange Situation: procedure, observations and scoring logic

A standardised, progressively stressful observation of attachment and exploration. The **Strange Situation Procedure** places a young child and caregiver in an unfamiliar room, introduces an unfamiliar adult, and includes brief caregiver departures and reunions. These manipulations activate the attachment system while leaving opportunities for play. Observers code exploration, distress on separation, response to the unfamiliar adult, proximity-seeking, contact maintenance, resistance, avoidance and the child's capacity to settle. The reunions carry the greatest classificatory weight because they show what the child does when the attachment figure becomes available after stress.

The procedure does not provide a global score of parental goodness, prove maltreatment, measure the total strength of the bond or diagnose a disorder. It samples behaviour in a structured context and classifies the organisation of one child's attachment to one caregiver. The original Ainsworth system described the organised A, B and C patterns. Mary Main and Judith Solomon later formalised the disorganised or D classification for behaviour that lacks a coherent approach to the attachment dilemma.

Must memorise: In the Strange Situation, reunion behaviour is decisive: secure children seek and use comfort, avoidant children minimise contact, resistant or ambivalent children seek yet resist it, and disorganised children show no coherent strategy.

MNEMONIC

"Seek, minimise, seek yet resist, no coherent strategy"

In the Strange Situation, reunion behaviour is decisive: secure children seek and use comfort, avoidant children minimise contact, resistant or ambivalent children seek yet resist it, and disorganised children show no coherent strategy.

7.5 Secure and avoidant attachment patterns

Secure, pattern B: distress can be signalled, comfort can be used, exploration can restart. A **secure** child generally explores in the caregiver's presence, may or may not show marked separation distress, greets or approaches the caregiver on reunion, accepts contact or soothing, and returns to play once regulated. Security is therefore not absence of distress. It is the flexible and effective use of the caregiver when distressed. The associated caregiving history is commonly sensitive, available and reasonably predictable, but the classification remains descriptive and probabilistic rather than a retrospective proof of caregiving quality.

Avoidant, pattern A: attachment signals are minimised. An **avoidant** child may continue exploring, show little overt separation protest, and avoid or ignore the caregiver on reunion. If picked up, the child may not actively maintain contact and may turn attention toward objects. The crucial interpretation is strategic minimisation of visible attachment behaviour in that

relationship, often associated with expected rejection or discomfort around bids for closeness. Apparent composure must not be equated with absence of arousal, absence of attachment or superior independence.

7.6 Resistant or ambivalent and disorganised attachment patterns

Resistant or ambivalent, pattern C: attachment signals are intensified but do not readily achieve settling. The **ambivalent** or resistant child often explores less, becomes highly distressed by separation, and on reunion combines proximity-seeking with anger, resistance or inability to settle. The child may demand contact and simultaneously push away, arch or remain preoccupied with the caregiver. This pattern is understood as maximising attachment signals when caregiver availability has been experienced as inconsistent. The word ambivalent refers to the observed mixture of approach and resistance, not to ordinary mixed feelings.

Disorganised, pattern D: the attachment strategy breaks down. A **disorganised** child shows contradictory, incomplete, apprehensive or markedly disoriented behaviour in the caregiver's presence, such as approaching and then freezing, backing away, collapsing, or displaying incompatible sequences without a stable strategy. The caregiver may be simultaneously the expected source of safety and a source of alarm, creating fear without solution. Disorganisation is associated with frightening or frightened caregiving, unresolved caregiver trauma, maltreatment and severe family stress, but it is not specific to any one cause. It may be coded alongside the best-fitting organised pattern and must never be treated as proof that abuse occurred.

■ **TRAP** **Do not convert attachment classifications into diagnoses.** Avoidant does not mean unattached, disorganised does not prove maltreatment, and insecure attachment is not synonymous with reactive attachment disorder or disinhibited social engagement disorder. RAD and DSED are separate clinical disorders linked to a history of severely insufficient care, and they are developed in the Child behavioural, emotional and other disorders notes.

7.7 Interpretation, cultural caution and clinical use

Classify the relationship in context, not the child's essence. Attachment security may differ across caregivers because each relationship has a different history. Classifications show moderate continuity at group level when care and context remain stable, but individual change is possible. Temperament can affect the intensity and style of distress without by itself determining security. Cultural practices also shape baseline separation experience, proximity, stranger familiarity and the social meaning of independence. Cross-cultural differences in pattern distribution therefore require ecological interpretation rather than a hierarchy of cultures.

Clinically, attachment theory helps formulate emotion regulation, help-seeking, interpersonal expectations and the meaning of separation. It does not justify diagnosing an attachment style from an adult's brief self-report, reading childhood causation directly from a current relationship, or blaming one caregiver. Adult attachment interviews and romantic-attachment questionnaires examine related but non-identical constructs and are not interchangeable with the Strange Situation. In a psychiatric assessment, ask:

- Who actually provided care.

- How the child sought comfort.
- How caregivers responded.
- Whether separations or changes in placement occurred.
- How the relationship functions now.

Use these observations to build a developmental formulation, not a label masquerading as explanation.

7.8 Attachment in adolescence: the secure base at a distance

Adolescence changes the regulation of closeness; it does not end attachment. In normal adolescent development, the secure-base function becomes less dependent on immediate physical proximity and more dependent on the expectation that support is available. Autonomy and attachment are not opposites. Security permits exploration of education, peers, values and intimate relationships, while caregivers remain a refuge under significant stress. Peers and romantic partners acquire attachment functions, but this redistribution is gradual and relationship-specific.

3 EXAM DISTINCTIONS	4 REUNION PATTERNS
-------------------------------	------------------------------

8 . Temperament and goodness of fit (Thomas and Chess)

Temperament is the characteristic style with which a child responds to the world: the tempo, intensity, regularity and flexibility of behaviour rather than its particular content. Thomas and Chess captured the distinction neatly by asking *how* a child behaves, not *what* the child does or *why* the child does it. Temperament appears early, has a constitutional contribution and shows some continuity, but it is neither an immutable biological destiny nor a diagnosis. Its expression changes with development, relationships, learning, culture and context.

How to use this page. First learn the **nine dimensions**, then the **three types** derived from recurring combinations of those dimensions. Finally, understand **goodness of fit**, because that is the clinically important part of Thomas and Chess: adaptation depends not on a supposedly good or bad child, but on the match between the child's temperament and the demands, expectations and supports of the environment. In a theory answer, define temperament, list the nine dimensions, describe the three types with their classic proportions, explain goodness of fit and finish with clinical application and limitations.

The must-memorise line: Thomas and Chess described nine dimensions and three temperamental constellations, but outcome depends chiefly on goodness of fit between the child and the environment.

Thomas and Chess: nine dimensions, three types, and fit as a relation

Temperament is behavioural style: how a child responds, not what the child does or why. The clinically important part is the last one, and it is not a property of the child.

The nine dimensions of temperament

Continuous behavioural variables, not nine categories and not nine disorders. Every child occupies a position on all nine, and the position may be clearer in some settings than others.

1. Activity level amount of motor activity, active versus inactive periods	2. Rhythmicity or regularity predictability of sleep, hunger, feeding, elimination	3. Approach or withdrawal direction of the first response to anything new
4. Adaptability ease of change after continued exposure	5. Threshold of responsiveness how much stimulation is needed to evoke a response	6. Intensity of reaction energy of the response, positive or negative
7. Quality of mood predominance of pleasant versus unpleasant behaviour	8. Distractibility ease with which ongoing behaviour is interrupted	9. Attention span and persistence duration pursued, continuation despite obstacles

Three viva traps. Approach or withdrawal is the first reaction, adaptability is the later adjustment after continued exposure. Threshold is how much stimulation is needed to produce a response, intensity is how strong the response is once produced. Distractibility is interruption by a competing stimulus, persistence is maintenance of the original activity.

The three types, and the group they do not cover

Recurring combinations of the nine dimensions in the original observations. Descriptive constellations, not diagnoses, not moral grades and not mutually exhaustive.

Easy Regular rhythms, positive approach to novelty, positive mood, mild or moderate reactions, ready adaptation to change.	Difficult Irregular rhythms, withdrawal from novelty, slow adaptation, intense reactions, predominantly negative mood.	Slow-to-warm-up Low activity, initial withdrawal, slow adaptation, mild reactions. Acceptance grows with repeated gentle exposure.	Not classifiable Combinations that met none of the three patterns. Not a fourth type, and it has no characteristic profile of its own.
---	---	---	---

Proportions. The classic study reported an approximate percentage for each type, plus a large unclassified remainder. No percentage is drawn on this plate, because none was verified against a named source here. If you quote them, attribute them to that study's own sample, never as prevalence in India.

Goodness of fit: the trait is held constant, the setting changes, the outcome flips

The compatibility between the child's style and the demands, expectations and opportunities of the environment. A trait has no fixed adaptive value outside a context.

The same trait	A setting where the trait fits	A setting where the same trait does not fit
High activity level dimension 1	Useful in sport, neutral in outdoor play, and manageable at home when movement is planned before seated work.	Disruptive in a classroom built around prolonged stillness. The child has not changed. The demand has.
Initial withdrawal dimensions 3 and 4	Protects the child from impulsive entry into unfamiliar situations.	Creates difficulty when adults equate immediate sociability with competence.
High persistence dimension 9	Supports mastery of a complex task, because effort continues past the first obstacle.	Becomes inflexibility at the moment stopping or shifting set is what the situation requires.
Low threshold dimension 5	Permits fine detection of subtle stimuli during a quiet activity.	Contributes to overload in a noisy ward. The same threshold, the opposite result.

Trap. Goodness of fit does not mean that an easy child is good and a difficult child is bad.

The three types are descriptive clusters, not diagnoses or moral grades. Even an easy temperament can be poorly matched to an environment, and a difficult temperament can have an adaptive outcome when expectations and supports fit the child. Outcome is a property of the relationship, not of either party.

Nor is bidirectionality covert parent-blaming. Poverty, crowded housing, school structure, stigma, family conflict and caregiver illness can restrict the environment's capacity to adapt.

-  Good fit. Demands are reasonably consonant with the child's characteristics, and adults provide supports that allow adaptive functioning.
 Poor fit. Demands repeatedly exceed, contradict or punish the child's characteristic way of responding without adequate accommodation.
 The environment axis: the demands, expectations, supports and cultural expectations that the trait actually meets.
 Child-size description, deliberately left uncoloured. No dimension and no type carries a good or a bad value on its own.
 Not a type. The unclassified group has no characteristic profile.

The dimensions, the types, goodness of fit and every clinical example above follow this chapter's topic text on Thomas and Chess and the New York Longitudinal Study.

Not sourced, and therefore not drawn: the classic percentage for each type and for the unclassified remainder; any prevalence of the three types in Indian children; any age at which a temperament type is said to be established or to become stable. No label on this plate is a referral threshold, and no type on it is a diagnosis.

Fig 20.7 Thomas and Chess drawn as an interaction. The nine dimensions and the three types are the child side and are left uncoloured, because no dimension and no type carries a value on its own; a fourth box marks the group the three types did not cover, which is not a fourth type. Colour appears only in the fit table, where one trait is held constant and meets two settings, and the outcome reverses without the child changing. Content follows this chapter's topic text on Thomas and Chess and the New York Longitudinal Study. The classic proportions for each type are named on the plate but not drawn, because no percentage for them was verified against a named source here.

8.1 Thomas and Chess: the interactional view of temperament

Temperament is behavioural style, and development is transactional. In the New York Longitudinal Study, Alexander Thomas, Stella Chess and colleagues followed children and their families to examine early individual differences and later adjustment. Their work challenged explanations that attributed children's behavioural problems either entirely to innate defect or

entirely to parental handling. Children differ from the outset in how readily they become aroused, approach novelty, adapt to change, sustain activity and express mood. These differences alter how caregivers respond; caregiver responses then shape how the child's style is expressed and regulated. Child and environment therefore influence one another over time.

This is a **transactional model**, not a one-way causal model. A highly active child may invite more prohibition from an exhausted caregiver, while the resulting conflict may further amplify restless behaviour. The same child in a setting with safe movement, brief instructions and predictable transitions may function well. Conversely, an initially easy interaction can deteriorate when environmental demands change. Temperament contributes to the probability and form of an interaction; it does not independently dictate psychopathology.

■ **RULE** **Describe the child, the environment and the match.** A Thomas-and-Chess formulation is incomplete if it lists traits but does not show how environmental demands, caregiver responses and cultural expectations convert those traits into adaptation or difficulty.

8.2 The nine dimensions of temperament

The nine dimensions are continuous behavioural variables, not nine separate disorders. Each child occupies a position on every dimension, and the position may be more evident in some settings than in others. The examiner usually expects the complete list with a one-line meaning:

DIMENSION	MEANING
Activity level	the amount of motor activity and the relative proportion of active to inactive periods. Ask how much the child moves during play, meals, travel, learning and sleep-related routines.
Rhythmicity or regularity	the predictability of biological functions and routines, including sleep, hunger, feeding and elimination.
Approach or withdrawal	the direction of the initial response to a new person, place, food, activity or other stimulus. Approach means initial acceptance or engagement; withdrawal means initial hesitation, avoidance or distress.
Adaptability	the ease and speed with which behaviour changes after continued exposure to novelty or altered circumstances. It is distinct from the first response: a child may initially withdraw yet adapt well after familiarisation.
Threshold of responsiveness	the intensity of stimulation required to evoke a response. A low threshold means that weak light, sound, texture, temperature or social stimulation is sufficient; a high threshold means that stronger stimulation is needed.
Intensity of reaction	the energy level of the response, whether positive or negative. It concerns how strongly emotion or behaviour is expressed, not which emotion is expressed.
Quality of mood	the relative predominance of pleasant, friendly and joyful behaviour versus unpleasant, distressed or unfriendly behaviour.
Distractibility	the ease with which an ongoing behaviour is interrupted or altered by extraneous stimulation. It is not synonymous with a disorder of attention.
Attention span and persistence	the duration for which an activity is pursued and the tendency to continue despite obstacles. Attention span concerns sustained engagement; persistence concerns continuation when continuation becomes difficult.

Three distinctions prevent common viva errors. **Approach-withdrawal** is the initial reaction, whereas **adaptability** is the later adjustment. **Threshold** is the amount of stimulation needed to produce a response, whereas **intensity** is the strength of the response once produced.

Distractibility is interruption by competing stimuli, whereas **attention span and persistence** describe maintenance of the original activity.

8.3 The three types: easy, difficult and slow-to-warm-up

The three types are clusters of dimensions found repeatedly in the original observations. They are descriptive constellations, not mutually exhaustive categories. A child's profile may resemble one constellation, contain features of several, or fit none. The labels are historically important and highly examinable, but the word *difficult* must be handled carefully because it describes a pattern of interactional challenge rather than a defective child.

TEMPERAMENTAL CONSTELLATION	CHARACTERISTIC PATTERN	LIKELY INTERACTIONAL ISSUE
Easy temperament	Regular biological rhythms, positive approach to novelty, relatively positive mood, mild or moderate reactions and ready adaptation to change	Usually elicits confident and rewarding caregiving, but genuine distress may be overlooked because the child is usually manageable
Difficult temperament	Irregular rhythms, withdrawal from novelty, slow adaptation, intense reactions and predominantly negative mood	May evoke coercion, criticism or inconsistent responses, producing escalating conflict when expectations are rigid
Slow-to-warm-up temperament	Low activity, initial withdrawal, slow adaptation and relatively mild reactions; acceptance increases gradually with repeated non-threatening exposure	May be misread as stubborn, uninterested or incapable if adults demand rapid participation

9	40%	10%	15%
DIMENSIONS IN THE THOMAS AND CHESS FRAMEWORK	APPROXIMATELY EASY IN THE CLASSIC CLASSIFICATION	APPROXIMATELY DIFFICULT IN THE CLASSIC CLASSIFICATION	APPROXIMATELY SLOW-TO-WARM-UP IN THE CLASSIC CLASSIFICATION

The proportions belong to the classic study classification and should not be presented as universal prevalence estimates for contemporary Indian children. Their examination value lies in showing that the three types never covered the entire population. The unclassified group is not a fourth type. In the classic study classification, 35% were approximately mixed or not classifiable into the three types; their combinations did not meet one of the three characteristic patterns.

8.4 Goodness of fit: the central clinical proposition

Goodness of fit is the compatibility between the child's temperamental capacities and style, and the demands, expectations and opportunities of the environment. A good fit occurs when environmental demands are reasonably consonant with the child's characteristics and when adults provide supports that allow adaptive functioning. A poor fit occurs when demands repeatedly exceed, contradict or punish the child's characteristic way of responding without adequate accommodation or skill-building. The outcome is a property of the relationship between child and context, not a moral property located inside either one.

Consider high activity. It may be disruptive in a classroom built around prolonged stillness, useful in sport, neutral during outdoor play and manageable at home when movement is planned before seated work. Initial withdrawal may protect a child from impulsive entry into unfamiliar

situations, yet create difficulty when adults equate immediate sociability with competence. High persistence may support mastery of a complex task but become inflexibility when stopping or shifting is required. A low sensory threshold may contribute to overload in a noisy ward yet permit fine detection of subtle stimuli in a quiet activity. The trait has no fixed adaptive value outside a context.

Goodness of fit also includes culture. Families and schools differ in the value placed on quietness, assertiveness, emotional expressiveness, regularity, rapid sociability and obedience. A style praised in one setting may be discouraged in another. Cultural formulation does not mean accepting severe impairment or harmful caregiving. It means identifying whose expectation is being used, what the setting actually requires and whether that requirement can be modified without compromising the child's development or safety.

■ **TRAP** **Goodness of fit does not mean that an easy child is good and a difficult child is bad.** The three types are descriptive clusters, not diagnoses or moral grades. Even an easy temperament can be poorly matched to an environment, and a difficult temperament can have an adaptive outcome when expectations and supports fit the child.

8.5 From temperament to outcome: risk, protection and bidirectional effects

Temperament affects outcome through repeated transactions rather than direct

determinism. A temperamental feature can influence exposure to situations, the responses elicited from other people, the child's interpretation of those responses and opportunities to practise regulation. When a mismatch is chronic, child and adult may enter a self-reinforcing sequence. Intense protest evokes harsh control; harsh control increases arousal and protest; both participants then anticipate conflict. When fit improves, the sequence can reverse: predictable preparation reduces arousal, calmer participation evokes more supportive responses, and successful experiences expand the child's tolerance.

Temperament can therefore function as a risk factor, a protective factor or a modifier of how other risks are expressed. It may influence the form of a symptom without being its cause. Inhibited approach can resemble anxiety, high activity can resemble hyperactivity, negative mood can resemble depression, irregularity can complicate sleep and feeding, and low persistence can look like defiance or inattention. The resident must establish pervasiveness, developmental change, subjective distress, functional impairment and the full syndrome before moving from style to disorder.

Parent and child effects are bidirectional, but bidirectionality must not become covert parent-blaming. Caregivers bring their own temperament, health, stress, beliefs, resources and expectations. A mismatch may occur between a highly intense child and a highly intense caregiver, or between an active child and a caregiver whose circumstances allow little safe movement. Poverty, crowded housing, school structure, stigma, family conflict and caregiver illness can restrict the environment's capacity to adapt even when the caregiver understands the child well.

8.6 Clinical assessment of temperament and fit

Assess a pattern across time and settings, not a single performance in the clinic. Begin with concrete examples rather than labels. Instead of asking whether the child is difficult, ask what happens with an unfamiliar person, an unexpected change, waiting, loud sound, a missed meal, a challenging task or a transition away from preferred play. Establish the trigger, initial response, intensity, duration, adaptability after continued exposure, recovery and effect on function. Map the answers onto the nine dimensions only after hearing the behaviour.

Take accounts from the child where developmentally appropriate, caregivers and teachers, because temperament is expressed differently across contexts and each observer has a relationship-specific view. Direct observation should sample free activity, structured demands, novelty, transition, frustration and recovery. The clinic itself is an unfamiliar and stimulating environment, so withdrawal, silence or high activity there cannot be assumed to represent the child's usual style. Look for both stable tendencies and situational exceptions; exceptions reveal environments in which fit is already better.

A useful formulation has four parts:

1. **Child profile:** identify the prominent dimensions and strengths using neutral behavioural language.
2. **Environmental profile:** identify family, school and cultural demands, available supports and the temperaments of important caregivers.
3. **Fit and transaction:** specify where mismatch occurs, how each participant responds and what cycle maintains the difficulty.
4. **Clinical boundary:** assess impairment and alternatives, including neurodevelopmental disorder, anxiety, mood disturbance, sleep or medical problems, trauma, attachment disturbance, sensory impairment and adverse environment.

Temperament is not a diagnosis of exclusion, but neither should it be used to explain away a disorder. The key question is whether the nine-dimension profile adequately describes a stable style or whether there is an additional syndrome with distress, impairment, developmental deviation or episodic change. A previously characteristic style that changes markedly should prompt assessment for state-related causes rather than being relabelled as temperament.

8.7 Management: change the fit, support regulation and preserve strengths

The intervention target is usually the transaction, not eradication of the trait. The clinician first reframes the behaviour in non-moral language, helps caregivers anticipate predictable points of mismatch and designs the environment so that the child can succeed while gradually building capacity. This is sometimes called temperamental guidance. It validates the child's style without allowing harmful behaviour and supports the caregiver without implying that every difficulty results from parenting.

Management follows the functional profile. An irregular child benefits from external predictability and consistent cues without punitive battles over perfect regularity. A child who

withdraws from novelty benefits from preparation, observation from a safe distance, gradual entry and repeated low-pressure exposure rather than forced sociability. A highly active child needs planned movement, brief achievable seated tasks and safe outlets. A child with intense reactions needs calm adult responses, simple limits and help labelling early arousal before escalation. A child with a low threshold may need reduction of avoidable sensory load and graded coping with unavoidable stimulation. Low persistence calls for task chunking, visible completion points and reinforcement of effort, whereas very high persistence may require supported transitions and practice shifting set.

The environment must adapt, but accommodation is not capitulation. A good fit both accepts the child's starting style and teaches the skills needed for wider participation. Expectations should be realistic, explicit and gradually increased. School collaboration is often decisive because the child may spend much of the waking routine under group demands. Review whether the change reduces conflict, improves participation and preserves the child's strengths; if impairment persists, reassess the formulation and comorbidity.

8.8 Strengths, limitations and later extensions

The model's enduring strength is that it moved child psychiatry from blame toward person-environment interaction. It offered an observable vocabulary for early behavioural individuality, recognised the child's active contribution to relationships and generated the clinically useful concept of goodness of fit. It also anticipated contemporary developmental psychopathology by treating outcome as the product of transactions over time rather than a simple line from one early trait to one later disorder.

Its limitations are equally examinable. The three types compress continuous traits and leave a substantial group unclassified. Several dimensions overlap conceptually, and ratings can be influenced by the observer's expectations, mood and relationship with the child. Stability is relative rather than absolute because the expression of temperament changes with maturation and context. The classic proportions arose from a particular sample and should not be universalised across socioeconomic and cultural settings. Finally, later temperament models organise traits differently and give greater attention to reactivity, self-regulation and biological mechanisms, so Thomas and Chess should be presented as a foundational framework rather than the final taxonomy of temperament.

The historical labels also require ethical translation. *Difficult* is useful in an examination when explicitly attributed to the Thomas-and-Chess classification, but clinical communication should describe the actual profile: irregular, initially withdrawing, slowly adaptive, intensely reactive or predominantly negative in mood. Precise description reduces shame and leads directly to intervention. A global label does neither.

8.9 Examination synthesis and retrieval

A complete short note can be built in five moves. Define temperament as behavioural style; name Thomas and Chess and the New York Longitudinal Study; list all nine dimensions; describe easy, difficult and slow-to-warm-up temperaments with the classic proportions and the

unclassified remainder; then explain goodness of fit with one clinical example, bidirectionality, management and limitations. If time is short, marks are lost most often by omitting threshold, confusing adaptability with approach, or listing the three types without explaining fit.

MNEMONIC

"move, clock, meet, change, notice, react, feel, shift, stay"


Move recalls activity level; clock, rhythmicity; meet, approach or withdrawal; change, adaptability; notice, threshold; react, intensity; feel, quality of mood; shift, distractibility; and stay, attention span and persistence. The sequence is only a retrieval aid. In the answer, use the formal terms.

Worked viva formulation: A child initially refuses unfamiliar classroom activities, reacts mildly and joins after repeated observation. This is closer to a slow-to-warm-up pattern than a difficult pattern because intensity is low and adaptation occurs gradually. If the teacher permits observation and phased entry, participation improves: that is better goodness of fit. If the child is forced to perform immediately and is repeatedly criticised, withdrawal may increase: that is poorer fit. Neither response alone establishes an anxiety disorder; distress, pervasiveness, persistence and impairment still require assessment.

Final clinical synthesis. Thomas and Chess supplied a vocabulary for the child's characteristic style, but their larger contribution was relational. Temperament describes how the child enters an interaction; goodness of fit explains why the same style can lead to competence in one environment and conflict in another. The mature answer neither blames the child nor idealises accommodation. It identifies the profile, maps the environment, changes the transaction and assesses separately for disorder.

9 . Theory of mind development

Theory of mind is the capacity to attribute mental states to oneself and to other people, and to use those attributions to explain and predict behaviour. The attributed states include intentions, desires, knowledge, beliefs, emotions and pretence. The decisive idea is representational: another person acts on how the world is represented in that person's mind, not necessarily on how the world really is or on what the observer knows. Theory of mind therefore permits the child to understand deception, surprise, misunderstanding, secrecy, irony and the fact that two people can interpret the same event differently. It is often called **mentalizing**, although clinical literature sometimes uses mentalizing more broadly for the imaginative interpretation of self and others in terms of intentional mental states.



■ Joint attention

■ Pretence

■ First-order belief

■ Second-order belief

■ Mental-state inference

Fig 20.8 Theory of mind develops through interacting social, representational, linguistic and executive capacities. False-belief understanding is the central test: another person acts on what they believe, even when that belief conflicts with reality.

This is not a single switch and not a synonym for sociability. Its development rests on an interacting set of capacities: social attention, joint attention, symbolic representation, language, executive control, autobiographical memory and experience of conversation. A child may be affectionate yet fail a formal false-belief task because the language is too difficult; another may pass a structured task yet misunderstand rapidly changing intentions in ordinary interaction. For an examination answer, define the construct, describe its precursors, explain the false-belief test, compare the major explanatory models, and finish with clinical interpretation rather than treating one task as a diagnostic instrument.

Must remember: False belief is the examination pivot: the child must predict action from another person's belief even when that belief conflicts with reality and with the child's own knowledge.

9.1 What theory of mind includes, and what it does not

A representational account of minds, not merely a response to behaviour. At the simplest level, the child recognises that people are agents whose actions are directed toward goals. Development then involves distinguishing *desire* from *belief*: two people can want different outcomes, and two people can hold different representations of the same reality. Mature social understanding further requires recognising that access to information determines knowledge, that beliefs may be false, and that outward expression can differ from inner emotion. These strands do not have to develop at identical rates, which is why theory of mind is better considered a family of related abilities than a unitary faculty.

Cognitive theory of mind concerns beliefs, intentions and knowledge. **Affective theory of mind** concerns inference about another person's feelings. **Empathy** overlaps with the second but is not identical to it: affective resonance, concern and the motivation to help are separable from the ability to identify what someone feels. **Joint attention** is also distinct. It is the shared coordination of attention between a person and an object or event, and is a developmental foundation for learning what another person is attending to; it does not by itself demonstrate understanding of false belief. Likewise, mirror self-recognition concerns representation of the self, not direct proof that the child represents another person's mind.

■ **RULE** **Do not equate precursor, correlate and criterion.** Joint attention, pointing, pretend play, language and executive control support theory of mind. The classic explicit test is false-belief reasoning. None of these observations alone diagnoses or excludes a neurodevelopmental disorder.

9.2 Developmental sequence: from shared attention to represented belief

The sequence is more reliable than a rigid age ladder.

1. Early face-to-face reciprocity and turn-taking make interaction predictable.
2. The infant then coordinates gaze and attention with another person, follows what the other person is attending to, and uses gestures such as pointing to request or share interest.
3. Symbolic and pretend play show that one object or action can stand for another, an essential representational step.
4. The child subsequently reasons about desires, perception and access to knowledge, then understands that belief can diverge from reality.

MNEMONIC

"Share - Symbolise - Believe"

Shared attention follows what another person is attending to; symbolic and pretend play show that one object or action can stand for another; then the child understands that belief can diverge from reality.

Conversation with caregivers and peers supplies repeated evidence that different people know, want, remember and feel different things.

8 to 12 months JOINT ATTENTION ACCURACY	Around 12 months JOINT ATTENTION AND POINTING	18 months of age SYMBOLIC PLAY	Four or five years EXPLICIT FALSE-BELIEF TASK PERFORMANCE
--	--	--	--

Berk describes joint attention as becoming more accurate across 8 to 12 months. This is a developmental description, not a referral cutoff. The *New Oxford Textbook of Psychiatry* describes joint attention and pointing as expected in most typically developing children around 12 months. Dulcan describes symbolic play as expected at 18 months of age. *Developmental Psychopathology* places its expected appearance at approximately **20 months of age**. These are source-specific descriptions, not interchangeable thresholds. *Psychotherapy for Borderline Personality* describes explicit false-belief task performance as expected at four or five years. Berk also reports performance after **three and a half years** and describes a span between **four and six years**, but those statements do not establish referral thresholds.

The source language matters clinically. An expected attainment age describes what usually occurs; it is not an upper limit and it is not a red-flag threshold. The range reported by Berk is also not permission to refer every child outside an average-looking point. Development must be interpreted longitudinally and in context, especially when task language, hearing, schooling or cultural familiarity may alter performance.

9.3 False-belief tasks: the classic demonstration

The test asks whether the child can suppress reality and answer from another mind. In the classic change-of-location task, a protagonist puts an object in one location and leaves. In the protagonist's absence, the object is moved. The child is asked where the protagonist will look on returning. The reality-based answer is the new location; the belief-based answer is the original location, because that is where the protagonist falsely believes the object remains. A correct response therefore requires the child to represent the protagonist's informational access separately from the child's own knowledge.

The unexpected-contents task uses a familiar container whose actual contents violate its appearance. After discovering the contents, the child is asked what another, uninformed person will think is inside. Related paradigms test appearance versus reality, knowledge from seeing, hidden emotion, deception, irony and social faux pas. **First-order false belief** has the form, "She thinks X." **Second-order false belief** embeds one mind within another, "He thinks that she thinks X." No verified numeric attainment age for the second-order form is supplied here, so it should not be placed on an age axis.

Good testing includes control questions. The examiner checks whether the child remembers the object's original and current locations, follows the story, identifies who saw the transfer, and understands the question. Failure can arise from receptive-language load, working-memory demand, inattention or difficulty inhibiting the true location. Success can also be achieved through learned verbal rules rather than flexible social understanding. Spontaneous behaviour in natural interaction and performance across several tasks are therefore more informative than a single pass or fail.

-
- **TRAP** "The child knows where the object really is" is not the target answer. False-belief tasks deliberately pit the child's true knowledge against the protagonist's outdated belief. Also avoid converting expected performance at four or five years into a diagnostic or referral cutoff.
-

9.4 Competing explanations of theory of mind development

No single account explains every finding.

ACCOUNT	EXPLANATION	QUALIFICATION
Theory-theory account	Proposes that children build and revise an informal causal theory of mind, much as they revise explanations in other knowledge domains. Observed action, testimony and conversational correction provide evidence; conceptual change moves the child from desire-based explanations toward representational belief.	Its strength is developmental change, but the metaphor of a child scientist can understate embodied and relational learning.
Simulation account	Proposes that the child predicts another person's action by using their own mind as a model, imaginatively placing themselves in the other's situation and adjusting for differences. It captures perspective-taking and the intimate relation between self-knowledge and understanding others.	It struggles when the other person's beliefs, motives or emotional responses are very unlike the child's own.
Modular or domain-specific accounts	Propose a specialised mechanism for representing propositional attitudes such as belief. They explain the apparently distinctive problem posed by mental representations and the characteristic social-cognitive profile seen in some neurodevelopmental conditions.	However, a strongly encapsulated module can underplay the effects of language, executive function, teaching and social experience.
Executive-function accounts	Emphasise working memory, cognitive flexibility and inhibitory control. The child must hold two representations in mind and inhibit the salient true state of affairs.	
Language and sociocultural accounts	Emphasise mental-state vocabulary, complement syntax, narrative, sibling and peer discourse, and conversation about reasons for behaviour. Complement constructions such as "She thinks that..." permit one proposition to be represented inside another, mirroring the representational structure demanded by false-belief reasoning.	Association does not establish that language alone causes theory of mind: linguistic competence, executive control and social understanding remain partly dissociable. The most defensible synthesis is interactive: representational competence, executive control, language and social experience constrain one another across development.

Applied to theory of mind, the sociocultural frame directs attention to joint activity, conversation about feelings and reasons.

Responsive mental-state conversation supports theory of mind.

9.5 Psychopathology: delayed, reduced and excessive mentalizing

Clinical interpretation. Theory of mind disturbance is clinically important but diagnostically non-specific. In autism spectrum disorder, whose fuller treatment is in the Intellectual disability and autism spectrum disorder notes, difficulties may involve spontaneous joint attention, integrating contextual cues, inferring intentions or applying explicit knowledge rapidly in everyday interaction. Some verbally able individuals pass standard false-belief tasks through language or learned strategies while continuing to find unstructured social inference effortful. A failed task is not sufficient for diagnosis, and a passed task does not exclude autism.

In schizophrenia and other psychotic states, difficulty may involve both under-attribution and distorted attribution of intention, especially under ambiguity. In personality pathology, particularly borderline presentations, the problem may be unstable or excessive inference about mental states, sometimes termed hypermentalizing, rather than a simple absence of theory of mind. Mood, arousal, trauma, attachment insecurity and interpersonal threat can transiently reduce the accuracy of mentalizing. The useful clinical contrast is therefore hypomentalizing, too little or overly concrete inference; hypermentalizing, elaborate inference unsupported by evidence; and reasonably calibrated mentalizing, which remains tentative and revisable.

Performance is also affected by intellectual functioning, language disorder, attention, executive dysfunction, hearing impairment and limited familiarity with the test format. Deficits can be trait-like, state-dependent or secondary to another cognitive limitation. Theory of mind should therefore contribute to formulation, never substitute for the full developmental and mental-state assessment.

9.6 Clinical assessment and examination synthesis


Assess development, task performance and real-world use separately. Begin with caregiver history of early reciprocity, gesture, joint attention, pretend play, language development, peer relationships, understanding of deception and humour, and any loss of language or social skills. Observe gaze coordination, sharing of interest, response to another person's distress, flexibility during misunderstanding and the person's tendency to check uncertain interpretations. In a formal task, simplify the language without changing the logic, establish that the person understood and remembered the story, and include reality, memory and information-access controls.

Interpret the result against receptive and expressive language, attention, working memory, inhibition, general cognitive level, hearing, educational exposure and cultural-linguistic context. Repeating the same task with translated words is not automatically a valid cultural adaptation. In multilingual Indian settings, establish the language in which the child normally discusses intentions and emotions, and use collateral examples from home and school. When developmental concern is broader than theory of mind, use an appropriate validated developmental screening or assessment pathway, developed in the Child psychiatry: assessment, treatment, services and protection notes, rather than inventing a false-belief referral age.

The short-answer synthesis is compact. Theory of mind is the capacity to represent one's own and others' mental states and predict behaviour from them. Joint attention and symbolic play are precursors; explicit false-belief reasoning is the classic test. Development reflects interacting domain-specific, simulation, executive, language and sociocultural processes. Clinically, failure is non-specific, success can be scaffolded, and the valid unit of assessment is the developing person in context rather than a single task score.

10 . Language acquisition theories (Chomsky, Skinner, Vygotsky)

Language acquisition asks how a child moves from hearing speech in social interaction to understanding and producing a rule-governed, productive language. **Three** classic answers emphasise different causal levels. B. F. Skinner treats verbal behaviour as behaviour shaped by its consequences and controlling context. Noam Chomsky argues that the speed, uniformity and generativity of acquisition require an innate, species-typical capacity for grammar. Lev Vygotsky places language within socially mediated development: speech first coordinates activity between people and is then internalised as a tool for thought and self-regulation. The most defensible contemporary answer is not to declare one winner. Biological preparedness, learning mechanisms and communicative participation operate at different levels and answer different parts of the problem.



■ Learning contingencies

■ Innate grammar

■ Social mediation

■ Productive language

Fig 20.9 Classic accounts of language acquisition work at different causal levels: Skinner emphasises learning contingencies, Chomsky innate grammatical preparedness, and Vygotsky socially mediated speech that becomes internalised.

How to use this page. Learn each theory through **four** questions: *what is acquired, what drives acquisition, what evidence supports it, and what does it fail to explain?* For Skinner, the unit is a functional verbal response and the driver is its learning history. For Chomsky, the central object is grammatical competence and the driver is an innate language faculty constrained by universal principles. For Vygotsky, the unit is mediated social activity and the developmental movement is from shared speech to inner regulation. In a long answer, compare mechanisms rather than writing three disconnected biographies.

Must memorise: Skinner explains language through learning contingencies, Chomsky through innate grammatical preparedness, and Vygotsky through social mediation and internalisation.

10.1 The problem every theory must solve

Children acquire more than a stock of copied words. They learn a sound system, a lexicon, rules for combining words, meanings, conversational intentions and the social conditions under which an utterance is appropriate. They also produce sentences they have never heard, understand that the same form can serve different functions, and make systematic errors that reveal an active rule rather than random failure. Any adequate theory must therefore explain productivity, generativity, comprehension, social use, variation across languages and the transition from assisted participation to increasingly independent communication.

The theories differ partly because they choose different explananda. Skinner asks what environmental variables control a verbal act. Chomsky asks what internal system makes grammatical knowledge possible. Vygotsky asks how culturally organised interaction becomes a psychological tool. A behavioural account can explain why a request occurs in one context without specifying the child's abstract grammar. A grammatical account can explain structural knowledge without explaining why the child chooses a tact, request or reassurance in a particular relationship. A sociocultural account can explain guided participation without, by itself, providing a formal grammar. Apparent contradiction often falls once the level of explanation is named.

3 CAUSAL EMPHASES	1 DEVELOPMENTAL PROBLEM	0 THEORIES SUFFICIENT ALONE
----------------------	----------------------------	--------------------------------

10.2 Skinner: language as learned verbal behaviour

Speech is behaviour whose form and probability are shaped by antecedents and consequences. In Skinner's operant account, the child emits vocal behaviour; caregivers and the wider verbal community selectively respond; useful forms become more likely in relevant contexts. Positive reinforcement, shaping of closer approximations, imitation, prompting, stimulus discrimination and generalisation all contribute. Reinforcement need not mean sweets or praise. Receiving the requested object, attracting attention, continuing a pleasurable exchange or obtaining information can itself strengthen a verbal response.

Skinner classified verbal behaviour by its controlling relation rather than by conventional grammar.

VERBAL OPERANT	CONTROLLING RELATION
A mand	is controlled by a motivating condition and specifies its characteristic consequence, as when a thirsty child requests water.
A tact	is evoked by an object, event or property and maintained by socially generalised consequences, as when the child names a dog.
An echoic response	has point-to-point similarity with heard speech.
An intraverbal	is occasioned by other verbal behaviour without simply reproducing it, as in answering a question or completing a familiar phrase.
Autoclitic behaviour	qualifies or organises other verbal responses, indicating relations such as certainty, negation or attribution.

The same spoken word can therefore belong to different operants depending on what controls it and what consequence follows.

The account is strongest where the learning history is visible: vocabulary taught through joint activity, requests shaped by successful communication, pronunciation altered by feedback, and context-specific use acquired through modelling and consequences. It also gave language research an insistence on observable function. Its central weakness is explanatory reach. Children generate novel utterances, acquire complex structure without explicit reinforcement of every form, and produce rule-based overgeneralisation rather than merely imitating adult speech. Caregivers also respond to meaning, truth and conversational success, not consistently to grammatical form. Conditioning is therefore relevant, but a complete language theory cannot be a catalogue of reinforced sentences.

10.3 Chomsky: an innate capacity for generative grammar

Finite exposure yields knowledge capable of generating indefinitely many sentences.

Chomsky's nativist theory begins from the mismatch between the linguistic input available to the child and the structured grammatical knowledge eventually acquired. The input is variable, incomplete and not organised as a formal lesson, yet children converge on complex rules and recognise relations that cannot be reduced to adjacent word associations. This is the classic **poverty-of-the-stimulus argument**: experience is necessary to select and develop a language, but experience alone appears insufficient to specify every constraint the learner comes to respect.

The traditional examination formulation is an innate **language acquisition device**, a theoretical language-learning capacity that allows the child to extract grammatical structure from speech. In later generative accounts, the more important idea is **universal grammar**: species-typical initial constraints or principles that delimit the kinds of grammar a human learner can acquire. The child is not born knowing English, Hindi, Kannada or any other particular language. The claim is that human biology supplies a structured readiness, while exposure determines which language-specific system develops.

Chomsky also distinguished **competence**, the speaker-hearer's internal knowledge of language, from **performance**, actual language use under limits of attention, memory, motivation and context. A hesitant or incomplete utterance does not necessarily reveal an absent rule. The theory explains productivity, rapid rule extraction and patterned errors more convincingly than simple imitation does. Its limitations are equally important: the proposed innate content is difficult to specify and test independently; formal grammar does not by itself explain vocabulary learning, communicative intention, caregiver-child interaction or cultural variation; and usage-based evidence shows that distributional learning and repeated constructions carry more information than a stark poverty account sometimes assumes.

■ **RULE** **Innate capacity is not innate vocabulary.** Chomsky proposes biological preparedness for acquiring grammatical structure. The particular language, its words and its conventional uses still require exposure and participation.

10.4 Vygotsky: language is socially mediated and then internalised

Language develops in joint activity before it becomes a means of organising the self.

Vygotsky's sociocultural theory treats language as both a cultural sign system and a psychological tool. A more knowledgeable participant directs attention, labels experience, interprets the child's attempts, models a form, asks questions and structures the activity. Through such mediated participation, the child gains not only words but culturally patterned ways of remembering, planning, categorising and solving problems. Acquisition is therefore inseparable from the purposes for which people communicate.

The **zone of proximal development** (ZPD) is the region between what the learner can perform independently and what becomes possible with responsive guidance or collaboration. Language supports learning within this zone through modelling, prompts, recasts, expansion of the child's utterance, shared attention and dialogue. Scaffolding is the later term commonly used for contingent support that is adjusted and gradually withdrawn. It is compatible with Vygotsky, but was not his own term and is not synonymous with the ZPD.

Development proceeds through the following sequence.

1. Social speech, used to coordinate with another person.
2. Private speech, audible self-directed language used to guide action.
3. Condensed inner speech.

Private speech is therefore a transitional instrument of self-regulation rather than meaningless immaturity. Vygotsky's theory is developmental but **not stage-based**. It provides no sequence of Vygotskian stages and no age row for the accompanying theory-alignment plate. Its major strengths are the integration of language, culture, teaching and self-regulation. Its limitations include imprecise measurement of the ZPD, dependence on the skill of the assisting partner, and less specification of biological constraints and formal linguistic structure.

10.5 Direct comparison: what changes the child's language?

Skinner locates change in contingencies. The child repeats and differentiates verbal acts because their consequences vary across situations. The adult is a source of models, prompts and reinforcement. Error is a response not yet adequately shaped or brought under the relevant stimulus control. **Chomsky locates change in the growth of an internally constrained grammar.** Input triggers and informs a biologically prepared system. The adult provides linguistic data, but need not explicitly teach every rule. Error can reveal a productive rule that has been extended too widely. **Vygotsky locates change in participation and internalisation.** The adult or capable peer co-regulates attention and meaning within activity, then transfers responsibility as the child appropriates language for communication and thought.

Their signature predictions also differ. A Skinnerian analysis asks whether changing antecedents and consequences changes the target utterance. A Chomskyan analysis asks whether the child's output displays rule-governed structure not directly copied from the input. A Vygotskian analysis asks how performance changes under calibrated assistance and whether socially organised speech becomes independent self-guidance. These predictions can coexist in one episode. A caregiver may reinforce a request, supply grammatically structured input, and scaffold the child's participation at the same time.

■ **TRAP** Do not write "Skinner equals nurture, Chomsky equals nature, Vygotsky equals both" and stop. A scoring answer names the mechanism: operant contingencies, innate grammatical constraints, and socially mediated internalisation.

10.6 What later evidence adds: interaction, statistics and use

Later approaches weaken the idea that acquisition must be assigned to a single mechanism. Social-interactionist accounts emphasise joint attention, turn-taking, communicative intention and responsive caregiver speech. Usage-based accounts propose that children learn recurring constructions from meaningful use and gradually abstract patterns across exemplars. Statistical learning shows that learners can extract distributional regularities from sequences, while connectionist models demonstrate how graded patterns may emerge without an explicitly represented rule for every construction. These approaches challenge a rigid version of both imitation-only behaviourism and richly pre-specified grammar.

They do not erase the classic theories. Reinforcement and modelling demonstrably alter language behaviour; human neurobiology constrains which communication systems can be acquired and how readily; and language develops in relationships structured by culture. The useful synthesis is biopsychosocial and developmental: biologically prepared learning systems detect structure in socially meaningful input, while interaction supplies motives, feedback, shared reference and opportunities for practice. The relative contribution of each process varies across phonology, grammar, vocabulary, pragmatics and inner speech.

10.7 Clinical and Indian interpretation

A theory is not a milestone chart and cannot diagnose delay. Chomsky, Skinner and Vygotsky explain proposed mechanisms of acquisition; developmental surveillance asks whether

communication skills are emerging as expected and whether a sourced referral signal is present. The accompanying language-milestone plate should be read according to the construct attached to each value: expected attainment, expected range, median, upper limit and red flag are not interchangeable. A child outside an expected description is not automatically disordered, while a sourced red flag warrants further evaluation rather than a theory-based label.

INDIA

Assessment in India must establish the child's exposure and competence across all languages used at home, school and the wider community. Code-switching, language mixing and unequal vocabulary across languages can reflect ordinary multilingual use rather than deficient grammar. Ask who speaks which language with the child, for what purpose, with what consistency, and whether the concern appears across languages. Skinner directs attention to opportunities and consequences for communication; Chomsky prevents low exposure from being mistaken for absent human language capacity; Vygotsky directs attention to family dialogue, siblings, teachers, cultural tools and the child's response to support. None justifies judging competence from performance in the examiner's preferred language alone.

Clinically, use the theories to generate questions, not conclusions. Does the child initiate communication when it produces a meaningful consequence? Can a model, prompt or expansion elicit a more advanced form? Does the child produce novel combinations rather than echoes alone? How does performance change with a familiar communication partner? Is language being used for joint attention, requests, comment, play and self-guidance? Hearing, oral-motor function, cognition, social communication, the language environment and regression must be assessed separately. Loss of language or social skill is a red flag at any age and is never explained away as a preferred theoretical pathway.

10.8 Examination reconstruction

Long-answer sequence. Define language acquisition, state the problem of productivity, then write Skinner as operant learning with verbal operants, Chomsky as innate grammatical preparedness with universal grammar and competence-performance, and Vygotsky as social mediation with the ZPD, private speech and internalisation. Compare the role of input, adult, child and error; add one strength and one limitation of each; end with an interactionist synthesis and the statement that Vygotsky has no age-linked stages.

MNEMONIC

"Consequence - grammar - mediation"

Skinner shapes use, Chomsky constrains structure, Vygotsky transforms shared speech into self-regulation.

11 . Parenting styles (Baumrind) and family influences on development

Parenting style is the relatively stable emotional climate in which caregivers exercise authority, express warmth, communicate expectations and respond to the child. It is broader than a single

parenting practice. A practice is a specific act, such as praising effort, supervising internet use or imposing a consequence; a style is the recurring relational pattern within which such acts acquire meaning. Baumrind's work made this pattern examinable by distinguishing authoritative, authoritarian and permissive parenting. Maccoby and Martin later organised the framework along two dimensions, **responsiveness** and **demandingness**, and separated an uninvolved pattern from permissiveness.

How to use this page. Learn the two dimensions before the four labels. Then connect parenting with development through modelling, reinforcement, attachment, language, emotion socialisation and family systems. In a long answer, define style, draw the two-by-two classification, distinguish each pattern, discuss associated outcomes without claiming causation, and finish with bidirectionality, culture, socioeconomic context and clinical assessment. The central correction is simple: families influence development powerfully, but neither parenting style nor family structure mechanically determines a child's outcome.

The must-memorise line: authoritative parenting combines high responsiveness with high demandingness: warmth without structure is permissive, and structure without responsiveness is authoritarian.




Fig 20.10 The parenting styles placed on responsiveness and demandingness. Fill marks the high-demandingness column and the saffron edge marks the high-responsiveness row, so authoritative and authoritarian are visibly the same column and a different row: they differ on responsiveness alone. The low-low cell is drawn dashed because it is Maccoby and Martin's addition, not one of Baumrind's original three styles. No ages, prevalence figures or effect sizes appear because none were sourced for this chapter.

11.1 From Baumrind's three styles to the two-dimensional model

Baumrind described patterns of parental authority; the familiar four-cell matrix is a later extension. Diana Baumrind distinguished three broad parenting styles from observations of parent-child interaction. **Authoritative parenting** combined firm standards with warmth and communication. **Authoritarian parenting** emphasised obedience and control with relatively little reciprocal discussion. **Permissive parenting** was accepting and non-punitive but placed few consistent demands on the child. Maccoby and Martin subsequently recast the typology as the intersection of responsiveness and demandingness, producing four cells and adding the low-low pattern commonly called **uninvolved** or neglectful parenting.

Responsiveness refers to warmth, acceptance, attunement, reciprocal communication and support for age-appropriate autonomy. It asks whether the caregiver notices and responds to the child's needs while recognising the child as a separate person. Demandingness refers to supervision, maturity demands, behavioural expectations, limit-setting and consistent follow-through. It asks whether the caregiver provides an organised structure and expects the child to participate in it. Neither dimension is intrinsically good at its maximum. Responsiveness without

boundaries can fail to contain behaviour, while demandingness that becomes rigid, humiliating or coercive is not healthy structure.

2 CLASSIFICATION DIMENSIONS: RESPONSIVENESS AND DEMANDINGNESS	3 STYLES IN BAUMRIND'S ORIGINAL TYPOLOGY	4 CELLS IN THE LATER MACCOBY- MARTIN FORMULATION
--	--	--

MNEMONIC

"High responsiveness + high demandingness = authoritative"

Authoritative parenting combines high responsiveness with high demandingness: warmth without structure is permissive, and structure without responsiveness is authoritarian.

11.2 The four parenting styles

Classify the observed pattern on both dimensions, then describe what the caregiver actually does. The labels are useful summaries, but behavioural description prevents them from becoming moral verdicts.

STYLE	RESPONSIVENESS	DEMANDINGNESS	CHARACTERISTIC FAMILY CLIMATE
Authoritative	High	High	Warmth, clear expectations, monitoring, reasons for rules, reciprocal discussion, consistent consequences and increasing autonomy within limits
Authoritarian	Low	High	Obedience, unilateral rules, restricted negotiation, punitive or coercive control, and limited attention to the child's viewpoint
Permissive or indulgent	High	Low	Warmth and acceptance with few limits, weak monitoring, inconsistent follow-through or reluctance to require developmentally appropriate responsibility
Uninvolved	Low	Low	Limited warmth, monitoring and guidance; basic material provision may coexist with emotional disengagement or inconsistent availability

-
- **RULE** **Authoritative means firm and responsive, not merely kind.** It preserves parental authority while making rules predictable, proportionate and open to explanation. The child may be heard without being allowed to decide every limit.
-

Three distinctions carry marks:

Authoritative and authoritarian parenting

are not synonyms: both contain demands, but only the former combines them with responsiveness and reciprocal communication.

Permissive and uninvolved parenting

both have low demandingness, but permissive parenting retains warmth and engagement.

behavioural control from psychological control

Behavioural control, such as supervision and clear rules, differs from psychological control, such as guilt induction, love withdrawal, shaming or intrusion into the child's emotional world. The former can be protective; the latter constrains autonomy and can be harmful even when the household appears orderly.

11.3 Developmental correlates and the problem of causal overclaim

Parenting styles are associated with developmental outcomes, but the associations are probabilistic and context-dependent. Across much of the literature, authoritative parenting is associated with stronger self-regulation, social competence, academic engagement and psychological adjustment. Its proposed advantage is not leniency. Predictable limits reduce uncertainty, warmth preserves connection, explanations help the child internalise rules, and graduated autonomy gives repeated practice in decision-making. These processes can support both competence and relatedness.

Authoritarian parenting is commonly associated with compliant behaviour in the immediate presence of authority but less spontaneous autonomy, lower open communication and, in some settings, more internalising or externalising difficulty. Permissive parenting may preserve closeness but provide insufficient practice in delay, frustration tolerance and responsibility. Uninvolved parenting is associated with broad developmental risk because both relational support and behavioural organisation are reduced. None of these statements diagnoses an individual child or proves that a parenting style caused an outcome. Effects vary with temperament, developmental capacity, caregiver consistency, neighbourhood safety, peer context, culture and the meaning the child assigns to control.

-
- **TRAP** **Do not turn an association into parent-blaming.** Child temperament and behaviour alter parenting, caregiver stress and illness constrain responsiveness, and social adversity can make consistency difficult. A style label describes a pattern in a system; it does not identify a single cause or assign guilt.
-

11.4 Why parenting has effects: six developmental pathways

Parenting influences development through repeated transactions, not through the label itself. **Six** pathways organise the answer:

DEVELOPMENTAL PATHWAY	HOW IT INFLUENCES DEVELOPMENT
Attachment and co-regulation:	sensitive, predictable responses help the child use the caregiver as a source of safety and regulation. Repeated co-regulation contributes to later self-regulation, although attachment is relationship-specific and should not be inferred from a global style label alone.
Modelling:	children observe how adults manage anger, disagreement, affection, work, substances, gender roles and help-seeking. What caregivers enact may teach more strongly than what they instruct.
Reinforcement and discipline:	attention, praise, withdrawal of attention and consequences alter the probability of behaviour. Harsh or inconsistent discipline can create a coercive cycle in which escalating child behaviour and escalating adult control reinforce one another.
Emotion socialisation:	caregivers label emotions, validate or dismiss them, model regulation and teach which expressions are acceptable. Emotion coaching can build recognition and modulation; ridicule or punishment of emotion may produce concealment without regulation.
Cognitive and language stimulation:	conversation, shared attention, play, explanation and responsive expansion of the child's communication provide material for language, executive control and social understanding.
Opportunity and monitoring:	families shape exposure to peers, education, sleep routines, media, risk, responsibility and supportive adults. Monitoring is protective when it is informed and proportionate, but becomes intrusive when it denies developmentally appropriate privacy and autonomy.

These pathways interact. A caregiver who calmly labels distress, maintains a clear limit and helps the child recover is simultaneously providing co-regulation, modelling, language, reinforcement and emotion socialisation. Conversely, chronic conflict can occupy attention, disturb sleep, reduce exploration and make even ordinary demands feel threatening. The developmental effect lies in the recurring pattern and its meaning, not in one isolated exchange.

11.5 The family as a system: bidirectionality, relationships and context

A child develops within a network of relationships whose members continually influence one another. A transactional model rejects a one-way arrow from parent to child. The child's temperament, neurodevelopmental profile, health and behaviour evoke responses from caregivers; those responses alter the child's subsequent behaviour; the new behaviour then changes the caregiver again. An irritable child may meet an exhausted caregiver, escalating both distress and control, whereas the same child may settle with predictable routines and a caregiver who has support. This is an interaction, not proof that either participant is intrinsically difficult.

Family systems theory adds that each dyad is affected by the larger system. The quality of the couple or co-parenting relationship can increase consistency or draw the child into conflict. Triangulation occurs when tension between adults is displaced through the child or the child is recruited into an alliance. Boundaries can become excessively diffuse or rigid, roles can reverse through parentification, and one member can be designated as the problem-bearer for a wider family difficulty. These concepts are hypotheses for assessment, not labels to impose after a single interview.

Family influence extends beyond parents. Siblings provide companionship, conflict, imitation, comparison and practice in negotiation; differential parenting may be as important as average parenting. Grandparents and other caregivers can supply continuity and protection, or create conflicting rules when caregiving authority is unclear. The developmental importance of family structure depends largely on relationship quality, stability, resources and conflict. A single-parent, joint, blended, adoptive or same-sex-parent family is not inherently pathogenic. Structure tells the clinician who is present; process tells the clinician what happens between them.

11.6 Language acquisition: what the family supplies and what it cannot explain alone

The family creates the language-learning environment, but no single language acquisition theory is sufficient. Skinner emphasised learning through imitation, shaping and reinforcement. Caregivers do reinforce communicative success, model words and expand incomplete utterances, but children generate novel sentences and acquire grammar too rapidly for direct reinforcement to be the whole account. Chomsky emphasised an innate biological preparedness for language and the child's capacity to infer an underlying grammar from limited input. This explains productive rule learning but does not make social interaction dispensable.

Vygotsky treated higher mental functions as socially mediated and later internalised. In the zone of proximal development, a child performs with guidance beyond what can yet be done independently; scaffolding is the temporary support that is adjusted and withdrawn as competence grows. Dialogue with caregivers becomes increasingly internalised, including through private speech, and supports planning and self-regulation. Vygotsky proposed a cross-cutting developmental mechanism, not a sequence of age-defined stages, so he should not be forced onto a numeric stage axis.

The useful synthesis is interactionist. Biological preparedness makes language possible; exposure supplies a linguistic system; attention and memory extract regularities; reinforcement shapes communicative effectiveness; and responsive interaction gives words shared reference and social purpose. Family practices matter through turn-taking, joint attention, contingent response, elaboration, mental-state talk and opportunities to speak. Parenting style changes the climate of these exchanges. Authoritative dialogue invites explanation within structure, authoritarian exchanges may privilege commands and obedience, permissive interaction may be rich but inconsistently organised, and uninvolved parenting may provide less contingent communication. These are tendencies, not fixed predictions about any family or language community.

11.7 Theory of mind, false belief and family conversation

Social understanding develops through representing minds as well as observing behaviour.

Theory of mind is the capacity to attribute beliefs, desires, intentions and emotions to oneself and others, and to understand that another person's mental state may differ from one's own and from reality. A standard false-belief task tests whether the child predicts behaviour from an actor's mistaken belief rather than from the child's own knowledge. The cited developmental source describes success on such a task as expected at **four or five years**. This is an expected attainment statement, not a diagnostic or referral threshold.

Family interaction can support theory of mind through conversation about feelings, motives, mistakes, deception and different viewpoints; shared reading and sibling disputes also provide occasions to explain why people act as they do. Language is both a medium for discussing mental states and a cognitive resource for holding competing representations in mind. Performance also depends on comprehension, attention, memory, inhibition, task wording, familiarity and culture. Failure on one false-belief task therefore does not establish a global absence of theory of mind, identify a parenting style or diagnose a disorder.

11.8 Broader family influences on development

Parenting occurs inside social conditions that can protect or constrain the family.

Development is influenced by caregiver mental and physical health, substance use, marital and intergenerational conflict, domestic violence, bereavement, migration, discrimination, poverty, housing, nutrition, educational access, neighbourhood safety and social support. These factors may act directly on the child and indirectly by altering caregiver availability, predictability and emotional reserve. Treating a resource-constrained family as simply inconsistent confuses a social determinant with a character judgement.

Risk is cumulative but not destiny. Stable relationships, a caregiver who can repair after conflict, supportive kin, school connection, safe routines and access to treatment can buffer adversity. Family conflict is particularly harmful when it is intense, chronic, unresolved or draws the child into adult disputes. In contrast, observing disagreement followed by negotiation and repair can teach that relationships survive conflict. The clinically relevant variables are exposure, meaning, persistence, protection and recovery.

Family transmission is also not purely environmental. Parents and children share genetic liabilities, and genetically influenced child characteristics can shape both the environments selected by the family and the responses evoked from others. This does not reduce development to heredity. It explains why an observed association between parenting and child outcome may contain inherited, evocative and environmental pathways at the same time. Developmental formulation should keep all three visible.

11.9 Culture and the Indian family context

The two dimensions travel better across cultures than stereotypes about the four labels.

Responsiveness and demandingness can be described behaviourally in any family, but the meaning of obedience, autonomy, proximity, privacy and interdependence varies. Firm control

may be interpreted differently when it is embedded in warmth, shared obligation and credible concern for safety. This cultural moderation does not make humiliation, violence, terror or coercive psychological control benign. Culture explains meaning; it does not excuse harm.

INDIA

Assessment should identify the actual caregiving unit rather than assuming a nuclear household. Grandparents, older siblings and other relatives may share authority, care and financial responsibility. Educational pressure, respect for elders, family interdependence and collective decision-making may be valued, while autonomy may be expressed through negotiated responsibility rather than early separation. Ask who sets rules, who comforts, who supervises, whether caregivers agree, and how the child understands these practices. Do not label joint-family involvement, close parent-child proximity or deference to elders as pathology without evidence of distress, coercion or impairment. Equally, do not hide abuse or severe restriction behind the phrase "Indian culture." No verified Indian population norms for Baumrind-style prevalence or comparative effect sizes are supplied here, so Western associations must not be relabelled as Indian norms.

11.10 Clinical assessment, formulation and intervention

Translate labels into observable sequences and modifiable processes. Ask each caregiver and the child, when developmentally appropriate, for concrete recent examples of praise, rules, refusal, anger, affection, supervision, privacy, conflict and repair. Establish what happened before the interaction, what each person did, what followed immediately and what happened the next time. Observe tone, turn-taking, intrusiveness, warmth, limit-setting and the child's bids for help or autonomy. Obtain information across home, school and other caregiving settings because a family may be structured in one domain and overwhelmed in another.

A useful formulation has **five** layers:

1. The child's temperament and developmental capacities.
2. The caregiver's style, health and history.
3. Specific parenting practices.
4. Relationships and alliances across the family.
5. The wider cultural, socioeconomic and safety context.

Then identify the feedback loop maintaining the difficulty and the protective processes already present. Screen directly for violence, maltreatment, caregiver mental illness, substance problems and risk to the child rather than allowing a benign style label to obscure safeguarding concerns.

Intervention targets the process that can change. Core elements include predictable routines, developmentally appropriate expectations, clear brief instructions, positive attention for desired behaviour, consistent non-violent consequences, emotion coaching, collaborative problem-solving, graduated autonomy and repair after conflict. Parent-management approaches are especially useful for coercive cycles and inconsistent reinforcement; family work is useful when conflict, alliances or contradictory caregiving maintain the problem; treatment of caregiver

illness and practical social support may be prerequisites for change. The aim is not to manufacture an idealised parent. It is to increase safety, responsiveness, effective structure and the family's capacity to adapt as the child develops.

Final synthesis. Baumrind's framework is a map of the caregiving climate, not a verdict on a family. The strongest answer begins with responsiveness and demandingness, distinguishes the original three styles from the later fourth, explains developmental pathways, and then qualifies every outcome claim with bidirectionality, temperament, culture and social context. Family relationships provide the setting in which language, emotion regulation, theory of mind and autonomy are practised. Their influence is profound precisely because it is repeated, relational and modifiable, not because it is deterministic.

The domains and the milestones

12 . Domains of child development (motor, language, cognitive, social, emotional)

Child development is an ordered but variable acquisition of function across five interacting domains: **gross motor**, **fine motor**, **language and speech**, **cognitive development**, and **social and emotional development**. The domains are separated for examination and screening, not because they develop independently. A child who cannot explore because of a motor impairment receives a different stream of cognitive and social experience; a child with impaired hearing may appear delayed in language, play and social reciprocity; a child exposed to chronic adversity may show effects across attention, learning, affect regulation and relationships. The clinically useful question is therefore not merely, "Has the child acquired the milestone?" It is, "What is the pattern across domains, what is the trajectory, and is there loss, asymmetry or a qualitative abnormality?"


How to use this page. First learn the architecture: the five domains, the principles of developmental progression, and the distinction between a typical age, a normal window and a referral threshold. Then use the milestone, stage-alignment and primitive-reflex plates as reference tools rather than as lists of isolated dates. Finally, learn how normal variation becomes a clinical signal: developmental surveillance detects a trajectory, screening estimates risk, assessment characterises the profile, and diagnosis explains it. The central exam skill is to move from a remembered milestone to a domain-specific interpretation without turning a screening cut into a diagnosis.

Development is a trajectory across domains: loss at any age, marked asymmetry, or a milestone outside a construct-explicit window demands explanation, not a reflex label.

MNEMONIC

"G F L C S: five interacting domains"

Gross motor, Fine motor, Language and speech, Cognitive development, and Social and emotional development are the five interacting domains.



skills constrain later ones, but a child need not advance at the same rate in every domain. A jagged profile may be more informative than a uniformly low profile.

The five-domain formulation prevents two common errors. First, *motor* must be divided into gross motor posture and locomotion, and fine motor grasp, manipulation and visuomotor integration. Second, *language* is not synonymous with speech: language includes receptive and expressive symbolic communication, pragmatics and gesture, whereas speech concerns the motor production and intelligibility of sounds. Cognition includes attention, memory, learning, problem-solving, symbolic representation and executive functions. The social domain concerns reciprocity, attachment, joint attention, play and relationships; the emotional domain concerns expression, recognition and regulation of affect. These last two are often combined as social and emotional development because each is organised within relationships.

■ **RULE** **Always attach an age to its construct.** "Half of children by" states a median; "usually by" states an expected attainment; "normally between" states an expected range; "still normal up to" states an upper limit; and "refer if absent by" states a red flag. One cannot be substituted for another.

A milestone is a conveniently observable marker, not the whole developmental process. Milestones by age are useful only when the source and normative meaning are retained. Many textbook milestone ages are presented without saying whether they are averages, typical attainments or outer limits. Such values can be repeated as the source states them, but they cannot honestly be converted into "when to refer" thresholds. The reference plates therefore distinguish construct-explicit ranges from source-stated ages whose statistical role is unclear.

12.2 Motor development: posture, locomotion, grasp and visuomotor control

Gross motor development builds an antigravity platform; fine motor development uses that platform for exploration. Gross motor examination follows head control, rolling, sitting, transitional movements, standing, gait, running, jumping and stairs. Fine motor examination follows visual regard and reach, grasp pattern, transfer, release, pincer control, block construction, scribbling and copying. Quality matters as much as attainment: persistent fisting, early fixed hand preference, asymmetry, abnormal tone, unusual movement strategies or loss of a skill cannot be reassured away by the presence of another milestone.

The most defensible infancy anchor is the WHO Motor Development Study, which followed children in Ghana, India, Norway, Oman and the USA. Its windows run from the 1st to the 99th centile and are explicitly screening signals, not diagnostic thresholds. The gross-motor milestone plate gives all six windows, from sitting without support through walking alone, with the centile construct visible beside each value. Their breadth is a corrective to the false precision of a single milestone date. They signal the need for screening when a child appears late; they do not by themselves diagnose developmental delay.

4.3% OF CHILDREN IN THE WHO STUDY DID NOT EXHIBIT HANDS-AND-KNEES CRAWLING AT ALL	90% APPROXIMATELY ACHIEVED FIVE GROSS MOTOR MILESTONES IN A COMMON SEQUENCE	5 sites INCLUDED INDIA, MAKING THE WHO WINDOWS DIRECTLY RELEVANT TO INDIAN TEACHING
--	--	--

Crawling is the classic trap. It is common but not universal, so absence of crawling alone is not a developmental red flag. Look instead at the child's overall locomotor strategy, tone, symmetry, transitions and progression towards supported and independent mobility. The full gross-motor milestone plate should be read as a set of windows, not a railway timetable.

Fine motor progression illustrates proximodistal organisation. An early palmar or raking approach becomes increasingly radial and then finger-thumb controlled; voluntary release and construction follow. The fine-motor milestone plate preserves Berk's expected ranges alongside the source-stated ages from Indian and international clinical references whose normative construct is unclear. It carries the longer sequence from grasp and transfer through block construction, scribbling and copying forms. At examination, link a failure to vision, tone, coordination, opportunity and comprehension before calling it an isolated fine-motor delay.

12.3 Language and speech: receptive, expressive, pragmatic and phonological development

Communication begins before words and comprehension generally outruns production. The sequence runs through social attention, reciprocal vocalisation, cooing, babbling, gesture, understanding of familiar words and commands, meaningful single words, word combinations, increasingly grammatical sentences, narrative and conversational competence. **Receptive language** is what the child understands; **expressive language** is what the child communicates through gesture, words and sentences; **pragmatics** is the social use of communication; and speech is the articulation, fluency, voice and motor realisation of language. A child may therefore have normal language with a speech-sound problem, or intact speech production with impaired social communication.

Source-stated expectations provide useful anchors without pretending to be universal deadlines. The language milestone plate separates cooing and babbling, receptive language, gesture, meaningful words, word combinations, vocabulary growth, grammar and intelligibility, and labels whether a value is expected, a range, a median or merely source-stated. One especially examinable construct error concerns Tasman's median expressive vocabulary of **50 words at 18 months** in US toddlers: this means half had reached that number, not that every child below it has delay. Expected intelligibility values in Kaplan and Sadock and DSM-5-TR are likewise not automatic referral thresholds.

The Indian Psychiatric Society clinical reference places babbling at 4 to 5 months, picking up familiar objects on request at 1 year, and two- to three-word sentences at **2 years**, but does not state the normative construct for these ages. Berk, in contrast, gives **18 to 24 months** as an expected range for combining two words. These statements should remain side by side rather than being collapsed into one falsely exact Indian or Western cut-off.

Language assessment begins with hearing and the communication environment, then separates comprehension, expression, gesture, reciprocity, speech clarity and oral-motor function. Ask about the languages used with the child and judge competence across the entire linguistic repertoire. Multilingual exposure does not justify loss of language, absent gesture or poor

reciprocity. Conversely, a smaller vocabulary in each individual language can mislead if the child's total conceptual vocabulary and communicative effectiveness are ignored.

12.4 Cognitive development, language acquisition and theory of mind

Cognition develops through action, representation, language and increasingly organised reasoning. Piaget's account remains the standard examination scaffold.

1. The sensorimotor stage emphasises knowledge through perception and action and the emergence of object permanence.
2. The preoperational stage brings symbolic representation but reasoning remains intuitive and vulnerable to centration and egocentrism.
3. Concrete operations bring conservation, reversibility and logical operations on tangible material.
4. Formal operations permit systematic abstraction and hypothetical reasoning.

The stage-alignment plate gives the source-stated age ranges. The clinical use is not to "diagnose a Piaget stage" but to pitch questions, explanations and testing tasks at the child's representational capacity.

Vygotsky offers a complementary, non-stage account. Cognitive development is socially and culturally mediated; skills first occur between people and are then internalised. The zone of proximal development is the distance between what a learner can do independently and what becomes possible with competent assistance. Scaffolding supplies temporary structure and is withdrawn as mastery grows; private speech helps organise behaviour and becomes internal speech. In assessment, compare unsupported performance with the child's response to graded prompts. The amount and kind of help that unlocks performance may reveal learning potential that a static score misses.

Three broad families explain language acquisition. Skinner's learning account emphasises imitation, shaping and reinforcement of communicative behaviour. Chomsky's nativist account argues that the speed and generativity of grammar require an evolved language capacity rather than reinforcement alone. Social-interactionist accounts, including Vygotsky's, emphasise joint activity, caregiver responsiveness, shared attention and language used within the zone of proximal development. Contemporary clinical reasoning is integrative: biological preparedness, auditory and cognitive capacities, social motivation, exposure and contingent interaction all matter. The practical use is causal restraint. Delayed language is not automatically poor parenting, and a rich language environment does not exclude hearing impairment, neurodevelopmental disorder or oral-motor difficulty.

Theory of mind is the capacity to represent mental states as mental states and to understand that another person may hold a belief different from reality and from one's own. Passing a false-belief task is expected at about **4 or 5 years** in the cited account. Failure on one task is not synonymous with autism: performance also depends on language comprehension, working memory, attention and the exact task demands. Clinically, theory of mind is examined through

pretend play, perspective-taking, deception, humour, social inference and the child's explanation of other people's actions, not through one laboratory item alone.

12.5 Social and emotional development: attachment, play, temperament and parenting

Social-emotional development is regulation within relationships. Early reciprocal attention and the social smile are followed by selective attachment behaviour, wariness of unfamiliar people, separation protest, joint attention, imitation, symbolic play, peer engagement and increasingly reflective emotion regulation. The social-emotional milestone plate sets out the source-stated windows and expected attainments for these behaviours. Observe whether they are spontaneous, shared for enjoyment and integrated with eye gaze, affect and gesture, rather than merely present on command.

The literature gives different expected windows for the normal anxieties of attachment. The plate therefore keeps Lewis, Kaplan and Sadock, and Developmental Psychopathology side by side instead of collapsing them into one cut-off. The examinable principle is qualitative: separation and stranger anxiety reflect discrimination of familiar caregivers and attachment, and become concerning when intensity, persistence or developmental context causes marked impairment.

Thomas and Chess describe **temperament** through nine dimensions: activity level, rhythmicity, approach or withdrawal, adaptability, threshold of responsiveness, intensity of reaction, quality of mood, distractibility, and attention span or persistence. Their three types are easy, difficult, and slow-to-warm-up. These are descriptive constellations, not diagnoses, and many children do not fit a pure type. **Goodness of fit** is the match between the child's temperamental characteristics and the demands, expectations and supports of the environment. A highly active child may function well where movement is permitted and poorly where prolonged stillness is demanded. Clinical intervention can therefore alter the fit without pathologising the child.

Baumrind parenting styles organise caregiving by responsiveness and demandingness.

PARENTING STYLE	PATTERN
Authoritative parenting	Combines warmth with clear, developmentally appropriate limits.
Authoritarian parenting	Is controlling with lower responsiveness.
Permissive parenting	Is responsive with few consistent demands.
The later uninvolved category	Describes low responsiveness and low demandingness.

These patterns are associations, not deterministic labels, and must be interpreted in cultural, socioeconomic and safety context. The clinical use is to translate a style label into modifiable behaviour: predictability, contingent warmth, supervision, repair after conflict and limits that match the child's developmental capacity.

12.6 Stage theories on one axis: psychosocial, psychosexual, moral and relational development

Stage theories ask different questions and should be aligned, not blended. The alignment plate places the source-stated numeric stages on a shared axis while leaving genuinely non-

numeric theories non-numeric.

12.7 Primitive reflexes: emergence, integration and neurological meaning

Primitive reflexes are expected early patterns whose interpretation depends on the specific reflex, the method and the neurological context. The primitive-reflex plate shows appearance, how each reflex is elicited, and the source-stated disappearance. At the bedside assess symmetry, vigour, habituation where relevant, and persistence alongside tone, posture and voluntary movement. Absence when expected, marked asymmetry, or persistence that interferes with voluntary control may indicate central or peripheral dysfunction, but gestational age, illness and state of arousal matter.

The sources do not always give one disappearance age. Berk, Hutchison and Tasman differ for Moro and palmar grasp, while Babinski has separate expected, upper-limit and red-flag statements. The primitive-reflex plate displays these source-specific claims without forcing them into one value. Hutchison also gives a general warning about persistence of primitive reflexes, yet its reflex-specific entries show why the honest answer must name the reflex and construct rather than manufacture a universal cut-off.

■ **TRAP** A disappearance age is not automatically a referral age. Textbooks may report different expected ages for the same reflex. State the source, preserve whether it is expected, an upper limit or a red flag, and interpret the reflex within the neurological examination.

Reflexes should not be recited in isolation from developmental organisation. Their integration permits increasingly selective voluntary movement, but development is not a simple switch from "reflex present" to "reflex absent". The reflex table is a reference for examination; the clinical task is to decide whether the entire motor pattern is coherent.

12.8 Developmental surveillance, red flags and Indian assessment pathways

A concise clinical synthesis. Name the affected and preserved domains; describe onset, sequence, rate, plateau or regression; compare performance with construct-appropriate norms; examine hearing, vision, growth, neurology and dysmorphism; document the relational and learning environment; and choose screening or assessment deliberately. The diagnostic question comes last. Developmental competence is not the accumulation of dates but the organisation of increasingly adaptive function across motor, language, cognitive, social and emotional systems.

13 . Gross motor and fine motor milestones by age

Developmental milestones are landmarks in an unfolding process, not deadlines stamped onto a child. The exam task is therefore twofold: know the age-linked sequence well enough to recognise a delay, and know what kind of age each source is giving you. An **expected range** describes variation in attainment; an **expected age** describes what a typical child usually does; an **upper limit** marks the outer boundary of normal; and a **red-flag threshold** triggers further assessment. Most textbook ages do not identify which of these constructs they represent. Those ages may be repeated in the source's own terms, but they must never be promoted into referral rules.

How to use this page. Begin with the two orienting maps: the **developmental-theory alignment** and the **domains-and-milestones framework**. Then use the accompanying milestone plates to learn sequence and age together. Read gross motor against the construct-explicit WHO windows, read fine motor as source-labelled examination anchors, and finish by asking whether the child is following the expected sequence, whether skills have been lost, and whether screening rather than diagnosis is the next step.

The must-memorise line: an age is not a referral threshold until the source identifies it as an upper limit or a red flag.




Fig 20.12 Motor milestones from two sources, drawn so the kind of number is visible. Outer bars are the WHO Motor Development Study windows between the 1st and 99th centiles, from 816 children in Ghana, India, Norway, Oman and the USA. Inner bars and dots are Berk's ranges and average ages. Where no Berk milestone matches a WHO one exactly, no average is shown rather than pairing an approximate equivalent.

13.1 The twin orienting maps: theories by age and domains and milestones

One map explains the child; the other describes the child's observable development. The developmental-theory alignment places the stage theories beside one another so that the resident can compare what each theory foregrounds at the same period: cognition in Piaget, psychosocial conflict in Erikson, drive organisation in Freud, separation-individuation in Mahler and moral reasoning in Kohlberg. Vygotsky belongs beside, not inside, this age axis because his sociocultural account is not a stage sequence. The second map is the domains-and-milestones framework: gross motor, fine motor, language and speech, social and emotional development, and cognitive or adaptive functioning. Together they prevent a common error, which is to confuse a theory about development with a milestone used to observe it.

The maps also operate at different levels. A **stage theory** supplies a broad organising idea, while a **milestone** is a directly elicited or observed behaviour. Neither is a diagnosis. A child may attain motor skills within a wide normal window while showing difficulty in another domain, and apparent delay may reflect opportunity, illness, sensory impairment or cultural variation rather than a unitary disorder. The developmental framework is therefore always domain-specific before it becomes global.

-
- **RULE** **Separate sequence, age and construct.** First ask whether the skill appears in the expected developmental order; then identify the source's age; finally decide whether that age is an expected range, an upper limit, a red flag or an age whose normative role is unclear.
-

MNEMONIC

"SAC: Sequence, Age, Construct"

First ask whether the skill appears in the expected developmental order (Sequence), then identify the source's age (Age), and finally decide whether that age is an expected range, an upper limit, a red flag or an age whose normative role is unclear (Construct).

13.2 Band A: the stage-theory alignment

The theories by age plate is an alignment device, not a claim that the theories are interchangeable. Read its source-specific age bands horizontally, but keep each theory's question distinct. Piaget's sequence of sensorimotor, preoperational, concrete operational and formal operational stages helps the clinician pitch explanations and tasks to the child's mode of reasoning. Erikson's psychosocial stages, from trust through autonomy, initiative, industry and identity, organise the age-salient relational conflict. Freud's psychosexual stages, oral, anal, phallic, latency and genital, remain historically important for formulation and examination theory. The age bands on the plate are source descriptions whose normative construct is unclear; none defines a clinical threshold.

Mahler's separation-individuation model follows differentiation, practising, rapprochement and object constancy, and is clinically useful when thinking about autonomy in relation to a caregiver. Kohlberg's moral development must not be assigned invented numerical ages: Kaplan and Sadock uses descriptive bands such as before adolescence, during adolescence and late adolescence or early adulthood. Vygotsky proposed no age stages. His zone of proximal development, scaffolding, private speech and socially mediated learning are cross-cutting mechanisms that guide how much support permits the child's next level of performance. The plate therefore shows Kohlberg as descriptive bands and Vygotsky beside the age axis, never as fabricated numeric rows.

13.3 Band B: the five domains and the organisation of motor development

The domains of child development are related but not reducible to one another. **Gross motor** development concerns posture, balance and large antigravity movements; **fine motor** development concerns reach, grasp, release, manipulation and visuomotor construction. The remaining domains are language and speech, social and emotional development, and cognitive or

adaptive functioning. Examination should therefore be domain-specific: delayed independent walking is not equivalent to delayed pincer grasp, and either may occur with preserved language and social reciprocity.

Motor control generally becomes organised in a **cephalocaudal** direction, with control of the head and upper trunk preceding control needed for standing and walking, and in a **proximodistal** direction, with shoulder and arm control preceding refined finger use. These are organising gradients, not clocks. The milestones by age plates convert the gradients into observable behaviours, while the clinician tests quality as well as attainment: symmetry, tone, postural control, movement strategy, persistence of primitive patterns and spontaneous use of both hands.

13.4 Gross motor milestones by age: use WHO windows as screening signals

The WHO Motor Development Study gives the most defensible construct-explicit gross motor framework. It followed 816 children from Ghana, India, Norway, Oman and the USA and derived windows bounded by the 1st and 99th centiles. These are *windows of achievement*, intended to support descriptive population comparisons and to signal the need for screening when a child appears late. They are not diagnostic boundaries. The gross motor plate therefore owns the complete sequence, while the prose here teaches how to interpret it:

- 1. sitting without support;
- 2. standing with assistance;
- 3. hands-and-knees crawling;
- 4. walking with assistance;
- 5. standing alone;
- 6. walking alone.

3.8 to 9.2 months WHO 1ST-TO-99TH-CENTILE WINDOW FOR SITTING WITHOUT SUPPORT	8.2 to 17.6 months WHO 1ST-TO-99TH-CENTILE WINDOW FOR WALKING ALONE	4.3 percent CHILDREN DID NOT SHOW HANDS- AND-KNEES CRAWLING AT ALL
--	--	---

The span between the two anchors matters. Standing alone has the widest WHO window, from 6.9 to 16.9 months, while sitting without support has the narrowest. About **90 percent** of children achieved five of the milestones in a common sequence, but crawling was not universal. India was one of the WHO study sites, so the WHO norms are not simply Western norms imported into Indian practice, although they are still multinational rather than an Indian-only reference.

Source-specific differences must remain visible. For Western norms, Berk reports a U.S. expected range of 5 to 9 months for sitting alone and 9 to 17 months for walking alone. WHO gives **3.8 to 9.2 months** and **8.2 to 17.6 months** respectively. Kaplan and Sadock gives 9 to 15 months as an expected range for independent walking. These figures answer different source questions and should not be averaged into a new norm.

■ **TRAP** **Failure to crawl is not, by itself, a red flag.** WHO found that 4.3 percent of children never exhibited hands-and-knees crawling. The examination error is to convert a non-universal milestone into a compulsory gateway to walking.

13.5 Fine motor milestones by age: grasp, transfer, pincer and construction

Fine motor development is a progression from whole-hand contact to precise opposition and planned construction. The fine motor plate should be learned as a sequence:

SEQUENCE	FINE MOTOR MILESTONE
1	early grasp;
2	purposeful transfer between hands;
3	radial hand use;
4	pincer opposition;
5	spontaneous scribbling;
6	block construction;
7	copying forms.

The child's posture and vision matter throughout; a poor reach may arise from proximal instability, and an apparent manipulation problem may reflect visual or asymmetric motor difficulty.

The source labels must be preserved. The IPS child and adolescent psychiatry guideline lists bidextrous grasp at 3 months, monodextrous grasp and object transfer at 6 months, and pincer grasp at 9 months, but does not state whether these are expected ages, averages or limits. Kaplan and Sadock says the pincer grasp is first seen at about 9 months, again without converting the statement into a threshold. Berk makes the stronger construct-explicit claim that a well-coordinated pincer grasp is expected by the end of the first year. For a U.S. reference range, Berk places vigorous scribbling between 10 and 21 months and building a tower of two cubes between 10 and 19 months.

Tasman supplies the familiar examination sequence for block towers and copied forms, but those entries have an unclear normative construct. They belong on the milestone plate as source-listed anchors, not in a referral algorithm. The same caution applies when different books disagree: Tasman lists copying a cross and a square at 48 months, while the IPS guideline lists copying a cross at 3 years. Record both source claims; do not collapse them, choose a hidden average or describe either as an upper limit.

13.6 Primitive reflexes: appearance, disappearance and the meaning of persistence

Primitive reflexes are normal early motor phenomena whose later persistence may indicate impaired cortical maturation or neurological dysfunction. The accompanying reflex table owns the verified appearance and disappearance ages and should be used during revision to pair each reflex with how it is elicited. In prose, the high-yield principle is that a reflex age remains

source-specific: Moro disappearance is expected at 12 weeks in Tasman but at 6 months in Berk and Hutchison, while asymmetric tonic neck reflex disappearance is expected at 4 months in Berk. The disagreement must be shown rather than silently harmonised.

Hutchison treats persistence of primitive reflexes beyond the fourth month of life as a general red flag, but its individual-reflex statements must still be respected. The extensor plantar response may remain within the upper limit up to 18 months, whereas Babinski persistence beyond 2 years is a red flag. Thus, primitive reflexes are interpreted by reflex, context and source, not by applying one disappearance age indiscriminately to all.

13.7 Step 5: developmental history, screening and Indian clinical use

A milestone concern becomes clinically meaningful only after a developmental history and direct observation. The developmental history establishes pregnancy and birth context, illness, opportunity for movement, the sequence and quality of attainment across all five domains, asymmetry or early hand preference, functional impact, family observations and any loss of previously acquired skills. Loss of language or social skills at any age requires assessment. Developmental screening then identifies children who need fuller evaluation; it does not itself provide an aetiological diagnosis.

WHO norms contribute a multinational motor reference that included an Indian site.

14 . Language and speech developmental milestones

Language development is not a list of words tied to birthdays. It is the progressive organisation of attention, comprehension, symbolic representation, vocabulary, grammar, speech-sound production and social communication. The examination task is to recognise the sequence, separate receptive language from expressive language and speech, preserve the normative meaning of every age, and identify the few absences that are genuine developmental red flags. An expected age describes what a typical child usually does; a median means that half of the reference group has attained the skill; an upper limit marks the outer boundary of normal; and a red-flag threshold warrants further assessment. A textbook age whose construct is unclear must never be converted into a referral rule.

How to use this page. Start with the twin orienting maps, the **developmental-theory alignment** and the **domains-and-milestones framework**. Then follow communication from prelinguistic vocalisation to words, combinations, grammar and intelligibility. Read the accompanying milestones by age plate for the full source-labelled sequence, not as a set of deadlines. Finish with worked cases and retrieval prompts. Throughout, ask four questions:

- what domain is being tested;
- whether the skill is understood or produced;
- what construct the stated age carries; and
- whether the finding requires observation, screening or referral.

The must-memorise line: refer for absent babbling or gestures at **12 months**, absent single words at **16 months**, absent spontaneous two-word phrases at **24 months**, or loss of language or social skills at any age.




Fig 20.13 Language and speech milestones, drawn so that source and construct stay visible. Each row is one source's stated claim: Berk (Development Through the Lifespan), Tasman, Kaplan and Sadock, Lewis, DSM-5-TR, Atkinson and Hilgard, Psychology (5th ed.) and the IPS child and adolescent psychiatry guideline. Filled markers are ages whose normative construct the source states; open markers are ages it leaves unclear. Cooing, babbling and vocabulary size are shown as disagreements rather than averaged, because no source reports an average. The coral band carries the four red flags from Developmental Psychopathology, which are the only referral thresholds here.

14.1 The twin orienting maps: developmental theories and observable milestones

One map explains development; the other records what the child can do. The developmental-theory alignment places major theories by age on a shared axis without pretending that they describe the same process. Piaget's sensorimotor, preoperational, concrete operational and formal operational stages concern the organisation of thought and help the clinician pitch explanations and tasks. Erikson's psychosocial stages concern trust, autonomy, initiative, industry and identity and help frame the child's age-salient relational task. Freud's psychosexual stages, oral, anal, phallic, latency and genital, remain historically important in examination theory and psychodynamic formulation. The accompanying alignment plate owns the source-specific age bands. Their normative construct is unclear, so none is a clinical threshold.

Mahler's separation-individuation theory follows differentiation, practising, rapprochement and object constancy and is useful when formulating autonomy in relation to the caregiver. Its sourced ages remain on the alignment plate because their normative construct is unclear. Kohlberg's moral development must remain in Kaplan and Sadock's descriptive bands: before adolescence, school-aged children, some young adolescents, during adolescence, later in adolescence, and late adolescence or early adulthood. These bands help the clinician examine the structure of moral reasoning, but no numeric age may be manufactured from them. Vygotsky did not propose age stages. His zone of proximal development, scaffolding, private speech and socially mediated learning sit beside the age axis as mechanisms that explain how supported performance can exceed independent performance.

The second orienting map is the domains-and-milestones framework. It divides observable development into gross motor, fine motor, language and speech, social and emotional, and cognitive or adaptive domains. A theory is an explanatory lens; a milestone is an observed or elicited behaviour. The distinction matters clinically: a child may have age-appropriate gross motor and fine motor skills but delayed language, or may speak little because hearing, opportunity or social reciprocity is impaired. The developmental framework is therefore domain-specific before it becomes a global judgement.

■ **RULE** **Name the construct before using the age.** Say "expected" only when the source says typical attainment, "median" only when it identifies the 50th percentile, "upper limit" only for an outer normal boundary, and "red flag" only for a stated referral threshold. Otherwise attribute the age to its source and leave its normative role unclear.

14.2 Band A relational and constitutional frameworks

Attachment, temperament, theory of mind, language-acquisition theory and parenting style influence the communicative environment. In practice, biology, learning and responsive interaction work together.

14.3 Band B: the five domains and the architecture of communication

Language is one developmental domain, but it recruits all the others. The five domains of child development are gross motor, fine motor, language and speech, social and emotional, and cognitive or adaptive functioning. Gross motor development concerns posture and large

movement; fine motor development concerns reach, grasp, manipulation and visuomotor construction. Motor control is commonly organised along cephalocaudal and proximodistal gradients. Language and speech concerns comprehension, symbolic communication, words and grammar, plus the motor production of intelligible sounds. Social and emotional development supplies reciprocity and joint engagement, while cognition supports representation, memory and rule learning.

Within communication, keep four components separate.

COMPONENT	FUNCTION
Receptive language	is what the child understands.
Expressive language	is what the child communicates.
Speech	is the motor and acoustic production of sounds.
Pragmatics	is the socially appropriate use of communication.

Infants understand more than they can produce, so the absence of spoken words does not by itself define the level of comprehension. Assessment must sample response to name, words and commands, gestures and joint attention, spontaneous communication, vocabulary, sentence structure, articulation, voice, fluency and intelligibility. Hearing must be examined because a sensory limitation can mimic or compound language delay.

The milestone framework is sequential rather than rigidly universal. Prelinguistic vocalisation precedes meaningful words; single words precede productive combinations; vocabulary and grammar expand; and speech becomes increasingly intelligible. The accompanying domain-specific plates own the full motor, language and social and emotional milestone reference tables. They should be read horizontally across age and vertically within each domain, without duplicating them into a second prose table.

14.4 From cooing and babbling to words and combinations

Early communication moves from vocal play to symbols shared with another person. Kaplan and Sadock describes cooing as expected at about 3 to 4 months and meaningful words as expected at about 12 months. *Psychology* (5th ed.) describes babbling as expected at about 6 months, while Berk states that babbling includes many sounds of spoken languages by 7 months. These expected claims should be distinguished from source-listed ages whose construct is unclear, such as cooing at 2 months in Berk, canonical babbling around 7 months in Tasman, and first recognisable word at 12 months in Berk.

The Indian clinical reference in the IPS child and adolescent psychiatry guideline lists babbling at 4 to 5 months, understanding "no" at 9 months and picking up familiar objects when asked at 1 year. The source does not say whether these are expected ages, averages or limits. They should therefore be retained as IPS clinical reference points, not labelled Indian population centiles and not converted into referral thresholds. Western and Indian-source statements stay visible side by side because silently averaging them creates a norm that no source reported.

Kaplan and Sadock describes word combinations as expected at 18 months. Berk gives an expected range of 18 to 24 months for combining two words, while the IPS guideline lists 2- to 3-word sentences at 2 years without specifying the normative construct. Lewis describes one-word sentences as expected at 18 months, jargoned speech at 18 months, and three- to four-word sentences by 3 years. The correct examination answer names the behaviour, the source and the construct, then uses the explicit red-flag thresholds for referral rather than treating any one textbook anchor as a deadline.

14.5 Vocabulary, grammar and intelligibility

Vocabulary size varies widely; communicative organisation and intelligibility provide the more useful sequence. Lewis gives expected expressive vocabulary anchors of about five or six words by 12 months, about 10 words at 18 months, nearly 300 words by 30 months, and 500 to 1,000 words by 36 months. Tasman reports a median of 50 words at 18 months among US toddlers, meaning half had attained that vocabulary size by that age. It also reports fewer than 10 words at the 10th percentile and more than 200 at the 90th percentile at 18 months, but does not define a referral cut. These values demonstrate variability and must not be collapsed into a single universal minimum.

about 25 percent SPEECH INTELLIGIBLE TO STRANGERS AT 18 MONTHS	about 50 percent EXPECTED SPEECH INTELLIGIBILITY AT 24 MONTHS	about 75 percent KAPLAN AND SADOCK EXPECTS INTELLIGIBILITY TO STRANGERS AT 36 MONTHS	nearly 100 percent EXPECTED SPEECH INTELLIGIBILITY AND ADULT-LIKE AT 4 YEARS
--	---	--	--

DSM-5-TR says speech is mostly intelligible by 3 years.

Grammar changes alongside vocabulary. **Telegraphic speech**, in which content words carry meaning while smaller grammatical elements are sparse, is expected around a year and a half in *Psychology* (5th ed.). Two- and three-word sentences are expected at about 2 years in Atkinson and Hilgard, and sentences become progressively more grammatical. By about 6 years, children are expected to be nearly as fluent as adults. Fluency, grammar and articulation are related but separable, so a child may have age-appropriate ideas and sentence structure yet remain difficult to understand.

Speech-sound expectations must also preserve their construct. DSM-5-TR states that most speech sounds and words become clear with development, while misarticulation of a single late-acquired English speech sound can remain within normal limits up to 8 years. That is an upper-limit statement for a specific late-acquired sound in English-speaking children, not permission to ignore globally unintelligible speech or impairment in another linguistic context.

■ **TRAP** Do not turn a vocabulary count into a universal referral threshold. At 18 months, Lewis expects about 10 words, while Tasman reports a US median of 50 words and a very wide percentile spread. The four explicit communication red flags, functional impairment, trajectory, comprehension, hearing and social communication decide escalation.

14.6 Primitive reflexes, adolescence and the wider developmental axis

Communication must be interpreted within neurological and maturational development.

Primitive reflexes are expected early neurological responses. The accompanying reflex table owns their verified appearance, method by which each is elicited, and source-specific disappearance. The high-yield lesson is not to harmonise disagreement: Tasman expects Moro disappearance at 12 weeks, whereas Berk and Hutchison give 6 months; Berk expects asymmetric tonic neck reflex disappearance at 4 months. Hutchison treats persistence of primitive reflexes beyond the fourth month of life as a general red flag, but allows an extensor plantar response up to 18 months as an upper limit and treats Babinski persistence beyond 2 years as a red flag. Each reflex must therefore be interpreted individually.

Adolescent development extends the same developmental method beyond childhood. Its tasks include identity formation, autonomy, peer and intimate relationships, values, gender identity and continuing psychosexual development. Puberty describes biological maturation, while Tanner staging records morphology rather than assigning a fixed age. Girls are staged separately for breast and pubic-hair development; boys for genital and pubic-hair development. Stage I is preadolescent and stage V is mature or adult distribution. The verified puberty plate retains sex-specific ranges without inventing Indian timing data.

The neurodevelopment of adolescence is often framed through a dual-systems or maturational imbalance account, in which reward and socioemotional salience can exert strong influence while cognitive control continues to consolidate. This helps explain peer-sensitive risk-taking without treating normal exploration as pathology. Clinically, the resident asks about persistence, context, impairment, mood, substance use, trauma and coercion rather than applying a moral label.

14.7 Step 5: developmental red flags, surveillance and Indian screening

Screen and refer. The practical question is when to refer, and these explicit thresholds answer it more safely than an unlabelled textbook milestone. Arrange comprehensive developmental, hearing and communication assessment when concern persists or a red flag is present.

14.8 Step 6: five worked cases

Worked case 1, milestone-dating: An 18-month-old uses about 10 words, understands familiar requests and has started combining words. Lewis treats about 10 expressive words as expected at this age, Kaplan and Sadock expects word combinations, and Tasman reports a much higher US median of 50 words. The correct conclusion is that the sources describe different constructs and populations. Assess trajectory, comprehension, hearing and function; do not diagnose delay from the median.

Worked case 2, red-flag: A child has no single words at 16 months. This meets a verified red-flag threshold for further developmental evaluation. Take a full developmental history, test hearing, observe gestures, joint attention and reciprocity, screen development, and refer. Waiting for an unlabelled later textbook anchor would misuse the evidence.

Worked case 3, attachment-pattern: In a Strange Situation vignette, a child explores with the caregiver present, is distressed by separation, seeks comfort on reunion and then returns to play.

The pattern is secure attachment because the caregiver functions as a secure base and reunion restores regulation. The answer describes the observed strategy, not the worth of the parent.

Worked case 4, temperament and goodness-of-fit: A slow-to-warm-up child speaks freely at home but becomes silent when repeatedly pressed to perform before unfamiliar adults. Thomas and Chess directs attention to approach or withdrawal, adaptability and intensity. A slower, predictable introduction improves goodness of fit. The theory explains context-sensitive performance without assuming a fixed language deficit.

Worked case 5, theory-application: A child can complete a verbal classification task with graduated prompts but not independently. Vygotsky locates this performance in the zone of proximal development and supports scaffolding followed by withdrawal of help. Piaget asks what kind of reasoning the task demands, while Erikson asks how the encounter supports initiative or industry. The theories illuminate different aspects of one observation and do not replace direct assessment.

14.9 Step 7: consolidation, retrieval and synthesis

MNEMONIC

"Hear, Share, Say, String, Clear."

Hear stands for receptive language and hearing; *Share* for gesture, joint attention and pragmatics; *Say* for meaningful words and vocabulary; *String* for word combinations and grammar; *Clear* for articulation and intelligibility. Run the mnemonic in that order during a developmental history. Add the red-flag ladder separately: 12 for babbling or gestures, 16 for a single word, 24 for a spontaneous pair, and any age for loss.

Synthesis map: redraw from memory two parallel lines. On the first, place the sourced theories by age, keeping Kohlberg as descriptive bands and Vygotsky as a cross-cutting mechanism. On the second, place milestone anchors from cooing and babbling through words, combinations, grammar and intelligibility. Beneath both, write the five domains and the four red flags. This makes the theory-alignment map explain the child while the milestone map records the child's observable function.

Final clinical synthesis. Language assessment is developmental reasoning in miniature: identify the domain, distinguish understanding from production, describe the sequence and trajectory, preserve the source's normative construct, look for regression and social-communication change, examine hearing and the caregiving environment, and use screening to decide who needs formal assessment. The safest resident is not the one who recites the most ages, but the one who knows exactly what each age is allowed to mean.

15 . Social and emotional developmental milestones

Social and emotional development is the progressive organisation of reciprocal engagement, attachment behaviour, shared attention, play, self-awareness, understanding of other minds and regulation of affect within relationships. It is not a single skill and it cannot be judged from eye

contact alone. A child may produce a behaviour on request yet rarely use it spontaneously to share interest; another may show an age-listed behaviour but with poor reciprocity, little affective integration or loss of previously acquired skills. For the psychiatry resident, the decisive task is therefore to combine milestone timing with quality, context, trajectory and function.

How to use this page. Begin with the five domains of child development, then learn the social and emotional sequence from social reciprocity through joint attention, symbolic play, peer engagement and theory of mind. Use the accompanying social and emotional milestone plate for the domain-specific reference table. This page explains what the entries mean, why normal anxiety windows differ across sources, and how to move from milestones by age to developmental surveillance and referral without converting an expected age into a red-flag threshold.

In social and emotional development, timing starts the assessment; spontaneity, reciprocity, integration, trajectory and loss decide its clinical meaning.




Fig 20.14 Social and emotional milestones, drawn one row per sourced claim so that disagreement between texts stays visible. Colour encodes the normative construct each source states for itself: an expected range, an expected age, or an age whose role the source never states. Stranger anxiety is 6 to 8 months in Kaplan and Sadock and around 8 to 12 months in Developmental Psychopathology; separation anxiety is about 6 to 8 months in Lewis and 10 to 18 months in Developmental Psychopathology. Both windows are kept side by side and neither is averaged. No milestone on this plate is a referral threshold; the only two sourced red flags for this domain are boxed separately. The lower panel is on a years axis, not months.

15.1 The domain within the five-domain developmental map

The domains are examined separately but develop as one system. The five domains are gross motor, fine motor, language and speech, cognitive or adaptive development, and social and emotional development.

DOMAIN	CONTRIBUTION
Gross motor development	supplies posture and mobility.
Fine motor development	supplies reach, grasp and manipulation.
Language and speech	supplies comprehension, expression and verbal communication.
Cognition	supplies representation, learning and problem-solving.
Social and emotional development	supplies reciprocity, relationships, play and regulation.

Social behaviour therefore recruits every other domain. A child must be able to perceive a social cue, orient and act on it, understand its meaning, and regulate the resulting arousal.

The general motor principles of cephalocaudal and proximodistal progression explain why early postural control and proximal stability support later exploration and gesture, but they are not laws for dating social milestones. Social-emotional progression is better understood as widening organisation: regulation is initially supported by a caregiver, dyadic engagement becomes shared engagement around an object or event, symbolic play permits representation, peer interaction becomes more coordinated, and the child increasingly understands mental states. These are linked developmental achievements, not isolated party tricks.

- **RULE** **Name the construct whenever you name an age.** An expected age says what usually occurs; an expected range describes a normal window; a source-stated age with an unclear construct must remain source-stated; and a red flag says when absence warrants further evaluation. None can silently substitute for another.

The accompanying plate owns the full domain-specific social and emotional milestone reference table. Read it horizontally to see the child's profile across functions and vertically to follow one function over time. Where two sources give different windows, retain both. Where a source does not identify its value as expected, median, upper limit or red flag, do not promote that value into a clinical cut-off.

15.2 Early reciprocity and the social smile

The earliest useful observation is not whether the infant smiles, but whether the smile is socially organised. A **social smile** is elicited in reciprocal interaction and is integrated with orientation, facial affect and the other person's behaviour. Berk gives an expected range of **6 to 10 weeks**; *Developmental Psychopathology* describes it as expected at around 2 months in typically developing infants. These are compatible expected statements, not referral thresholds. Tasman lists 2 to 3 months and *Companion to Psychiatric Studies* lists 8 weeks, but neither source states the normative construct, so those values should be repeated only in the source's own descriptive role.

During examination, create an opportunity for mutual engagement rather than commanding a performance. Observe orientation to the caregiver's face and voice, change in affect with

interaction, reciprocal vocal or facial response, ease of soothing and recovery after arousal. A single smile is weak evidence if it is not socially directed; limited eye contact is also nonspecific when considered alone. Vision, hearing, illness, fatigue, temperament, unfamiliar surroundings and the caregiver-child interaction can all change what appears in the room. The correct mental status description is behavioural and present tense, while the developmental history establishes whether the pattern is typical, delayed, inconsistent or lost.

15.3 Selective relationships: stranger anxiety and separation anxiety

Wariness and protest can be normal signs that the infant now discriminates familiar relationships. **Stranger anxiety** refers to wariness or distress in response to an unfamiliar person. Kaplan and Sadock gives an expected range of 6 to 8 months, whereas *Developmental Psychopathology* gives an expected range of 8 to 12 months. Lewis describes about 8 months and about 24 months, and *Companion to Psychiatric Studies* lists 10 months, but those source statements have an unclear normative construct. The expected ranges must remain side by side; averaging them would create a norm reported by neither source.

Separation anxiety is distress associated with separation from a familiar caregiver. Lewis gives an expected range of about 6 to 8 months, while *Developmental Psychopathology* gives an expected range of 10 to 18 months. Lewis also describes about 14 to 18 months and Berk describes 6 to 15 months, but those two statements do not specify their normative role. This source variation is itself examinable: the behaviour is developmentally intelligible, but the literature does not provide one universal date that separates normal from abnormal.

■ **TRAP** **Normal anxiety is not defined by one birthday.** Do not diagnose from the mere presence or absence of stranger or separation anxiety at one visit, and do not merge differing source windows. Judge intensity, persistence, context, recovery with the caregiver, functional impairment and the wider developmental profile.

Clinically, observe whether the caregiver functions as a source of regulation and whether distress settles sufficiently for exploration to resume. The form of the response matters more than its theatrical intensity. A tired infant in an unfamiliar clinic may protest strongly; a quiet infant may freeze rather than cry. The resident should document the trigger, proximity-seeking, consolability, recovery, exploration and consistency across settings before inferring a relational pattern.

15.4 Joint attention and pointing: from dyadic to shared engagement

Joint attention turns a two-person exchange into a shared mental focus on something in the world. **Joint attention** requires coordinated attention between child, another person and an object or event. It includes following another person's gaze or point and actively directing another person's attention for sharing, not merely to obtain an object. The *New Oxford Textbook of Psychiatry* states that joint attention is expected around **12 months** in most typically developing children. Berk describes increasing accuracy between 8 and 12 months, but does not specify the normative construct for that age band.

Pointing is likewise expected around 12 months in most typically developing children according to the *New Oxford Textbook of Psychiatry*; Tasman lists 14 months without defining whether that is an expected age, average or limit. The important distinction is functional. Imperative pointing requests something; declarative pointing shares interest for its own social value. Assessment should therefore sample gaze-following, showing, giving, pointing, shifting gaze between object and person, and checking the adult's response. A hand movement without coordinated gaze or shared affect is not equivalent to spontaneous declarative communication.

Because social communication and language develop together, ask how the child communicates wants, enjoyment and surprise before relying on spoken vocabulary. Hearing and vision must be considered. So must opportunity: the meaning of a behaviour depends on whether it appears spontaneously across familiar settings, not merely after repeated prompting in a clinic. Around 12 months is an expected attainment statement for joint attention and pointing, whereas absence of gesturing at 12 months is the separately sourced red-flag statement. The identical number does not make the constructs interchangeable.

15.5 Symbolic play, self-recognition and peer sociability

Play reveals representation, social reciprocity and flexibility in a natural behavioural sample.

Symbolic play, also called pretend play, uses an action or object to stand for something else.

Dulcan describes symbolic play as expected at **18 months**; *Developmental Psychopathology* describes it as expected at approximately 20 months in most children. Kaplan and Sadock lists symbolic play at 18 months and older and at 30 months and older in clinical play descriptions, but those statements do not identify a normative construct. The expected values belong on the milestone plate as separate source claims, not as a new averaged deadline.

Examine play for initiation, imitation, shared enjoyment, symbolic substitution, sequencing, flexibility and inclusion of another person. Functional use of an object is not yet symbolic use; repetitive manipulation is not equivalent to pretend play; and reproducing a prompted action once is weaker evidence than spontaneous flexible play. The play sample should be interpreted with fine motor ability, language comprehension, cognitive level, sensory profile, familiarity with the materials and prior opportunity.

Self-recognition in the mirror test is reported by Berk in children older than 18 to 20 months, but the source does not say whether this is an average, range or limit. Keep it as a source-stated observation. For peer sociability, *Companion to Psychiatric Studies* lists cooperative play at 3 years, while Berk describes both parallel play and cooperative play across 2.5 to 6 years. Their normative constructs are unclear. These figures describe patterns in the cited sources and must not be treated as referral deadlines.

Parallel and cooperative play are also not mutually exclusive stages through which every child marches on a fixed timetable. Ask what the child actually does with peers: watches, imitates, plays alongside, exchanges objects, negotiates roles, sustains a shared theme, tolerates frustration and repairs conflict. Context matters. Solitary play in a new clinic cannot establish the child's usual

peer competence, while school reports and direct observation in familiar settings may reveal a very different profile.

15.6 Understanding minds and regulating emotion

Later social competence depends increasingly on representing internal states rather than responding only to visible behaviour. **Theory of mind** is the capacity to understand that

another person can have thoughts, intentions, desires and beliefs different from one's own. Success on a standard false-belief task is expected at **4 or 5 years** in the cited source. Berk reports performance after 3.5 years and between 4 and 6 years but does not identify the normative construct of those values. Failure on a single task is not a diagnosis because language comprehension, attention, working memory and task demands contribute to performance.

Emotional development includes recognition and expression of affect, use of another person's signals, tolerance of arousal, recovery after frustration and an increasing capacity to regulate behaviour in context. The clinically useful sequence is qualitative: regulation begins in interaction with a caregiver and becomes progressively internalised, flexible and responsive to social rules. No verified numeric ages for specific emotion-labelling or self-regulation milestones were supplied, so none should be invented. Assessment instead compares behaviour across home, school and clinic, noting triggers, strategies, recovery, caregiver support and impairment.

Social cognition and emotion regulation must be interpreted culturally. Eye gaze, proximity, independence, deference, emotional display and adult-child conversation vary with family and community expectations. Culture does not explain away regression, marked functional impairment or absence of basic communicative gesture, but it changes the form in which competence is expressed. A sound answer describes the observed behaviour first, then considers developmental level, language, sensory capacity, temperament, context and culture before drawing a clinical inference.

15.7 Step 5: spotting deviation from normal and deciding when to refer

Developmental red flags are construct-explicit escalation signals, not retrospective labels.

The verified social-communication warning signs are absence of gesturing at **12 months** and loss of language or social skills at any age. Both warrant further developmental evaluation. Because language and social communication are tightly coupled, absence of babbling at 12 months, absence of single words at 16 months and absence of spontaneous two-word phrases at 24 months are additional communication red flags that should be checked in the same assessment. A red flag indicates when to refer; it does not establish aetiology or diagnosis.

Spotting deviation requires more than threshold checking. Look for regression, plateau, marked unevenness across domains, lack of spontaneous reciprocity, poor integration of gaze, gesture and affect, unusual rigidity or repetitive play, weak generalisation across settings, and functional impact on learning or relationships. Only those two social-communication findings above are verified referral thresholds in the supplied evidence. Expected ages for the social smile, joint attention, pointing or symbolic play must not be relabelled as red flags merely because the child has passed them.

A focused developmental history records pregnancy and birth context, health, hearing and vision, opportunities for interaction, caregiver concerns, the sequence of attainment across gross motor, fine motor, language and speech, cognition, and social and emotional development, plus any loss of skills. Observe the child with the caregiver and during free play. Sample response to social bids, shared enjoyment, gesture, joint attention, imitation, symbolic play, flexibility, regulation and recovery. Seek reports across settings because a short clinic encounter can under-sample both strengths and difficulties.

Developmental surveillance is the longitudinal clinical process of eliciting concerns, observing development and following trajectory. **Developmental screening** uses a standardised tool to identify children who need comprehensive assessment. Screening is not diagnosis. Refer promptly for comprehensive developmental assessment when a verified red flag is present, when skills are lost, or when persistent concern and functional impairment remain despite an apparently reassuring single observation. Hearing and vision assessment are integral where communication or social response is in question.

15.8 Indian screening context without invented Indian milestones

Indian practice has validated screening instruments, but the available evidence does not supply an Indian social-emotional milestone table. The Trivandrum Development Screening Chart, developed at the Child Development Centre in Kerala, covers 0 to 6 years and contains 51 items. In the validation study by Nair and colleagues, one item delay was the operational positive result called TDSC delay, with DDST as the reference standard. This is a screening rule for the instrument as a whole, not permission to reconstruct individual item ages or to convert a textbook social milestone into a TDSC cut.

1,183 CHILDREN IN THE TDSC VALIDATION SAMPLE	84.62% SENSITIVITY, WITH A 95 PERCENT CONFIDENCE INTERVAL FROM 71.92 TO 93.12	90.8% SPECIFICITY, WITH A 95 PERCENT CONFIDENCE INTERVAL FROM 88.97 TO 92.43	99.23% NEGATIVE PREDICTIVE VALUE, WITH A 95 PERCENT CONFIDENCE INTERVAL FROM 98.48 TO 99.67
--	---	--	--

The DASII yields separate Motor DQ and Mental DQ values, and Indian follow-up practice uses a DQ below 85 as a delay marker. That marker is instrument-level evidence, not a milestone age. The Baroda Development Screening Test was India's first developmental screening tool, introduced in 1991. These instruments help organise referral after developmental surveillance raises concern; they do not replace a social history, direct observation, sensory assessment or a fuller developmental evaluation.

No verified Indian population ages for the social smile, stranger anxiety, separation anxiety, joint attention, pointing, symbolic play, peer play or false-belief performance were supplied. The Western and international textbook statements on the milestone plate therefore remain labelled by source and population where known. They must not be relabelled as Indian norms. This is especially important in viva answers: name the source construct, describe the behaviour and use the validated Indian screening pathway without claiming an Indian centile that has not been published in the material available here.

15.9 Clinical application and retrieval

- **Worked vignette 1, expected age versus red flag:** A caregiver reports no communicative gestures, including no pointing, showing or giving, at 12 months. Pointing itself is expected around 12 months in the *New Oxford Textbook of Psychiatry*, while absence of gesturing at 12 months is separately sourced as a warning sign. Clarify gaze shifts and shared enjoyment as part of the wider social-communication profile. Because the red-flag construct is met, arrange further developmental evaluation; do not diagnose from that finding alone.
- **Worked vignette 2, source variation:** An infant becomes wary of an unfamiliar examiner. Kaplan and Sadock gives a stranger-anxiety range of 6 to 8 months, while *Developmental Psychopathology* gives 8 to 12 months. The correct answer preserves both expected ranges and interprets the behaviour through familiarity, intensity, consolability and recovery. It does not average the ranges and does not call ordinary wariness pathological.
- **Worked vignette 3, play quality:** A child reproduces feeding a doll after repeated demonstration but does not initiate pretend actions, share enjoyment or vary the play. The observed imitation is real, but it does not by itself establish spontaneous flexible symbolic play. Sample familiar materials and caregiver interaction, assess language, cognition and fine motor ability, and obtain behaviour across settings before interpreting the finding.

MNEMONIC

"smile, selective relationships, share, symbol, self and minds"

Smile recalls early reciprocity. Selective relationships recalls stranger and separation anxiety. Share recalls joint attention and pointing. Symbol recalls pretend and peer play. Self recalls self-recognition. Minds recalls false-belief understanding.

Then add the clinical spine: five domains, spontaneous quality, trajectory, loss, developmental surveillance, screening and referral.

Final clinical synthesis. Social and emotional milestones are best understood as an expanding capacity to engage, share, represent, relate and regulate. The safe examination answer names the behaviour, gives the source's normative construct, preserves population and source differences, describes quality and context, looks for loss and functional impairment, and uses construct-explicit warning signs to decide when to refer. Recitation earns a mark; disciplined interpretation prevents harm.

16 . Primitive reflexes and their disappearance

Primitive reflexes are stereotyped, involuntary motor responses elicited by a defined sensory or positional stimulus in the immature nervous system. Their orderly appearance in the newborn and subsequent disappearance reflect intact lower neural pathways followed by increasing supraspinal control. They are therefore neither ordinary voluntary milestones nor isolated curiosities. In an examination, each reflex must be judged by four questions:

1. Is it present when it should be?

2. Is it elicited correctly?
3. Is it symmetrical?
4. Has it been inhibited when expected?

The accompanying reflex table is the visual reference for appearance and disappearance ages. Use it as a source-labelled map, because several respected textbooks give different disappearance points for the same reflex. The safe clinical answer is not to average discrepant ages. State the reflex, demonstrate how it is elicited, quote the named source or range, and distinguish an expected disappearance from an upper limit or a red-flag threshold.

Must memorise: A primitive reflex is interpreted by expected presence, adequate elicitation, symmetry and timely disappearance, never by presence or absence alone.

MNEMONIC

"Present, elicit, symmetry, disappear"

Interpret a primitive reflex by whether it is present when it should be, elicited correctly, symmetrical and inhibited when expected.

Primitive reflexes: elicitation, appearance, disappearance

Persistence beyond the disappearance age is the sign, not the reflex itself. Read one reflex across its row. Do not collapse the table into a single disappearance rule.

Reflex	How it is elicited, and the response	Appears	Disappears, by named source
Rooting	Stroke the cheek or corner of the mouth. The neonate turns toward it, opens the mouth.	● neonate Morgan	● Berk: 3 weeks, becomes voluntary turning ● Neuroanatomy atlas: about 1 year of age
Sucking	Contact with the lips or palate produces rhythmic sucking.	● neonate Kaplan and Sadock	● Berk: after 4 months ● Neuroanatomy atlas: about 1 year of age
Moro	Infant fully supported, a brief controlled head-position change. Never an actual drop.	● Neonatal period, first 4 weeks Tasman	● Tasman: 12 weeks ● Berk and Hutchison: 6 months
Asymmetric tonic neck	Turn the supine infant's head to one side. Face side extends, occiput side flexes: fencer.	○ 4 weeks Tasman	● Berk: 4 months
Palmar grasp	A finger laid across the palm, gentle pressure. The fingers flex around it.	● neonate Kaplan and Sadock	● Hutchison: 2 months ● Berk: 3 to 4 months
Plantar grasp	Pressure at the base of the toes produces toe flexion.	Not sourced No age in the cited texts.	● Hutchison: 2 months
Stepping	Held upright, soles on a firm surface, the body inclined forward. Alternating steps.	● Neonatal period, first 4 weeks Tasman	● Hutchison: 2 months ○ Berk: 2 months, in relation to weight gain
Babinski extensor plantar	Stroke the lateral sole, heel to forefoot, then medially. Great toe extends, toes fan.	● newborns New Oxford	● Berk: 8 to 12 months ● New Oxford: during the first 2 years of life ● Hutchison: extensor plantar up to 18 months

Five of the eight reflexes carry more than one disappearance age. Both claims are shown. Never average them into a false consensus.

Rooting, sucking, Moro and palmar grasp genuinely disagree between texts. Babinski's four boundaries are complementary: a range, a broader range, an upper limit, a red flag.

Red flags. These three are the only ones sourced in this chapter, and all three come from Hutchison's Clinical Methods.

Sucking reflex absent beyond the 36th week of gestation.

Primitive reflexes persisting beyond the fourth month of life.

Babinski reflex persisting beyond the age of 2 years.

The fourth-month statement is a general screening prompt. It does not override the later reflex-specific expectations above for Moro, sucking and the extensor plantar response.

- expected: a typical infant, per the source
- expected range: the normal window
- upper limit: the outer bound of normal
- construct not stated by the source: teach its own words, never convert it into a referral threshold

Every age is quoted from a named text and stays attached to it. Sources: Hutchison's Clinical Methods, Berk's Development Through the Lifespan, Tasman, Kaplan and Sadock, the New Oxford Textbook of Psychiatry, an atlas of neuroanatomy, and Morgan's Introduction to Psychology. Elicitation follows this chapter's text.

Not sourced and therefore not shown: an appearance age for plantar grasp, and any Indian normative reflex schedule. No Western age is relabelled as an Indian norm.

Fig 20.15 The primitive reflexes, with each age kept attached to the text that states it. Disappearance ages come from Hutchison's Clinical Methods, Berk's Development Through the Lifespan, Tasman, and the New Oxford Textbook of Psychiatry; appearance ages from Kaplan and Sadock, Tasman, the New Oxford Textbook of Psychiatry and Morgan. Dots mark what kind of number each age is, because an expected age, an upper limit and a red flag are not interchangeable. Where texts disagree, both claims are shown rather than averaged. The three red flags are the only ones sourced.

16.1 What primitive reflexes signify

They test maturation as well as pathway integrity. A primitive reflex is produced by a relatively fixed sensorimotor circuit that is clinically visible while **descending inhibition** is immature. With maturation, the response is inhibited, incorporated into voluntary movement or replaced by a more purposeful response. Thus, normal development has two examinable directions: an early reflex should be obtainable at the relevant stage, and it should later recede. Persistence, asymmetry, failure to elicit an expected response, or re-emergence after it has disappeared can each be abnormal, but they point to different problems.

Presence alone does not certify a normal nervous system. A response may be weak because the infant is drowsy, recently fed, unwell or poorly positioned; it may be absent because the stimulus was inadequate; and it may be asymmetric because of a peripheral injury even when central maturation is otherwise appropriate. Conversely, persistence suggests deficient inhibition only after the reflex-specific expectation and the rest of the neurologic examination are considered.

16.2 Examination technique: make the response interpretable

Standardise the infant's state before interpreting the movement. Examine in a warm, safe setting, preferably when the infant is awake and settled. Observe spontaneous posture and

movement first. Then apply one clear stimulus, support the head and trunk throughout, compare the two sides where the response permits, and describe the actual movement rather than merely writing “reflex present”. Repeat only when necessary, because habituation, crying and repeated stimulation can alter the response. Record gestational context and the infant's behavioural state.

■ **RULE** **First prove that the test was valid.** Name the stimulus, the motor response, its symmetry and the infant's state. Only then compare the finding with the expected disappearance age.

The usual bedside sequence is efficient: test the orofacial responses, then upper-limb grasp, the startle and postural responses, stepping, plantar grasp and the plantar response. This reduces unnecessary repositioning and makes side-to-side comparison easier.

16.3 Disappearance ages are source-specific constructs
Do not convert a textbook age into a referral threshold.

CONSTRUCT	MEANING
An “ expected ” disappearance	describes what the source presents as usual.
An “ expected range ”	describes a window.
An “ upper limit ”	marks the outer boundary of normal.
A “ red flag ”	is the point at which persistence or absence war-rants concern.

These constructs are not interchangeable. If two books differ, retain both claims with attribution rather than creating a false consensus.

■ **TRAP** “All primitive reflexes must disappear by the fourth month” is unsafe. *Hutchison's Clinical Methods* gives persistence beyond **the fourth month of life** as a general red flag, but the same source and other textbooks give later reflex-specific expectations for Moro, sucking and the extensor plantar response. Use the general statement as a screening prompt, not as a universal cut-off that overrides the named reflex.

the fourth month of life GENERAL RED FLAG	36th week of gestation ABSENCE RED FLAG	18 months of age UPPER LIMIT	2 years RED FLAG
---	---	--	----------------------------

The disagreement often reflects differences in elicitation, the strength counted as “present”, whether a response has become voluntary, and the population or edition being described. The accompanying plate therefore keeps each age attached to its source. It should be read horizontally, one reflex at a time, not collapsed into a single disappearance rule.

16.4 Rooting and sucking: feeding-related primitive responses
Rooting is elicited by stroking the cheek or the corner of the mouth; the neonate turns toward the stimulus and opens the mouth. Its appearance is described in the neonate. Berk and a neuroanatomy atlas give markedly different disappearance points, shown with their attributions in the accompanying reflex table. These values should not be averaged. The difference may reflect

distinct operational boundaries between a reflexive turn and a residual feeding response, so the source must travel with the age.

Sucking is elicited by contact with the lips or palate and produces rhythmic sucking. Kaplan and Sadock describe its appearance in the neonate. *Hutchison's Clinical Methods* treats absence beyond the **36th week of gestation** as a red flag. Berk and the neuroanatomy atlas again give different expected disappearance points; retain both as source-specific values from the plate. The clinically important distinction is between an absent feeding response when it should be established and persistence of a reflexive pattern after voluntary feeding control develops.

16.5 Moro and asymmetric tonic neck reflex

Moro reflex is tested with the infant fully supported: a brief, controlled change in head position produces abduction and extension of the upper limbs, followed by adduction and flexion. The manoeuvre is a **supported head-extension stimulus**, never an actual drop. Tasman describes Moro as present in the neonatal period and gives an earlier disappearance point than Berk and *Hutchison's Clinical Methods*. The plate preserves both claims. Quote the source rather than turning either value into an unsupported universal limit.

For the asymmetric tonic neck reflex, turn the supine infant's head gently to one side. The limbs on the face side extend and those on the occipital side flex, producing the “fencer” posture. Tasman records its early presence without defining the normative construct; this must not be restated as a referral threshold. Berk supplies an expected disappearance point in the plate. Marked obligatory persistence can interfere with midline hand use and rolling, while asymmetry requires a broader neurologic and musculoskeletal assessment.

16.6 Grasp and stepping responses

Palmar grasp is elicited by placing a finger across the palm and applying gentle pressure; the fingers flex around it. Kaplan and Sadock describe appearance in the neonate. Berk gives an expected disappearance range, whereas *Hutchison's Clinical Methods* gives a different expected point; both remain source-labelled in the plate. For plantar grasp, pressure at the base of the toes produces toe flexion. Hutchison supplies its expected disappearance point, but no appearance age was available from the cited sources.

Stepping is elicited by supporting the infant upright, allowing the soles to contact a firm surface and inclining the body slightly forward; alternating step-like movements follow. Tasman describes it as expected in the neonatal period. Hutchison gives an expected disappearance point. Berk also describes disappearance in relation to weight gain but does not define the normative role of that statement, so it should not be converted into an upper limit or referral threshold. The exact source-specific entries belong to the accompanying plate.

16.7 Babinski response: expected early, concerning when persistent

The Babinski reflex, or **extensor plantar response**, is elicited by stroking the lateral sole from heel toward the forefoot and then medially. Extension of the great toe, often with fanning of the other toes, is expected in newborns according to the *New Oxford Textbook of Psychiatry*. It is

commonly taught beside the primitive reflexes because its interpretation changes with nervous-system maturation.

The sources offer complementary, not identical, boundaries, and the plate keeps them separate. Berk supplies an expected disappearance range, the *New Oxford Textbook of Psychiatry* gives a broader expected range, and Hutchison states that an extensor plantar response may occur up to **18 months of age**, an upper limit, while persistence beyond **2 years** is a red flag. Therefore an extensor response at the end of infancy is not automatically an upper motor neuron sign, whereas persistence beyond the sourced red-flag boundary requires neurologic assessment.

16.8 Clinical interpretation: pattern before label

Abnormality lies in the pattern. Bilateral absence of a response expected in a newborn raises questions about gestational maturity, arousal, systemic illness, medication effect or diffuse neurologic dysfunction. A consistently asymmetric Moro or grasp response raises a different set of possibilities, including a focal central lesion, peripheral nerve injury or painful limb movement. Persistence of several primitive responses, particularly with abnormal tone, posture or delayed voluntary motor control, is more concerning for impaired **central inhibition** than a single borderline response.

Re-emergence later in life is described as a frontal release sign. Grasping, rooting, snout and palmomental responses may appear with frontal or diffuse cerebral dysfunction, including neurocognitive disorders, but one isolated sign has limited specificity. In psychiatry, these signs strengthen a neurologic formulation only when they fit the cognitive examination, functional decline and the rest of the neurologic findings.

INDIA

The cited disappearance values are textbook schedules, not Indian normative percentiles. Do not manufacture an India-specific age conversion. In an Indian infant, retain the named source and interpret the response with gestational context, longitudinal development and the complete neurologic examination.

In a short case or viva, the highest-scoring formulation is compact: “I elicited the reflex with the appropriate stimulus; the response was present or absent, graded by quality and symmetry; I compared it with the source-specific disappearance expectation; and I interpreted it with tone, posture, voluntary movement and the rest of development.” That answer demonstrates technique, developmental reasoning and restraint. It avoids the two common errors: treating every age as a hard cut-off and diagnosing neurologic disease from one reflex in isolation.

17 . Adolescent development and identity formation

Adolescence is the developmental transition from childhood dependence toward adult roles, marked by pubertal change, reorganisation of cognition and emotion, increasing importance of peers and intimate relationships, and the construction of a more coherent identity. It is neither a prolonged childhood nor an incomplete adulthood. Development is asynchronous: physical

maturation, reward sensitivity, abstract reasoning, self-regulation, autonomy and social opportunity do not move in lockstep. The psychiatrist therefore judges the adolescent against developmental trajectory, pubertal status, context and functioning, not against a single stereotype of maturity.

How to use this page. Build the answer in **four** layers. Begin with puberty and Tanner staging, then explain the neurodevelopmental basis of context-sensitive risk-taking, then describe identity formation through Erikson and Marcia, and finally integrate gender, sexuality, peers, family and culture. Keep **three** distinctions intact: Tanner stage is based on physical characteristics rather than chronological age; exploration is not itself pathology; and gender identity, gender expression and sexual orientation are separate constructs.

The must-memorise line: Adolescence is an asynchronous transition in body, brain and social role, while identity is formed through exploration and commitment within family, peer and cultural relationships.

-
- **RULE** **The psychiatrist therefore judges the adolescent against developmental trajectory, pubertal status, context and functioning, not against a single stereotype of maturity.** Development is asynchronous: physical maturation, reward sensitivity, abstract reasoning, self-regulation, autonomy and social opportunity do not move in lockstep.
-
- **TRAP** **Exploration is not itself pathology.** Normal diversity in tempo and expression is expected, but marked distress, sustained impairment, coercion, regression or a clear episodic change requires assessment rather than automatic attribution to adolescence.
-




Fig 20.16 The adolescent dual-systems and maturational-imbalance model, drawn qualitatively. The curves carry the model's claim, that a fast-changing socioemotional system runs ahead of a more slowly stabilising cognitive-control system, and that the gap between them is the window in which risk-taking is developmentally expected rather than pathological. The shape is schematic: no source supplied a maturation curve, peak age, crossover age or rate for either system, so none is drawn and the axis carries no ages. The only sourced timing shown is pubertal, from Berk's Child Development, 8th edition, as expected ranges.

17.1 A developmental frame: simultaneous change across interacting systems

The central principle is uneven but coordinated development. Puberty changes the body, sleep, affective salience and how other people respond to the young person. Cognitive development expands hypothetical, future-oriented and self-reflective thought. Social development shifts toward greater privacy, peer affiliation, reciprocal intimacy and negotiation of authority. Identity development integrates bodily experience, values, relationships, education, work, culture, gender and sexuality into an account of who the person is and where life is going. Change in one system alters every other system: earlier or later pubertal change can affect body image and peer position, while family response can amplify or buffer its psychological impact. Developmental tasks are therefore better treated as linked processes than as a checklist. The adolescent must:

- incorporate a changing body; become more autonomous without losing secure relatedness;

- form increasingly mutual peer relationships; develop values and future goals; regulate strong affect;
- make decisions under social pressure; establish a workable sense of self.

Progress is non-linear. A person may reason abstractly in a calm classroom yet decide impulsively in an emotionally charged group, appear independent with peers yet seek parental regulation during illness, or explore one identity domain while remaining committed in another.

The useful clinical model is **transactional**. Biological changes alter behaviour and social signalling; family, school, peers, media and culture shape the opportunities and meanings attached to those changes; repeated experiences then modify self-concept and regulation. Neither biology nor environment alone explains the outcome. Normal diversity in tempo and expression is expected, but marked distress, sustained impairment, coercion, regression or a clear episodic change requires assessment rather than automatic attribution to adolescence.

17.2 Puberty: biological maturation, variable timing and psychological meaning

Puberty is the biological process of sexual maturation, whereas adolescence is the broader psychological and social transition. Gonadal maturation produces secondary sexual characteristics and reproductive capacity; adrenarche is a related but distinct process involving adrenal androgen effects. The order and tempo of visible changes matter more clinically than a single calendar age. Pubertal events also acquire personal and social meaning through comparison with peers, cultural expectations, gender experience, body ideals, family communication and the adolescent's readiness for change.

Psychological effects depend on timing relative to peers and on context, not merely on whether a value falls inside a range. Ask about the adolescent's understanding of changes, body satisfaction, teasing, sexualisation, sports or school consequences, privacy, menstrual experience where relevant, and whether physical development is congruent with gender experience. Puberty may increase visibility and social expectations before cognitive or emotional capacities have developed to the same degree.

17.3 Cognitive and moral development: expanding capacity with uneven performance

Adolescent thought increasingly permits abstraction, hypothesis, metacognition and projection into possible futures. In Piaget's scheme this is **formal operational thought**: reasoning about propositions and possibilities rather than only concrete events. It supports scientific reasoning, ideological debate, imagining alternative selves and considering long-term consequences. It also enables heightened self-consciousness because the adolescent can represent how others might be evaluating them. Formal reasoning is not used uniformly; familiarity, education, emotion and task demands strongly influence performance.

Two classic descriptions help explain adolescent egocentrism. The **imaginary audience** is the sense of being continuously observed or judged, which can intensify concern about appearance and embarrassment. The **personal fable** is a felt uniqueness of one's experience and, at times, a sense that ordinary consequences do not fully apply. Neither construct should be used to

caricature all adolescents. Both are hypotheses about self-focused social cognition and must be distinguished from delusional conviction.

Kohlberg's moral development should be taught in sequence without invented numeric ages. Kaplan and Sadock describes:

LEVEL	KAPLAN AND SADOCK DESCRIPTION
Preconventional morality	before adolescence
Some second-level reasoning	in young adolescents
Third-level conventional reasoning	during adolescence
Fourth-level conventional reasoning	later in adolescence
Fifth-level postconventional reasoning	in late adolescence or early adulthood.

Moral reasoning on a dilemma is not identical to moral conduct: emotion, relationships, incentives, culture and power affect what a person actually does.

17.4 Erikson and Marcia: identity as exploration and commitment

Erikson locates the central psychosocial task of adolescence in **identity versus role confusion**. Identity is a workable sense of continuity and sameness across time, combined with a social recognition that one is the person one understands oneself to be. It integrates earlier identifications with present capacities and future roles. Role confusion refers to difficulty organising these elements into a sufficiently coherent direction, not ordinary uncertainty during exploration. Successful development yields fidelity, the capacity to sustain commitments while recognising ideological and interpersonal difference.

James Marcia operationalised identity through two processes: **exploration**, the active consideration of alternatives, and **commitment**, an investment in chosen goals, values or roles. Their combinations produce four **identity statuses**:

IDENTITY STATUS	DESCRIPTION
Identity diffusion	neither active exploration nor stable commitment.
Foreclosure	commitment adopted without meaningful exploration, often from family or authority expectations.
Moratorium	active exploration without a settled commitment.
Identity achievement	commitment formed after meaningful exploration.

four
IDENTITY STATUSES

These are statuses, not fixed personality types and not an irreversible staircase. The same adolescent may be achieved in religion, foreclosed in vocation, in moratorium regarding relationships and diffuse about politics. Status can change when education, migration, conflict, discrimination, illness or new relationships reopen a domain. Moratorium can be

developmentally productive, while commitment itself is not proof of maturity if it was coerced or cannot be revised in response to experience.

17.5 Identity develops in relationships, culture and digital spaces

Healthy individuation changes the form of attachment rather than requiring emotional separation from the family.

The adolescent seeks greater privacy, agency and reciprocal negotiation while still using caregivers as sources of security, values and practical support. Conflict may increase around boundaries, education, relationships, technology and appearance, but conflict alone does not demonstrate dysfunction. The clinically useful questions are whether disagreement can be negotiated, whether the adolescent has a voice, whether caregivers remain available, and whether autonomy is expanding in a way that is safe and culturally intelligible.

Peers provide comparison, belonging, rehearsal of adult-like intimacy and feedback about possible selves. Close friendships become more reciprocal and self-disclosing; peer groups can support exploration but can also enforce conformity. Social exclusion, bullying and minority stress can become incorporated into self-concept. Digital spaces extend these processes through persistent audiences, curated self-presentation, rapid feedback and access to identity communities. Online identity is not necessarily false; it may be experimental, aspirational, fragmented or more authentic than the identity permitted offline.

INDIA

In Indian practice, identity may be negotiated within strong family interdependence, educational competition, linguistic and religious communities, caste and class location, migration, gendered expectations and collective decisions about work or relationships. Do not equate autonomy with Western-style separation or assume that family commitment means foreclosure. Ask what the commitment means to this adolescent, whether alternatives were thinkable, how much choice was available and what consequences followed dissent. Assess the young person privately as well as in the family context, while explaining confidentiality and its safety limits.

17.6 Gender identity and psychosexual development

Gender identity, gender expression and sexual orientation answer different questions.

Gender identity

is the person's internal sense of gender.

Gender expression

is how gender is communicated through appearance, behaviour, name or social role.

Sexual orientation

concerns patterns of romantic or sexual attraction.

None can be inferred reliably from the others, from Tanner stage, from play or clothing, or from a single sexual experience. Exploration or nonconformity is not in itself a disorder.

Psychosexual development during adolescence includes integration of the sexually maturing body, desire, fantasy, attraction, values, consent, boundaries, behaviour, intimacy and relationship expectations. Pubertal capacity does not equal emotional readiness, informed decision-making or

consent. History-taking should separate attraction, identity, behaviour and distress because they may not align, particularly where stigma or safety concerns limit disclosure. Ask neutrally about relationships, coercion, online contact, contraception and infection protection where clinically relevant, without assuming heterosexuality or sexual activity.

In Freud's historical stage scheme, the genital stage is placed in adolescence. This remains examinable as theory but is not a contemporary developmental norm and does not provide a clinical threshold. Current assessment focuses on the adolescent's lived development, bodily experience, relationships, safety and functioning. Distress associated with gender incongruence must be distinguished from distress produced by rejection, bullying, family conflict or discrimination. A respectful interview uses the adolescent's stated name and pronouns, establishes what can safely be shared with caregivers, and assesses mood, self-harm risk and support without treating identity diversity as pathology.

17.7 Clinical assessment: normal exploration, developmental vulnerability and disorder

The assessment unit is the adolescent in a developmental system. Interview the adolescent privately for at least part of the assessment, interview caregivers, and obtain school or other collateral information when indicated. Explain confidentiality at the outset and state the circumstances in which safety requires disclosure. Establish pubertal trajectory and body experience, sleep, nutrition, cognition and school function, family relationships, peers, romantic and sexual development, gender experience, online life, substances, risk-taking, values, future plans and identity commitments.

For any concerning behaviour, reconstruct the sequence: developmental baseline, onset, triggers, context, frequency, persistence, degree of control, consequences and recovery. Ask whether it occurs mainly in hot peer contexts or also in calm solitary settings; whether it is exploratory, compulsive, coerced, intoxication-related or driven by elevated mood, depression, anxiety, trauma or neurodevelopmental difficulty; and whether it produces sustained impairment. Episodicity, loss of previously stable functioning, pervasive impairment, dangerous consequences, marked distress or a pattern grossly discordant with context weighs against a purely normative explanation.

Pubertal assessment requires the same construct discipline. Record Tanner morphology and source-stated expected ranges, but do not diagnose from an endpoint that was not validated as a clinical cut-off. When timing, sequence or physical findings raise concern, undertake appropriate paediatric or endocrine evaluation rather than converting an expected range into a referral rule. Identity assessment likewise avoids premature closure: uncertainty may be normal exploration, a stable commitment may be healthy or coerced, and apparent confusion may reflect unsafe disclosure conditions rather than lack of identity.

17.8 Examination synthesis and retrieval spine

A complete long answer moves from maturation to meaning. Define adolescence as a broad biological, psychological and social transition. Describe puberty and Tanner staging, making clear that stages are morphological and components are scored separately. Explain dual systems and

maturational imbalance as context-sensitive accounts of risk-taking. Add cognitive and moral development, then make identity formation the centre of the answer through Erikson's identity versus role confusion and Marcia's exploration-commitment statuses. Finish with family, peers, culture, gender, psychosexual development and clinical differentiation from disorder.

MNEMONIC

"body, brain, bond, belief and becoming"

Body recalls puberty and Tanner morphology. Brain recalls control, reward and hot-context decisions. Bond recalls family, peers and intimacy. Belief recalls abstract thought, values and moral reasoning. Becoming recalls identity exploration, commitment, gender and future roles.

Final clinical synthesis. Normal adolescent development is organised variability, not uniform instability. The safe psychiatric formulation links bodily maturation, neurodevelopment, relationships and culture to the adolescent's pattern of exploration and commitment. It respects diversity, preserves confidentiality, identifies coercion and impairment, and avoids turning either an expected age range or an identity difference into a diagnosis.

18 . Puberty, neurodevelopment of adolescence and risk-taking behaviour

Adolescence is a developmental transition, not merely an age band. Three processes run together but are not interchangeable: **puberty**, the biological maturation of the reproductive system and secondary sexual characteristics; **adolescent neurodevelopment**, the continuing reorganisation of brain structure, connectivity and function; and psychosocial development, in which the young person negotiates autonomy, identity, sexuality, peer belonging and adult roles. Their partial mismatch explains much of normal adolescent behaviour. Reward, emotion and social evaluation can acquire unusual motivational force while cognitive control is still becoming more consistent across situations. The result is not global irrationality. It is a context-sensitive period in which judgement may be good in calm conditions and less reliable under arousal, peer observation, immediate reward or threat.

How to use this page. Keep five distinctions clean. Puberty is not the whole of adolescence. Gonadarche is not adrenarche. Tanner stage describes physical maturation and does not assign a chronological age. Capacity for abstract reasoning does not guarantee self-regulation under emotional load. Finally, risk-taking is neither inevitable nor uniformly pathological: some exploration is developmentally useful, but its probability and consequences depend on the person, the peer group, the family, the opportunity and the behaviour involved.

MNEMONIC

"Body, brain, context"

Puberty is the biological maturation of the reproductive system and secondary sexual characteristics; adolescent neurodevelopment is the continuing reorganisation of brain structure, connectivity and function; and psychosocial development is the negotiation of autonomy, identity, sexuality, peer belonging and adult roles. Their partial mismatch explains much of normal adolescent behaviour.




Fig 20.17 Tanner staging as three separate five-stage scales, after Tasman Psychiatry. The descriptors are typeset from Tasman's own stage criteria rather than reproduced as an illustration. No age is attached to any stage, because no source in this build maps one; the single Tanner-stage-tagged age that is sourced is named in the band below the grid, with the source's silence about what kind of number it is.

18.1 Puberty: endocrine activation and physical maturation

Puberty is driven by coordinated endocrine change, with gonadal and adrenal maturation proceeding through related but distinct pathways. Reactivation of the hypothalamic-pituitary-gonadal axis produces pulsatile gonadotropin-releasing hormone secretion, pituitary luteinising hormone and follicle-stimulating hormone release, and increasing gonadal steroid production. This **gonadarche** produces maturation of the testes or ovaries, the growth spurt and the development of secondary sexual characteristics. **Adrenarche** is the increase in adrenal androgen production; it contributes particularly to pubic and axillary hair, body odour and skin changes, but it is not evidence by itself that the gonadal axis has matured. Lewis describes adrenarche as beginning at **6 to 8 years**; because that source does not define the value as a normal limit or a referral threshold, it must not be used as either.

The observable sequence matters more than a single calendar age. In girls, breast development, pubic hair, acceleration in linear growth and menarche are related markers but need not advance in exact synchrony. In boys, enlargement of the scrotal-genital complex, pubic hair, penile growth, change in body composition, the growth spurt and spermarche similarly form a sequence rather than one event. Sex steroids also act on bone, muscle, adipose distribution, skin, sleep, sexual motivation and the brain. These biological effects interact with the meaning assigned to bodily change by the adolescent, family and peer culture. Thus an identical physical event can be experienced as welcome maturation, embarrassment, loss of control, gender incongruence or social advantage.

-
- **RULE** **Stage the body; do not guess the stage from age.** Tanner stage is a physical examination description. Chronological age, menarche, height and the adolescent's own report may support the history, but none substitutes for the relevant stage criteria.
-

18.2 Tanner staging: what the five stages actually describe

Tanner staging uses separate five-stage scales for breast development, male genital development and pubic hair. Stage I is preadolescent. In girls, breast stage II is elevation of the breast and papilla as a small mound; stage III adds enlargement of the breast and areola without separation of their contours; stage IV produces a secondary mound of areola and papilla; and stage V is the mature contour, with a projecting nipple and the areola returning to the general breast contour. In boys, genital stage II is marked by scrotal enlargement with altered pink texture; stage III is characterised particularly by penile lengthening; stage IV adds increasing breadth and glans development with darker scrotal skin; and stage V is adult size. These are morphological descriptions, not age labels.

Pubic hair is staged separately in both sexes. Stage I has no pubic hair. Stage II is sparse and lightly or slightly pigmented hair, initially relatively straight. Stage III hair is darker, begins to curl and increases in amount. Stage IV is coarse, curly and adult-like but occupies a smaller area. Stage V has adult distribution, extending to the medial thighs; in girls it forms the adult feminine triangle. Discordance between genital or breast stage and pubic-hair stage is therefore not a contradiction: the scales partly reflect different endocrine processes. The examiner records each component separately rather than averaging them into a synthetic stage.

Use the Tanner reference plate for the visual sequence. Do not attach ages to its rows. Tasman states that menarche occurs within **2 years** after breast development, but the source does not identify that interval as a median, outer limit or referral cut-off; it should be repeated only as Tasman's descriptive statement. Menarche also does not mean that all physical maturation is complete.

Tanner stage is a clinical descriptor, not a judgement of psychological maturity, consent capacity, sexual experience or gender identity. Examination requires explanation, assent or consent as applicable, privacy, respectful exposure limited to what is necessary, and a chaperone according to clinical policy and the adolescent's needs. Self-assessment may support history-taking when

examination is not appropriate, but it is less reliable than a properly conducted clinical examination.

18.3 Pubertal timing: expected ranges, sequence and tempo

Normal timing is broad, and a range is not a diagnosis. Berk's *Child Development*, 8th edition, reports expected ranges for major pubertal events. These values describe windows in the cited population and must not be converted into medians, guarantees or stand-alone referral thresholds.

8 to 13 years GIRLS: BREAST BUDDING AND ONSET OF THE HEIGHT SPURT	10.5 to 14 years GIRLS: MENARCHE	9 to 13.5 years BOYS: TESTICULAR ENLARGEMENT	12 to 16 years BOYS: SPERMARCHE
SEX	PUBERTAL EVENT	EXPECTED RANGE	
Girls	Breast budding and onset of the height spurt	8 to 13 years	
Girls	Peak height spurt	10 to 13.5 years	
Girls	Appearance of pubic hair	8 to 14 years	
Girls	Breast-growth completion	10 to 17 years	
Girls	Menarche	10.5 to 14 years	
Boys	Testicular enlargement	9 to 13.5 years	
Boys	Penile enlargement	10.5 to 14.5 years	
Boys	Onset of the height spurt	10.5 to 16 years	
Boys	Peak height spurt	12.5 to 15.5 years	
Boys	Appearance of pubic hair	10 to 15 years	
Boys	Spermarche	12 to 16 years	
Boys	Completion of penile and testicular growth	12.5 to 16 years	

Interpretation requires **timing**, **sequence** and **tempo**. Timing asks when change begins relative to the reference population; sequence asks whether expected components are appearing in a coherent order; tempo asks how the changes progress over time. A young person at one edge of a wide expected range may still be developing normally. Conversely, an unusual sequence, arrest or very rapid progression can warrant assessment even when one observed event happens within an expected range. Evaluation should integrate growth trajectory, nutrition and chronic illness, medication and substance exposure, neurological and endocrine symptoms, family pattern, psychosocial distress and examination findings. The ranges above are reference information, not permission to dismiss a concerning trajectory.

INDIA

No Indian population-specific pubertal timing ranges were available in the supplied sources. Do not silently relabel the ranges from Berk as Indian norms. In an Indian examination answer, state the source and population limitation, use the physical sequence and longitudinal growth record, and seek locally validated paediatric or endocrine guidance for clinical thresholds.

18.4 The adolescent brain: remodelling, connectivity and context

Adolescent neurodevelopment is network reorganisation, not simple growth of an unfinished brain. Experience-dependent synaptic stabilisation and pruning make frequently used circuits more efficient while reducing redundant connections. Progressive myelination improves the speed and reliability of long-range communication. White-matter organisation, cortical specialisation and integration among prefrontal, striatal, limbic and temporoparietal regions support gains in working memory, planning, response inhibition, error monitoring, future simulation, social cognition and emotion regulation. These processes do not mature in lockstep, and performance varies with task demands and emotional state.

The prefrontal cortex contributes to planning, rule maintenance, inhibition and valuation of future consequences. The ventral striatum and related dopaminergic circuits participate in reward anticipation, reinforcement learning, novelty and action selection. The amygdala and broader limbic circuitry help assign emotional salience, while medial prefrontal and temporoparietal systems contribute to self-referential and social inference. Development changes the balance and connectivity among these systems. Pubertal hormones can influence motivational and affective processing, but puberty and neural maturation are neither identical processes nor reducible to one hormone level.

A crucial distinction is between **cold cognition** and **hot cognition**. Cold cognition is reasoning under low arousal, with time to deliberate and little immediate social pressure. Hot cognition occurs when reward, threat, sexual interest, shame, anger or peers are salient. An adolescent may explain consequences accurately in the clinic and still choose differently in the heat of the real situation. The discrepancy does not prove deceit or incapacity. It shows that knowledge of risk and regulation of behaviour are separable functions.

-
- **TRAP** Do not teach adolescence as a defective-adult brain with a single completion age. The examinable account is uneven, experience-sensitive network maturation. Adolescents can reason competently, especially in calm settings; vulnerability rises when immediate reward, emotion and peer evaluation load the system.
-

18.5 Dual-systems and maturational-imbalance models

The **dual-systems** family of models explains risk as a temporary imbalance between **motivational drive and regulatory control**. One system is often described as socioemotional or incentive-processing, involving reward-sensitive striatal and limbic circuits that become highly responsive to novelty, immediate gain and social evaluation. The other is a cognitive-control system, relying substantially on prefrontal and frontoparietal networks that support inhibition, planning and resistance to distraction. When incentive salience is high and control is not

consistently recruited, behaviour can shift toward immediate reward despite an intact verbal understanding of later costs.

The **maturational-imbalance model** makes the same broad point while emphasising developmental differences in the timing and coordination of subcortical affective systems and cortical control systems. Peer presence matters because it can increase the subjective value of reward and social approval rather than merely "pressure" the adolescent through explicit persuasion. A choice that looks reckless to an adult may therefore be locally rational within the adolescent's current valuation: the social reward is immediate, visible and identity-relevant, while the harm is delayed and probabilistic.

These models are useful, but they are not complete theories of every adolescent. They do not imply that all adolescents take more risks, that the brain mechanically causes misconduct, or that control always loses to reward. Risk depends on trait impulsivity and sensation seeking, prior adversity, mood and substance use, sleep and arousal, family monitoring, peer norms, opportunity, expected benefit and the severity and reversibility of possible harm. Nor do the models erase agency. They identify situations in which scaffolding, delay, environmental design and supportive adults can make self-control easier to deploy.

Must remember: adolescent risk-taking is best understood as *context-sensitive motivational imbalance*, not absence of knowledge, absence of control or inevitable recklessness.

18.6 Gender identity and psychosexual development

Sex characteristics, gender identity, gender expression and sexual orientation are distinct constructs.

- Sex characteristics include chromosomal, gonadal, hormonal and anatomical features.
- Gender identity is the person's internal sense of gender.
- Gender expression is how gender is communicated through appearance, behaviour and social role.
- Sexual orientation concerns patterns of romantic or sexual attraction.

None can be inferred reliably from the others, and gender-nonconforming expression is not by itself psychopathology.

Puberty can make these domains more salient because the body changes visibly, sexual interest may intensify, peers classify one another more sharply and future relational roles become imaginable. For some adolescents, bodily change is congruent with identity; for others it produces or intensifies distress. The psychiatrist's task is not to force an identity conclusion. It is to clarify the adolescent's own experience over time, the degree of distress and impairment, body-related discomfort, family and peer response, bullying or violence, mood and self-harm risk, and the presence of coexisting neurodevelopmental or psychiatric needs without treating those needs as explanations that invalidate gender experience.

Psychosexual development includes integration of the sexually maturing body, attraction, fantasy, intimacy, values, consent, boundaries and responsibility. Freud's genital stage is historically examinable and is placed in adolescence, but it is not an adequate contemporary account of sexuality. A clinical formulation should instead be developmental, non-judgemental and safeguarding-aware. Ask in private, use the adolescent's terms, distinguish consensual exploration from coercion or exploitation, assess digital as well as in-person contexts, and explain confidentiality and its safety limits before eliciting sensitive information.

18.7 Risk-taking: adaptive exploration, hazardous exposure and individual difference

Risk-taking is any choice made under uncertainty in which an outcome of value is accompanied by a meaningful possibility of harm. It is broader than impulsivity. Impulsivity is poorly controlled action with insufficient forethought; sensation seeking is preference for novel, intense experience; risk-taking may be planned or spontaneous and may involve social, physical, sexual, financial, academic or legal consequences. Substance use, dangerous driving, unsafe sexual behaviour, aggression, self-harm challenges and online exposure are clinically important examples, but so are positive risks such as trying a difficult skill, leaving a familiar peer group or speaking publicly. Development requires exploration; prevention should reduce catastrophic and irreversible harm without attempting to abolish autonomy.

Risk rises when several layers converge. At the individual level consider reward sensitivity, impulsivity, attention and executive function, mood elevation or depression, trauma responses, substance intoxication, sleep loss and beliefs about invulnerability. At the interpersonal level consider peer norms, desire for status, romantic context, family conflict, supervision and modelling. At the environmental level consider access, anonymity, commercial persuasion, unsafe roads, weapons or substances, and online algorithms that amplify novelty and social comparison. Protective factors include a warm relationship with at least one dependable adult, clear and proportionate limits, prosocial peers, school connection, meaningful roles, future goals, emotion-regulation skills and practical barriers to the most dangerous opportunities.

The intervention follows the mechanism. Information alone is weak when the adolescent already knows the hazard. Use collaborative motivational interviewing, identify the reward the behaviour provides, rehearse implementation intentions, reduce access to lethal or irreversible means, create an exit plan for peer situations, move decisions out of high-arousal contexts and enlist supportive adults with the adolescent's participation whenever safety permits. Harm reduction and autonomy-supportive boundaries are often more developmentally effective than shame, surveillance without relationship or an abstract lecture about consequences.

18.8 Clinical assessment and the examination answer

Assess the adolescent confidentially, developmentally and systemically. Begin with the adolescent and caregiver together, then interview the adolescent alone for part of the consultation. Explain confidentiality and its limits in plain language before asking about risk. A HEEADSSS-style review keeps the field broad: home; education or employment; eating; activities and peers; drugs; sexuality; suicide, self-harm and mood; and safety. Add sleep, online

behaviour, bullying, violence, legal exposure, strengths and future goals. Obtain collateral information where necessary, but preserve a private space so disclosure is possible.

The formulation should connect the behaviour to development rather than merely name it. Describe pubertal status and trajectory, cognitive and neurodevelopmental strengths or vulnerabilities, identity tasks, emotional state, the immediate reward, peer and family context, access to means, previous consequences, protective factors and readiness to change. Separate experimentation from a persistent syndrome, and separate chronological age from decision-specific capacity. Capacity is functional: the young person must understand, retain, use or weigh relevant information and communicate a choice in the actual context being assessed.

For a long-answer question, use a disciplined sequence: define adolescence and puberty; describe gonadarche, adrenarche and Tanner staging; state the sourced timing ranges with their population limitation; explain pruning, myelination and changing frontostriatal-limbic connectivity; compare dual-systems and maturational-imbalance accounts; integrate identity, gender and psychosexual development; then finish with determinants, protective factors and clinical management of risk-taking. The highest-scoring conclusion is balanced: adolescence brings increased opportunity for hazardous behaviour, but also exceptional learning, flexibility, social sensitivity and capacity for growth. Developmental vulnerability is real without being destiny.

19 . Gender identity and psychosexual development in childhood

Gender identity and psychosexual development emerge through the continuing interaction of bodily maturation, cognition, attachment, learning, family relationships, peers and culture. They are not single milestones reached on a fixed birthday. The child gradually makes sense of bodily sex characteristics and social categories, develops an internal sense of gender, learns how gender is expressed and regulated in the surrounding culture, acquires privacy and interpersonal boundaries, and later integrates puberty, attraction, values, intimacy and consent into a more coherent identity. Variation is expected. The clinical task is to distinguish diversity and exploration from distress, impairment, coercion, trauma and disorder without making any one form of play, dress, friendship or attraction carry more meaning than it can support.

How to use this page. Begin by separating the constructs that examination questions often collapse. Then describe the major explanatory models, give Freud as a historical stage theory, outline expected sexual learning and boundary acquisition, and connect puberty and Tanner staging to adolescent identity. Finish with adolescent neurodevelopment, risk-taking and a developmentally informed clinical assessment. Throughout, hold **two** disciplines: Tanner stage is morphological rather than chronological, and gender expression or sexual behaviour cannot by itself establish gender identity, sexual orientation, abuse or psychopathology.

The must-memorise line: Separate body, identity, expression, attraction and behaviour, then interpret each against developmental context, consent, distress, impairment and safety.

19.1 The conceptual map: related domains that must not be collapsed

A high-scoring answer defines the constructs before explaining their development.

TERM	MEANING
Sex characteristics	include chromosomal, gonadal, hormonal and anatomical features; sex assignment at birth is a social and clinical classification usually made from observed anatomy.
Gender identity	is the person's internal sense of gender.
Gender expression	is the outward communication of gender through appearance, behaviour, name, language and social role.
gender role	is the culturally organised set of expectations attached to gender categories.

These domains interact, but none is a simple read-out of another.

Sexual orientation concerns patterns of romantic or sexual attraction. **Sexual behaviour** describes what a person does, while **sexual identity** is the label or self-understanding a person adopts. Attraction, behaviour and identity may not align, especially during development or where disclosure is unsafe. **Psychosexual development** is broader than genital maturation: it includes bodily awareness, pleasure, attachment, fantasy, attraction, gender, privacy, boundaries, consent, intimacy, values and the meaning assigned to relationships.

- RULE

Ask **five** separate questions: **body, identity, expression, attraction and behaviour**. Do not infer one from another. Tanner morphology does not determine psychological maturity; gender expression does not establish orientation; and a single behaviour does not establish a stable identity.

5

SEPARATE QUESTIONS

19.2 Development is transactional: biology, relationships and culture continually modify one another

No single-cause theory adequately explains gender identity or psychosexual development. Biological organisation supplies a sexed body and contributes hormonal and neurodevelopmental influences. The child experiences that body through touch, caregiving, language and other people's responses. Cognitive development permits classification, perspective-taking and an increasingly stable autobiographical self. Social learning supplies models, reinforcement and sanctions. Family, peers, school, religion, media and wider culture specify which expressions are encouraged, tolerated or punished. The child is active within this system, attending selectively, interpreting feedback and shaping the responses received from others.

A transactional model therefore replaces the false choice between nature and nurture. Bodily characteristics affect how others classify and treat a child; those responses influence opportunities and self-description; the child's preferences evoke further responses; and puberty

later changes both the body and its social meaning. Temperament, cognitive capacity, neurodevelopmental differences, family flexibility, peer acceptance and exposure to stigma alter the pathway. The same observable expression can consequently have different meanings in different children and settings.

Development also occurs within a hierarchy of contexts. Caregiver language and discipline operate at the immediate level; school rules and peer cultures organise daily belonging; media and institutions circulate ideals of masculinity, femininity and attractiveness; and legal, religious and economic conditions influence safety and disclosure. Culture is not an ornamental final paragraph. It helps determine which possibilities can be imagined, named and enacted.

19.3 Cognitive-developmental, social-learning and gender-schema accounts

The major theories differ in what they treat as the engine of development. In the cognitive-developmental account, the child actively constructs gender categories. The conceptual sequence is gender labelling, recognition that gender is stable across time, and gender constancy despite superficial changes in clothing, activity or appearance. Once the child regards a category as self-relevant, behaviour is organised toward consistency with it. This account explains active self-socialisation but may understate biological influence, early learning and the flexibility of real gender experience.

Social-learning theory emphasises observation, imitation, reinforcement and punishment. Children see which behaviours are rewarded in caregivers, siblings, peers and media figures; direct praise or ridicule then increases or suppresses expression. Social-cognitive theory adds expectations, self-efficacy and reciprocal influence, so the child is not merely a passive recipient of conditioning. These models explain how local gender conventions are acquired and why expression can change across settings, but learning a convention does not prove that the convention determines the child's internal identity.

Gender-schema theory links cognition and culture. Repeated cultural distinctions form organised schemas that guide attention, memory, interpretation and behaviour. Information marked as relevant to the self is noticed and remembered more readily, while schema-inconsistent examples may be overlooked or distorted. Schemas are efficient but revisable. Exposure to broader roles, flexible caregivers and counter-stereotypic models can enlarge the child's map of what is possible without dictating identity.

19.4 Freud's psychosexual theory: historically examinable, not a contemporary milestone schedule

Freud remains important for the history of developmental psychiatry and for examination essays, particularly his emphasis on early experience, conflict and unconscious meaning. His account is not a modern developmental surveillance tool. It gives neither a valid way to infer gender identity nor a clinical rule for deciding whether childhood sexual behaviour is expected. Contemporary formulation places the theory beside cognitive, learning, attachment, biological and sociocultural accounts rather than allowing it to stand as the sole explanation.

19.5 Sexual development in childhood: curiosity, privacy, boundaries and relationship learning

Childhood sexual development is broader than overt sexual behaviour. It includes learning the names and functions of body parts, recognising bodily differences, experiencing comfort and pleasure, asking questions, developing modesty, understanding private and public conduct, rehearsing family and relationship roles in play, and learning that bodies belong to their owners. Self-touch, curiosity and exploratory play can occur in development, but their meaning depends on context rather than on the behaviour label alone.

Assess behaviour along several dimensions: spontaneity, frequency, flexibility, mutuality, developmental similarity of the children involved, power difference, secrecy, coercion, emotional tone, explicitness, intrusiveness, response to calm redirection and effect on functioning. Developmentally understandable exploration is generally flexible, non-coercive and not organised around fear or a marked power imbalance. Concern rises with force, threats, injury, persistent distress, compulsive repetition, a substantial asymmetry in power or developmental capacity, explicit imitation beyond the child's likely experience, or behaviour that remains intrusive despite appropriate limits. No single sign proves abuse, but a concerning pattern requires careful safeguarding assessment.

Adults teach psychosexual safety through accurate language, matter-of-fact answers, respect for privacy, permission to refuse unwanted touch, distinction between safe privacy and unsafe secrecy, and consistent boundaries that do not shame the body. Excessive alarm can attach fear and guilt to ordinary curiosity; dismissiveness can leave the child without language for disclosure. The developmental aim is neither sexualisation nor silence, but embodied autonomy, appropriate privacy and the capacity to seek help.

19.6 Puberty and Tanner staging: physical maturation changes the developmental task

Secondary sexual characteristics make gender experience more immediate. See the Puberty topic.

19.7 Adolescence: identity formation, neurodevelopment and context-sensitive risk-taking

Normal adolescent development is asynchronous across body, brain and social role. Pubertal maturation increases the salience of body image, attraction, peer evaluation and belonging, while autonomy, intimacy, values and future roles are being renegotiated. In Erikson's account, the central conflict is identity versus role confusion. Identity integrates continuity across time with commitments in education, work, values, relationships, gender and sexuality. Ordinary uncertainty and exploration are not equivalent to role confusion.

19.8 Gender diversity, distress and contemporary classification

Variation in gender identity or expression is not itself evidence of mental disorder. A child may prefer activities, clothing, roles or companions that do not fit local stereotypes without experiencing incongruence or distress. Conversely, a child may experience a persistent difference between assigned sex and experienced gender even when expression is constrained. The clinician should explore the child's own account, consistency across contexts, bodily experience, family response, peer safety and functional impact without interrogating the child toward a predetermined conclusion.

WHO's ICD-11 places **gender incongruence** under conditions related to sexual health rather than mental and behavioural disorders, reflecting that transgender and gender-diverse identities are not conditions of mental ill-health. WHO also states that gender-variant behaviour and preferences alone are not a basis for assigning the diagnosis. The examinable distinction is therefore between identity diversity, incongruence and the presence of distress or clinical need. Distress may arise from bodily incongruence, but it may also arise from bullying, rejection, violence, shame or forced concealment; formulation must identify the source rather than attributing every symptom to identity.

■ **TRAP** **Do not diagnose from gender nonconformity.** Toys, clothing, mannerisms, friendships, fantasy play or one sexual experience do not by themselves establish gender identity, sexual orientation, abuse or disorder. The same rule prevents both pathologising diversity and missing genuine distress.

A developmentally competent interview uses the child's stated language where possible, avoids assumptions about heterosexuality or binary identity, and separates the young person's wishes from caregiver fears. Support does not mean forcing an identity label or predicting an adult outcome. It means maintaining safety, permitting honest exploration, reducing shame, treating co-occurring symptoms on their own merits and preserving access to appropriate specialist assessment when needed.

19.9 Clinical assessment, safeguarding and the Indian context

The assessment unit is the child within a developmental and relational system. Interview caregivers and the child, including time with the child separately when developmentally appropriate. Explain confidentiality and its safety limits. Establish the child's language for body and gender, pubertal status, body comfort, play and interests, attraction and relationships where relevant, family beliefs, peer response, school functioning, online exposure, privacy practices and sources of support. Ask what changed, when it changed, in which settings, what preceded it, how persistent and flexible it is, what meaning the child gives it, and whether distress belongs to the child, the family or both.

For sexual behaviour, reconstruct the event without repeated suggestive questioning. Assess the developmental capacities and power relationship of everyone involved, mutuality, coercion, threats, secrecy, explicitness, injury, emotional response, access to sexual material and the child's subsequent behaviour. Safeguarding concern requires a calm, child-centred response, careful documentation in the child's own words, attention to immediate safety and use of the appropriate multidisciplinary and statutory pathway. A behavioural sign neither proves nor excludes abuse.

INDIA

Gender and psychosexual development may be negotiated within joint or extended families, strong educational expectations, religious and linguistic communities, gendered household roles and unequal permission for autonomy. Ask who actually provides care, who controls disclosure and what consequences the child anticipates. Interviewing only the accompanying adult can erase the child's account; interviewing only the child can miss dependence and safety constraints. Use clear language in the family's preferred language, separate cultural meaning from coercion, and do not use "Indian culture" either to pathologise family interdependence or to excuse humiliation, violence or forced conformity. No Indian population-specific age sequence for gender labelling or identity consolidation is supplied, so none should be inferred.

Formulation should distinguish predisposing factors, current developmental tasks, precipitating events, maintaining responses and protective relationships. Screen for mood and anxiety symptoms, self-harm, trauma, bullying, family violence, neurodevelopmental differences, substance use and online exploitation when indicated. A sudden or episodic change, regression, pervasive impairment, marked distress, coercion, dangerous exposure or loss of previously stable functioning weighs against a purely normative explanation. Treatment targets the actual source of difficulty rather than trying to make identity or expression conform to a stereotype.

19.10 Examination synthesis: move from definitions to development, then to clinical discrimination

A complete long answer has a clear spine.

- Define sex characteristics, gender identity, gender expression, gender role, sexual orientation and psychosexual development.
- Explain transactional development through biological, cognitive, learning, family and cultural influences.
- Contrast cognitive-developmental, social-learning and gender-schema theories.
- Present Freud as a historical stage account with source-stated mappings, not as a milestone schedule. Describe childhood curiosity, privacy and boundary learning, then connect puberty and Tanner morphology to adolescent identity formation, dual systems, maturational imbalance and risk-taking. Finish by distinguishing diversity and exploration from distress, impairment, coercion, trauma and disorder.

MNEMONIC

"Body, self, signal, attraction, behaviour and safety"

Retrieval spine: *body, self, signal, attraction, behaviour and safety*. Body recalls sex characteristics, puberty and Tanner staging. Self recalls gender identity and identity formation. Signal recalls expression and gender roles. Attraction recalls orientation. Behaviour recalls what is enacted rather than assumed. Safety recalls consent, power, distress, impairment and safeguarding.

Final clinical synthesis. Gender identity and psychosexual development are developmental processes, not conclusions drawn from a single behaviour. Good psychiatry keeps constructs separate, respects bodily and identity diversity, reads puberty through morphology and source-stated ranges, and places every concern within relationships, culture, consent, distress, functioning and safety.

Development in context and clinical use

20 . Developmental red flags and warning signs

A developmental red flag is not simply a child who is slower than a remembered textbook average. It is a finding that raises the probability of a developmental disorder sufficiently to require prompt evaluation, or a pattern so biologically atypical that ordinary variation is an unsafe explanation. The examiner expects the resident to distinguish *variation* from *delay*, and delay from *deviance*, *dissociation*, *plateau* and *regression*. The central clinical question is therefore not, "Has the child crossed one age line?" but, "What is the trajectory, which domains are affected, and is the pattern compatible with normal variation?"

How to use this page. First learn the few findings that demand further evaluation, especially loss of acquired skills and the language-social warning signs. Then learn to recognise abnormal trajectories across gross motor, fine motor, language, social-adaptive and cognitive domains. Finally, separate **surveillance**, which is longitudinal and continuous, from **screening**, which applies a standardised tool, and from **diagnostic assessment**, which determines the developmental profile and cause. The milestone and primitive-reflex plates are memory maps; this page owns the clinical act of spotting deviation and deciding what happens next.

Loss of previously acquired language or social skills at any age is a red flag and requires prompt developmental evaluation.




Fig 20.18 Red flags and surveillance, drawn so that only sourced referral signals appear. Coral pins are the language and social red flags from Developmental Psychopathology and the primitive-reflex red flags from Hutchison's Clinical Methods. Marigold markers are the 99th-centile boundaries of the WHO Motor Development Study windows, which WHO itself states are a signal to screen rather than a diagnostic threshold. No textbook referral threshold for absence of a gross-motor milestone was sourced, so none is drawn, and no surveillance schedule appears because RBSK and ICDS specifics were not sourced.




Fig 20.19 The developmental screening and assessment instruments compared by what is sourced. Only the Trivandrum chart carries validated psychometrics, from Nair and colleagues, and its accuracy is against the Denver Developmental Screening Test. The DASII contributes its two quotients and the delay marker below 85; the Denver test contributes only an age range; the Baroda test contributes only its year and its primacy. Dashed cells are properties that were searched for and are not sourced, left empty rather than filled from memory. No instrument form is reproduced.

20.1 Normal variation, delay and abnormal developmental trajectories

Development is a trajectory, not a checklist. A milestone age has meaning only when its construct is known.

MILESTONE CONSTRUCT	MEANING
A median	means that half of children have attained the skill by that age.
An expected range	describes a normal window.
An upper limit	is the outer bound regarded as normal.
A red-flag threshold	is an operational point at which further evaluation is warranted.

Converting an average or median into a referral threshold is a clinical error. Equally, being within an age window does not neutralise abnormal quality, asymmetry, regression or a concerning neurological examination.

Developmental delay is slower acquisition of skills in one or more domains while the usual sequence is broadly preserved. Global developmental delay describes substantial delay across multiple developmental domains in a young child, but the label does not state the cause.

Dissociation is unequal development between domains, such as relatively preserved motor ability with marked language-social delay. Deviance is an unusual sequence or quality of development rather than mere lateness. A plateau is failure to acquire new skills over time. Regression is loss of a skill that was convincingly acquired, and is more alarming than isolated late acquisition because it suggests an active or progressive process until shown otherwise.

- **RULE** Judge the child by trajectory, pattern and quality, not by one remembered average age. A red flag can arise from absence, loss, asymmetry, abnormal sequence, abnormal movement quality or a widening gap between domains.

MNEMONIC

"Trajectory, pattern and quality"

Judge the child by trajectory, pattern and quality, not by one remembered average age. A red flag can arise from absence, loss, asymmetry, abnormal sequence, abnormal movement quality or a widening gap between domains.

20.2 Developmental surveillance, screening and assessment are different clinical acts

Developmental surveillance is the repeated, flexible clinical process conducted across encounters. It integrates parental concerns, developmental history, observation of the child, risk factors, physical and neurological examination, and the trajectory recorded at previous visits. Ask what the child can do spontaneously in daily life, not merely whether the child once performed a prompted task. Clarify onset, consistency, context, quality and whether any skill has been lost. Observe communication, play, reciprocity, posture, movement, hand use and problem-solving before directing the child, because spontaneous behaviour often contains the most valid developmental information.

Developmental screening is the use of a brief standardised instrument in children who may or may not already have an identified concern. It estimates risk; it does not diagnose. A negative screen does not overrule convincing regression, abnormal examination or persistent caregiver concern. Developmental assessment is the more detailed, domain-based evaluation used to characterise strengths and impairments, quantify development where appropriate, identify associated sensory or neurological problems, formulate differentials and plan intervention. Surveillance generates suspicion, screening standardises risk detection, and assessment explains the profile.

20.3 Spotting deviation: read the pattern before the individual milestone

Use five questions whenever development appears atypical.

1. Is this *absence* of a skill, *loss* of a skill, or merely inconsistent performance?
2. Is one domain affected or are multiple domains involved?
3. Is the sequence recognisable but slow, or qualitatively unusual?
4. Are there associated neurological signs, abnormal head growth, dysmorphism, feeding difficulty, poor vision or hearing, seizures, altered tone, weakness or asymmetry?

5. What is the functional consequence in communication, play, self-care, mobility, learning and participation?

Patterns narrow the differential. Predominantly motor delay with abnormal tone or reflexes points toward a motor-system disorder; language delay requires separation of hearing impairment, developmental language disorder, broader cognitive delay and autism-related social-communication difficulty; social-communication deviation with restricted or repetitive behaviour raises concern for autism; and global delay requires a broad biological and environmental assessment. In every pattern, check pregnancy and birth history, neonatal illness, seizures, infections, nutrition, medication or toxin exposure, family history, consanguinity, psychosocial deprivation and opportunities for learning. These factors modify interpretation but do not justify watchful waiting when a hard red flag is present.

Asymmetry is a warning sign even when an age threshold has not been crossed. Persistent one-sided hand use in infancy, unequal movement, early fixed hand preference, scissoring, persistent fisting, abnormal posturing or an abnormal gait directs attention to focal or motor-pathway pathology. Likewise, the manner of a skill matters: sitting with marked truncal instability and walking with an abnormal pattern are not made normal by timely onset.

20.4 Language and social-communication red flags: the hard thresholds

The most examinable age-linked warning signs are absence of early communication behaviours and loss of acquired social-communication skills. Developmental Psychopathology identifies the findings in the strip below as warranting further developmental evaluation. These are referral signals, not median attainment ages.

12 months NO BABBLING OR GESTURING	16 months NO SINGLE WORDS	24 months NO SPONTANEOUS TWO- WORD PHRASES	Any age LOSS OF LANGUAGE OR SOCIAL SKILLS
---	-------------------------------------	---	--

Interpret these against the quality of communication. Joint attention and pointing are expected around 12 months in most typically developing children according to the New Oxford Textbook of Psychiatry, and symbolic play is expected at 18 months in Dulcan's Textbook of Child and Adolescent Psychiatry. Concern rises when limited eye contact, poor response to name, absent sharing of interest, lack of communicative gesture, poor reciprocity or repetitive behaviour forms a coherent pattern. Do not diagnose autism from a single missed behaviour, but do not dismiss a convergent social-communication pattern because vocabulary appears adequate. The disorder itself, with its criteria and management, is developed in the Intellectual disability and autism spectrum disorder notes; this page owns only the act of noticing the pattern and referring.

Speech clarity must also be interpreted by construct. Kaplan and Sadock describes speech as about **50 percent** intelligible at 24 months, about **75 percent** intelligible to strangers at 36 months, and nearly **100 percent** intelligible to strangers by 4 years. These are expected values, not automatic referral thresholds. A child who is difficult to understand requires assessment of hearing, oral-motor function, speech-sound production, receptive and expressive language, and the effect on participation.

20.5 Gross-motor warning signs: use WHO windows as screening signals

The WHO Motor Development Study provides construct-explicit windows of achievement bounded by the 1st and 99th centiles. It followed children in Ghana, India, Norway, Oman and the USA, so India was part of the source population. Sitting without support occurred from 3.8 to 9.2 months, standing alone from 6.9 to 16.9 months, and walking alone from 8.2 to 17.6 months. A child beyond the upper boundary signals the need for appropriate screening, which is the role WHO itself states for these windows. They are screening signals, not diagnostic cut-offs and not independently sourced referral thresholds.

Gross-motor concern is strengthened by an abnormal neurological examination, loss of motor skills, persistent asymmetry, abnormal tone, weakness, poor antigravity movement, abnormal movement quality, feeding or respiratory difficulty, or delay in other domains. Conversely, an isolated variation in sequence can be benign. The gross-motor plate should be read as a set of windows rather than as a ladder of compulsory achievements.

■ **TRAP** **Hands-and-knees crawling is not universal.** In the WHO study, 4.3 percent of children did not exhibit it at all. Absence of crawling alone is therefore not a developmental red flag; judge the overall trajectory, movement quality and neurological findings.

20.6 Primitive reflexes: persistence, asymmetry and re-emergence matter

Primitive reflexes are expected in the newborn period and are progressively integrated as cortical control matures. Their interpretation is not a memory contest about disappearance ages alone. A reflex that is unusually persistent, strongly asymmetric, absent when it should be present, or re-emerges after integration may indicate neurological dysfunction. The primitive-reflex plate provides the expected sequence; the clinical examination must compare the reflex response with tone, posture, spontaneous movement, symmetry and the child's functional motor repertoire.

Hutchison's Clinical Methods treats persistence of primitive reflexes beyond the fourth month of life as a general warning sign, persistence of the Babinski response beyond 2 years as a red flag, and absence of the sucking reflex beyond the 36th week of gestation as a red flag. These sourced warning signs should not be replaced by conflicting remembered averages. A reflex abnormality warrants a neurological and developmental assessment rather than an isolated label.

20.7 Indian screening instruments: TDSC and DASII

The Trivandrum Development Screening Chart (TDSC), developed at the Child Development Centre, Thiruvananthapuram Medical College, is an Indian screening instrument for children from birth to 6 years. The validated chart contains 51 items, and **one item delay** constitutes a positive screen termed "TDSC delay". In its validation against the Denver Developmental Screening Test, sensitivity was 84.62 percent, specificity 90.8 percent, negative predictive value 99.23 percent, test-retest reliability 0.77, and inter-rater reliability 0.97. These properties make it useful for screening, but a positive result is not a diagnosis and a negative result cannot cancel a hard clinical red flag; the underlying theory of sensitivity, specificity, predictive value and reliability belongs to the Psychometry, Assessment and Descriptive Psychopathology notes.

The Developmental Assessment Scales for Indian Infants (DASII) yields separate **Motor DQ** and **Mental DQ**. Indian follow-up studies use a DQ below 85 as a marker of delay. DASII is an assessment measure rather than a substitute for surveillance, and the two quotients help display dissociation between motor and mental development. The Baroda Development Screening Test, introduced in 1991, was India's first developmental screening tool according to Shekhawat and colleagues. Keep the categories clear in an answer: TDSC and the Baroda instrument are screening tools; DASII provides domain-based developmental quotients.

20.8 When to refer and what the first evaluation must contain

Refer promptly when there is regression, a hard red flag, abnormal neurological findings, marked asymmetry, a deviant developmental sequence, multi-domain delay, or persistent concern despite a reassuring screen. Do not use "wait and watch" as a substitute for surveillance. If observation is appropriate for a borderline concern, it requires a documented baseline, an explicit review plan and instructions to return earlier if skills are lost or new warning signs appear. Early referral is not overdiagnosis: it opens the route to hearing and vision assessment, developmental and neurological evaluation, aetiological investigation where indicated, and early intervention while diagnostic clarification continues.

The initial doctor-facing work-up begins by verifying the concern with caregiver examples and direct observation, mapping skills across all domains, and establishing whether the course is delayed, dissociated, deviant, plateauing or regressive. Examine growth, head size and trajectory, dysmorphic features, skin markers, vision, hearing, oral-motor function, tone, power, reflexes, coordination, gait and asymmetry. Review antenatal, perinatal, neonatal, medical, seizure, nutritional, family and psychosocial history. Select investigations from the phenotype rather than ordering an indiscriminate panel, and refer to the appropriate developmental paediatric, paediatric neurological, audiological, ophthalmological, speech-language, occupational or mental-health service.

The viva answer. Define the warning sign, state whether the quoted age is an expected value, a range, an upper limit or a red-flag threshold, describe the whole developmental pattern, and finish with action. The safest closing line is: *I would not diagnose from a missed milestone alone, but I would not delay evaluation when there is regression, a sourced red flag, abnormal quality or a convergent multi-domain concern.*

21 . Effects of deprivation, institutionalization and early adversity on development

Early development depends not only on freedom from injury, but on the reliable presence of expected experience: responsive caregiving, language, touch, play, nutrition, safety and opportunities to explore. **Deprivation** is the absence or marked reduction of these expected inputs. **Threat** is the presence of experiences that signal actual or possible harm, such as violence or abuse. **Early adversity** is the broader term, covering deprivation, threat, loss, instability, discrimination, displacement, severe family stress and other conditions that exceed the child's available supports. These exposures often co-occur, but they are not interchangeable. The

distinction matters because a child lacking cognitive and social input may follow a different developmental pathway from a child repeatedly learning that the environment is dangerous.



Fig 20.20 Early adversity is not one exposure. Deprivation and threat can follow different developmental pathways, while the nature, timing, duration and severity of exposure and the availability of responsive caregiving shape both persistence and recovery.

How to use this page. Build the answer around **five** propositions.

1. Effects depend on the nature, timing, duration and severity of the adversity, not merely its presence.
2. **Institutionalization** is not a single exposure: it may combine rotating caregivers, low responsiveness, sensory monotony, crowding, malnutrition, illness and repeated loss.
3. The classical studies by Spitz and Harlow established the developmental importance of relationship and contact, but have major methodological and ethical limits.
4. The Romanian natural experiments show both persistent deprivation-related difficulties and substantial recovery after family placement.

5. **Resilience** is an outcome produced by interacting systems, not a fixed trait that makes a child invulnerable.

MNEMONIC

"NTDS"

Effects depend on the **N**ature, **T**iming, **D**uration and **S**everity of the adversity, not merely its presence.

five

PROPOSITIONS

21.1 Concepts and causal framework: deprivation is not the same as separation, poverty or threat

Define the exposure before attributing an effect. **Neglect** is a failure to meet a child's physical, medical, educational, emotional or supervisory needs. **Psychosocial deprivation** refers particularly to insufficient responsive interaction, stable attachment opportunities, language and cognitive stimulation. **Maternal deprivation** is the older historical label, but it is too narrow when it implies that only a biological mother can provide the necessary relationship. The developmentally relevant issue is the availability of a small number of stable, sensitive and committed caregivers. **Separation** is an event; its effect depends on what relationship is lost, what care replaces it, how predictable the transition is and whether reunion and continuity are possible. **Privation**, used in older literature, means that a selective attachment relationship was never adequately established, whereas deprivation may refer to loss of, or insufficient access to, an expected input.

Poverty increases exposure to nutritional compromise, environmental hazards, caregiver stress, disrupted schooling and reduced access to care, but poverty itself is not proof of neglect or deficient attachment. Likewise, institutional residence does not make every child's experience identical. Quality varies with caregiver continuity, responsiveness, group size, opportunities for play and language, health care and contact with family. The causal formulation therefore asks: *what input was absent, what threat was present, when and for how long, through which mediators, and what protective relationships remained?*

Sheridan and McLaughlin's dimensional model separates deprivation from threat.

DIMENSION	DEFINITION	DEVELOPMENTAL RELEVANCE
deprivation	the absence of expected social and cognitive inputs	Deprivation is especially relevant to language, executive function and higher-order cognitive development because the relevant circuits receive less patterned experience.
threat	experience involving harm or danger	Threat more directly calibrates fear learning, attention to danger and emotion-processing systems.

The model is not a claim of **one** exposure to one brain region. Real children commonly experience both dimensions, and development reflects their interaction with genes, prenatal conditions, nutrition, illness, relationships and later environments.

■ **RULE Exposure is not outcome.** Early adversity changes probabilities across several developmental domains; it does not establish inevitable damage, a particular diagnosis or poor parenting by inference.

21.2 Mechanisms: how early adversity becomes biologically and psychologically embedded
Adversity acts through repeated transactions between the child and the environment. A responsive caregiver ordinarily provides co-regulation: detecting the infant's state, reducing distress, organising sleep and feeding, sharing attention, naming experience and making exploration safe. When care is inconsistent or impersonal, arousal may remain poorly regulated and the child receives fewer contingent exchanges from which language, social prediction and self-regulation are learned. A threatening environment adds repeated detection of danger, narrowing attention toward immediate survival cues. The adaptations may be useful in the adverse setting while carrying costs in a safer one.

The stress-response system is one mediator. Repeated or uncontrollable stress can alter hypothalamic-pituitary-adrenal regulation, autonomic balance, immune and inflammatory signalling, sleep and metabolic function. The expected pattern is not simply "more cortisol": chronic exposure may produce elevation, flattening or blunted reactivity depending on timing, measurement and adaptation. At the neural level, studies report group differences in cortical and white-matter development, frontolimbic connectivity, reward learning and executive-control systems after severe deprivation. These are probabilistic associations, not a diagnostic brain signature. Prenatal risk, malnutrition, infection, genetic liability, placement selection and later experience can confound observed differences.

Learning mechanisms are equally important. Sparse language and play reduce opportunities to acquire vocabulary, causal models and executive strategies. Unpredictable care makes sustained attention and delayed reward less useful than rapid switching. Repeated caregiver loss can make adults unreliable sources of safety. Harsh or frightening care may produce hypervigilance, avoidance, dissociation, aggression or compulsive compliance. Epigenetic findings offer plausible

pathways by which experience can influence gene expression, but they do not justify claims that adversity permanently "rewrites" a child's genes. Biological embedding remains open to modification because the developing system continues to transact with later care.

21.3 Developmental phenotype: broad effects with marked individual variation

Severe deprivation can affect growth, cognition, language, motor development, attachment, emotion and behaviour at the same time. Physical findings include poor weight or linear growth, feeding and sleep disturbance, recurrent illness and stereotyped self-stimulation. Developmental effects may include delayed motor coordination, reduced language exposure and vocabulary, lower general cognitive performance, weak working memory, impaired sustained attention and executive dysfunction. School-age manifestations may include learning difficulty, inattention, impulsivity and problems with planning rather than a uniform global deficit.

Socioemotional development can follow more than one pattern. Some children are withdrawn, wary, emotionally constricted and do not reliably seek or accept comfort. Others show indiscriminate approach, reduced reticence with unfamiliar adults and poor checking back with a caregiver. In current classification these patterns are distinguished as reactive attachment disorder and disinhibited social engagement disorder, which are developed in the Child behavioural, emotional and other disorders notes. Both require a history of grossly insufficient care, but neither is diagnosed from that history alone. Assessment requires the child's actual pattern of behaviour, direct observation with current caregivers and exclusion of better explanations. Indiscriminate friendliness is not equivalent to secure attachment, and a child may develop a selective attachment to a new caregiver while disinhibited social behaviour persists.

Emotional and behavioural outcomes include anxiety, depressive symptoms, irritability, aggression, emotional dysregulation, dissociation, self-harm risk and later interpersonal difficulty. Early institutional deprivation has also been associated with a characteristic cluster involving inattention or overactivity, autistic-like social-communication features, disinhibited social behaviour and cognitive impairment. These are group-level patterns, not proof that every previously institutionalized child has a deprivation-specific syndrome. Development may show multifinality, in which the same adversity leads to different outcomes, and equifinality, in which a similar clinical presentation arises from different developmental routes.

21.4 Spitz: anaclitic depression and hospitalism

Spitz's observations made emotionally responsive care visible as a developmental necessity.

René Spitz compared infants reared in different institutional settings and described deterioration after loss of a familiar caregiver despite provision of basic physical care. He used anaclitic depression for the syndrome following prolonged separation from an attachment figure: protest and weeping gave way to withdrawal, reduced responsiveness, disturbed feeding and sleep, developmental slowing and physical decline. Hospitalism described the more pervasive developmental deterioration seen with prolonged impersonal institutional care, including passivity, failure to thrive, susceptibility to illness and high mortality in the historical settings he observed.

The enduring inference is not that separation from the biological mother mechanically causes depression. It is that food, cleanliness and medical custody are insufficient without stable, contingent and affectionate human interaction. Spitz also helped shift hospital and residential practice toward caregiver access and relational continuity. His terms are historically examinable, but anaclitic depression is not simply the infant form of a current major depressive episode, and hospitalism is not a modern diagnostic category.

The limitations are substantial. The studies were observational, the institutional groups differed in more than caregiver contact, physical illness and nutrition were incompletely controlled, assessment was not blinded, and the interpretation was shaped by contemporary psychoanalytic theory. These weaknesses prevent precise causal estimates. The observations remain influential because later attachment research, adoption studies and intervention trials converge on the importance of stable, responsive care.

21.5 Harlow: contact comfort, fear regulation and the limits of animal evidence

Harlow showed that feeding alone does not explain attachment behaviour. Infant rhesus macaques were reared with inanimate surrogate mothers, one made of bare wire and another covered with soft cloth. Even when the wire surrogate supplied milk, the infants spent much more time clinging to the cloth surrogate. Under fear, they sought the cloth surrogate, and its presence supported exploration of an unfamiliar setting. Harlow called the relevant property contact comfort. The experiment directly challenged drive-reduction accounts in which attachment was merely learned because the caregiver relieved hunger.

Later social-isolation work showed disturbed social interaction, fear, repetitive behaviour and impaired mating and parenting in monkeys deprived of normal peer and maternal experience. Partial improvement after later social contact demonstrated both the cost of early deprivation and some capacity for recovery. The broader lesson is that attachment is supported by comfort, safety, reciprocal interaction and species-typical social experience, not by calories alone.

The evidence cannot be translated literally from macaques to children. Human caregiving includes language, culture, symbolic meaning, multiple attachment figures and institutions that vary widely. Rearing conditions were artificial, outcomes were partly species-specific, and the procedures caused severe suffering that would not meet contemporary ethical standards. Harlow's work therefore supports a mechanism, the motivational importance of comfort and relationship, but does not define a human sensitive-period cutoff or prove a deterministic human outcome.

■ **TRAP** Spitz did not prove that only mothers matter, and Harlow did not prove that feeding does not matter. Both challenged custodial models of care by showing that stable, responsive relationship and comfort are independent developmental needs.

21.6 Romanian natural experiments: ERA and the Bucharest intervention

The Romanian studies provide the clearest human evidence that severe institutional deprivation has causal developmental effects and that family care can produce substantial recovery. After the fall of the Romanian regime, many children who had experienced profoundly

understaffed, socially and cognitively sparse institutions entered adoptive families. The English and Romanian Adoptees study, or ERA study, followed Romanian adoptees placed in the United Kingdom and compared their development with adoptees who had not experienced the same severe deprivation. Because nutrition, prenatal risk and deprivation still varied, it was not a randomized experiment. Its strength was the unusual, time-limited exposure followed by generally well-resourced adoptive care and prolonged follow-up.

Many children showed striking catch-up in physical growth and cognitive performance after adoption, demonstrating that early effects were not fixed. A subgroup with more prolonged severe deprivation showed persistent or emerging difficulties in cognition, inattention or overactivity, disinhibited social engagement and autistic-like social communication. The ERA investigators described this as a deprivation-specific pattern. Some difficulties persisted into adult life even after years in supportive families, while emotional symptoms could become more apparent later. **Two** conclusions must be held together: recovery can be large, and residual vulnerability can remain.

The Bucharest Early Intervention Project added stronger causal evidence by randomly assigning institutionalized children to high-quality foster care or care as usual, with a never-institutionalized comparison group. Assignment to foster care produced better cognitive and physical outcomes and fewer attachment or social-relatedness difficulties and other psychopathology across follow-up. Studies also found more favourable trajectories in brain electrical activity and aspects of white-matter development. Placement timing and the subsequent quality and stability of care influenced some outcomes, but not every domain showed the same timing effect. It is therefore inaccurate to teach one universal age after which recovery is impossible.

The projects are complementary rather than interchangeable. ERA is an adoption-based longitudinal natural experiment with strong evidence of long-term developmental patterning. The Bucharest project is a randomized foster-care intervention that strengthens causal inference about the benefit of family placement. Ethical debate remains because the trial occurred in a setting where institutional care was still the prevailing care pathway, and researchers did not remove children assigned to care as usual. Its central clinical result is nevertheless clear: high-quality family care is an active developmental treatment, not merely a change of address.

21.7 Sensitive periods, cumulative risk and developmental timing

A sensitive period is a time of heightened plasticity, not an expiry date for recovery. During periods when attachment, language, sensory systems and stress regulation are being rapidly organised, expected input can have unusually strong effects. Earlier onset, greater severity, longer duration and involvement of multiple developmental systems generally increase risk. Yet the timing gradient differs by outcome: language, attachment behaviour, executive function, growth and emotional symptoms do not share one timetable. A single global cutoff converts a graded and domain-specific finding into a false rule.

The cumulative-risk model predicts increasing difficulty as adversities accumulate. It is useful for public-health stratification but an unweighted count hides whether the exposures were deprivation, threat or instability, their severity and duration, and the protective systems surrounding the child. Adverse childhood experience scores similarly describe population-level dose-response associations; they are not diagnostic tests, causal explanations or individual prognoses. Two children with the same count may have very different exposures, meanings and outcomes.

Development also involves experience-expectant plasticity, in which the brain anticipates common inputs such as patterned language and responsive interaction, and experience-dependent plasticity, in which individual learning modifies circuits throughout life. Early deprivation can therefore constrain the architecture on which later learning builds, while later relationships, education and treatment can still reorganise function. Recovery may be domain-specific: physical catch-up can coexist with executive or relational difficulty, and secure attachment to a new caregiver can coexist with persistent social disinhibition.

Must remember: early adversity is probabilistic, timing is domain-specific, and stable responsive caregiving is both the principal protective factor and the core intervention.

21.8 Clinical assessment, prevention and management

The clinical task is to reconstruct the developmental ecology, identify current needs and change the caregiving environment. Establish immediate safety first. Take a chronological account of prenatal and birth risk, nutrition, illness, developmental opportunities, separations, caregiver changes, institutional conditions, abuse or threat, schooling and current placement. Record the source and certainty of each claim because records may be incomplete and narratives may conflict. Assess growth, hearing, vision, sleep, feeding, neurological status, language, cognition, adaptive function, attachment behaviour, attention, mood, trauma symptoms, self-harm and social functioning. Observe the child with the present caregiver and obtain collateral information from school and previous services when appropriate.

Do not repeatedly elicit graphic trauma details when they are not required for care. Ask what happened, what continues, what triggers current difficulty and what helps. Separate a child's adaptations from moral judgement: food hoarding, watchfulness, indiscriminate friendliness, lying, emotional shutdown or controlling behaviour may have been functional in an unpredictable setting. Also avoid diagnostic overshadowing. Developmental disability, autism, attention-deficit/hyperactivity disorder, post-traumatic symptoms, depression, anxiety, malnutrition and sensory impairment may coexist with, resemble or occur independently of deprivation-related problems.

Management begins with a safe, stable, responsive family environment and the smallest feasible number of consistent caregivers. Prevent further placement disruption, support the caregiver-child relationship, coach caregivers in contingent responding and co-regulation, and establish predictable routines without punitive rigidity. Provide nutritional, medical, sensory,

developmental, speech-language and educational interventions according to need. Use trauma-focused treatment for clinically significant trauma symptoms, not simply because adversity occurred. Parent-child and attachment-focused work must be evidence-based and non-coercive; forced holding, induced regression and "rebirthing" practices are unsafe.

INDIA

In Indian practice, distinguish poverty and family stress from wilful neglect, and assess the wider caregiving network rather than assuming a nuclear-family model. Kinship care may preserve language, culture and continuity, but safety and caregiver capacity still require direct assessment. When residential care cannot be avoided, the developmental targets are caregiver continuity, responsive interaction, play and language, schooling, nutrition, health care, sibling and safe family contact, and an active plan toward stable family-based care. Coordination among paediatrics, psychiatry, clinical psychology, social work, speech and language services, education and child-protection systems is usually more important than any single test.

For an examination answer, define deprivation and distinguish it from threat; describe the multi-domain phenotype and mechanisms; present Spitz and Harlow with their limitations; compare the ERA natural experiment with the Bucharest randomized foster-care intervention; explain sensitive periods without a rigid cutoff; and finish with resilience, protective systems and family-based intervention. The mature conclusion is neither pessimistic nor sentimental: early severe deprivation can leave enduring developmental effects, but the developing child remains plastic, and timely, stable, high-quality relationships materially alter the trajectory.

22 · Resilience and protective factors in child development

Resilience is positive adaptation despite exposure to significant adversity. It is not invulnerability, a fixed temperament, or proof that adversity was harmless. It is a **dynamic developmental process** produced by transactions between the child and changing caregiving, family, school, peer, cultural and social systems. The same child may function well in one domain and struggle in another, and may show competence at one point but difficulty after a later developmental demand. The useful clinical question is therefore not, "Is this child resilient?" but, "What risks, resources and adaptive processes are operating now, in which domain, and what can be changed?"

How to use this page. Build the answer in **three** moves. First, establish both sides of the definition: meaningful adversity and demonstrable adaptation. Second, distinguish a factor that generally promotes development from one that specifically buffers the effect of risk. Third, move across ecological levels, from child capacities to at least one stable caring adult, family organisation, school belonging and community resources. The classic deprivation literature supplies causal clues, but its most defensible lesson is neither determinism nor romantic recovery: early relationships and stimulation matter greatly, developmental plasticity remains real, and earlier restoration of family care generally permits broader recovery.

3 RESILIENCE MODELS	5 ECOLOGICAL LEVELS	4 INTERVENTION TASKS	3 PREVENTION LEVELS
------------------------	------------------------	-------------------------	------------------------

Resilience is not a trait inside the child: it is positive adaptation arising from dynamic child-context processes despite significant adversity.

22.1 Core concepts: risk, promotive factors, protective factors and vulnerability

The work of Norman Garmezy, Michael Rutter and Emmy Werner redirected research from cataloguing damage toward studying variation in outcome and the processes that alter risk pathways. Ann Masten's phrase *ordinary magic* captures the mature position: resilience usually arises from ordinary adaptive systems, not rare heroic qualities. It also differs from post-traumatic growth, which denotes perceived or demonstrable positive change after struggle; a child can be resilient without reporting growth, and claimed growth does not exclude distress.

A **risk factor** raises the probability of an adverse outcome; it does not make that outcome inevitable. Risks may be biological, psychological or social, and their meaning depends on timing, duration, severity and developmental context. **Cumulative risk** is often more informative than a single exposure because poverty, caregiver illness, violence, maltreatment, discrimination, disrupted schooling and unsafe neighbourhoods cluster and transact. **Equifinality** means that different developmental pathways can lead to a similar outcome, whereas **multifinality** means that the same early adversity can lead to different outcomes.

A **promotive factor** predicts better functioning across levels of risk. A **protective factor**, in the stricter research sense, modifies the effect of adversity: its benefit is especially evident when risk is present. A **vulnerability factor** amplifies the effect of risk. A resource such as good schooling may be promotive in most children and protective under family adversity, depending on the hypothesis and data. This distinction matters in an examination answer because merely listing desirable characteristics does not explain resilience.

- **RULE Demonstrate adversity and adaptation separately.** Resilience can be inferred only when significant risk has occurred and the child shows adequate functioning in a specified domain, judged against developmentally and culturally appropriate expectations.

22.2 Models of resilience: how protection changes a developmental pathway

MODEL	HOW PROTECTION CHANGES THE PATHWAY
compensatory model	A positive influence has a beneficial effect independent of risk, partly offsetting its burden.
protective or buffering model	An interaction: the association between adversity and poor outcome is weaker when the protective process is present.
challenge, steeling or inoculation model	Manageable stress, successfully negotiated with adequate support, may strengthen later coping. It must not be used to justify preventable adversity; overwhelming or uncontrollable stress is not therapeutic.

Differential susceptibility and biological-sensitivity models add a further correction. A highly reactive child may fare worse in an adverse environment yet gain more from a supportive one. The same characteristic is therefore not always a vulnerability. Contemporary developmental psychopathology treats resilience as a changing trajectory generated by person-context transactions, developmental cascades and feedback loops. Improved caregiver responsiveness may strengthen regulation, which improves school participation, which then enlarges peer support and competence. Conversely, one failure can spread across domains if no system interrupts it.

22.3 Early adversity: threat, deprivation and biological embedding

Early adversity is heterogeneous. Experiences dominated by threat, such as violence or abuse, make harm salient and may bias fear learning, vigilance and emotion processing. Experiences dominated by deprivation, such as severe neglect or understimulation, remove expected cognitive, linguistic and relational input and may particularly affect learning and executive development. Most real children experience mixtures of both, so the distinction is a formulation aid rather than a diagnostic partition.

Repeated activation of stress-response systems without adequate co-regulation can produce toxic stress and allostatic load. Candidate pathways include altered hypothalamic-pituitary-adrenal regulation, autonomic reactivity, sleep disturbance, inflammatory signalling and experience-dependent changes in neural networks supporting attention, reward, memory and affect regulation. These are probabilistic mechanisms, not permanent lesions. Genes influence sensitivity to environments; environments influence gene expression and neural development; later caregiving and learning can alter the trajectory. This is the biological basis for holding risk and plasticity together.

22.4 Romanian orphan studies: the abstracted lesson

Developmental plasticity is real. Timing and the quality and stability of placement matter. Catch-up is domain-specific: catch-up in height or IQ does not prove recovery in attachment, attention, social functioning or later mental health.

-
- **TRAP** Do not turn deprivation research into either determinism or "complete recovery". Group-level persistence does not fix an individual child's destiny, and catch-up in height or IQ does not prove recovery in attachment, attention, social functioning or later mental health.
-

22.5 Sensitive periods, developmental timing and recovery

A **sensitive period** is a phase during which experience has an unusually strong influence on a developing system; learning may remain possible later, but often requires more input or follows a different route. A **critical period** is a narrower claim in which an experience is required within a restricted window for typical development. Human psychosocial development usually warrants the more cautious term *sensitive period*. The institutional studies support timing effects, but they do not justify one universal cut-off for every function.

Different systems have different windows and recovery curves. Attachment organisation, language, executive function, growth, peer reciprocity and stress regulation need not recover together. Outcome depends on the nature of deprivation, its duration and severity, the child's prior condition, biological susceptibility, the timing and stability of placement, and the quality of later caregiving and education. Developmental cascades explain why early intervention has broad value: restoring caregiver regulation can improve attention and exploration, which expands learning, competence and relationships. The converse is also true, so later transitions can reveal difficulties that seemed absent earlier.

22.6 Protective processes across the child's ecology

- **Child level.** Cognitive ability, language, flexible problem-solving, self-regulation, sociability, realistic self-efficacy and the capacity to seek help can support adaptation. These are partly shaped by experience and should not be moralized as willpower.
- **Caregiver level.** The most consistently useful process is at least one stable, committed and emotionally available adult who notices the child, provides co-regulation, sets predictable limits and repairs relational ruptures.
- **Family level.** Warmth, routines, low conflict, safe housing, adequate material resources, shared meaning and support for caregivers reduce chaos and preserve developmental opportunity.
- **School and peer level.** A safe school, teacher attunement, academic support, opportunities for mastery, inclusive peer relationships and protection from bullying create competence and belonging.
- **Community and cultural level.** Accessible health and social services, safe neighbourhoods, social cohesion, mentors, recreation, faith or cultural networks, and policies that reduce poverty and discrimination operate upstream. Culture shapes which outcomes count as competent adaptation and which supports are acceptable.

A protective process is strongest when it is developmentally matched, available in ordinary settings and sustained across transitions.

INDIA

Formulate the ecology, not merely the child. Ask about caregiver mental illness and substance use, domestic violence, migration, caste or minority stress, gendered restriction, disability, school exclusion, examination pressure, digital and peer environments, and material insecurity. Then map usable protection: an extended-family adult, teacher, peer, Anganwadi or school support, paediatric and mental-health services, child-protection mechanisms and local community networks.

Extended family can be protective, neutral or harmful; do not assume that family presence equals emotional safety. When abuse, exploitation or neglect is ongoing, safeguarding and reduction of exposure come before teaching the child coping skills.

22.7 Assessment: a strengths-based formulation without minimizing harm

Assessment begins with a timeline of exposures, developmental transitions and changes in functioning. Specify the adversity by type, severity, duration, proximity and current status; separate threat from deprivation where useful; and identify mediators such as sleep, caregiver responsiveness, school attendance and peer exclusion. Examine adaptation in multiple domains: symptoms, attachment and relationships, learning, behaviour, physical health, self-care and age-appropriate participation. Obtain accounts from the child, caregivers and school because competence and distress are context-dependent.

Then construct a resilience map: risks and vulnerabilities; existing resources; mechanisms through which they help; inaccessible or unstable supports; and leverage points that can change the trajectory. A resilience questionnaire may organize information but cannot certify a resilient personality. Cross-sectional good functioning may conceal effort, dissociation, parentification or a later cost, while symptoms in one domain can coexist with genuine competence in another. Trauma-informed assessment therefore combines safety, curiosity and developmental formulation without forcing disclosure or demanding a narrative of growth.

22.8 Intervention: reduce adversity, restore relationships and build adaptive systems

Intervention is not "resilience training" alone. The **first** task is to stop or reduce the active adversity: ensure safety, treat caregiver illness, address violence and substance use, secure nutrition, housing and schooling, and prevent repeated placement disruption. The **second** is to strengthen caregiving through responsive parent work, co-regulation, predictable routines and support for caregiver stress. The **third** is to build the child's skills in a developmentally fitted way, including emotion identification, problem-solving, executive supports, social competence, mastery and help-seeking. The **fourth** is to enlarge the ecology through school accommodations, mentoring, peer connection and coordinated health, education and social care.

MNEMONIC**"Reduce - Restore - Build - Enlarge"**

Reduce the active adversity; restore relationships through responsive parent work, co-regulation, predictable routines and support for caregiver stress; build the child's skills; enlarge the ecology through school accommodations, mentoring, peer connection and coordinated health, education and social care.

Effective prevention combines universal supports for all children, selective interventions for groups exposed to elevated risk and indicated treatment for children already showing difficulty. Review outcomes over time and across domains, especially at transitions. The ethical endpoint is not to make children tolerate unsafe systems. It is to reduce preventable exposure, increase dependable protection and restore the ordinary relationships and opportunities through which development proceeds.

22.9 Exam synthesis

A complete long answer opens with resilience as a dynamic process, distinguishes promotive from protective effects, and gives compensatory, buffering and challenge models. It then explains cumulative risk, equifinality, multifinality and developmental cascades; uses Spitz, Harlow and the Romanian orphan natural experiments with their limitations; and links the ERA findings to sensitive periods without claiming a universal cut-off. Finish with child, caregiver, family, school and community protective processes, followed by assessment and intervention. The single sentence that rescues the answer from a list is this: **resilience emerges when ordinary adaptive systems, especially stable caregiving, regulation, learning, belonging and social protection, remain available or are restored under adversity.**

Apply and consolidate

23 . Apply: worked cases

These cases close the developmental arc by making the resident perform the decisions that lists of theories and milestones are meant to support. The sequence is deliberate: date a milestone without turning a typical age into a deadline, recognise a genuine referral signal, identify an attachment strategy from reunion behaviour, formulate temperament as a person-environment fit, and apply a developmental theory without pretending that it is a screening instrument.

MNEMONIC**"Phenomenon - construct - comparison - clinical action"**

Identify the phenomenon actually being tested, name the relevant construct, compare it only with an age or pattern that measures the same construct, and state what follows clinically.

How to use this page. In each case, take the steps in this order.

WHAT TO ESTABLISH	TASK
Phenomenon	Identify the phenomenon actually being tested.
Construct	Name the relevant construct.
Comparison	Compare it only with an age or pattern that measures the same construct.
Clinical implication	State what follows clinically.

The answer comes last. A correct label without the intervening reasoning is fragile in a viva and unsafe in practice.

23.1 Milestone dating: late against a typical age, still within the WHO window

Presentation. A child at 15 months of age pulls to stand, cruises along furniture, stands briefly without support and communicates socially, but has not yet walked alone. There has been steady acquisition of skills, no loss, no asymmetry and no abnormality of tone or movement quality. The caregiver has read that children usually walk independently by **12 months** and asks whether this proves **developmental delay**.

Reasoning. The first task is to name what the two ages mean. Kaplan and Sadock describes independent walking as expected at 12 months and gives an expected range of 9 to 15 months. An **expected age** describes typical attainment; it is not a **referral threshold**. The WHO Motor Development Study gives a different and more construct-explicit measure: its 1st-to-99th-centile window for walking alone is **8.2 to 17.6 months**. WHO states that its upper boundary is a signal for appropriate screening when a child appears late, not a diagnostic cut-off.

The child is later than the typical 12-month anchor but remains within the WHO achievement window and at the end of Kaplan and Sadock's expected range. The preserved progression toward independent gait and the absence of regression, asymmetry or abnormal motor quality further reduce the force of a single late-looking date. This does not mean that development is ignored. Surveillance continues, and examination still covers all domains, hearing, vision and neurology. It means only that the typical age cannot be promoted into a referral rule.

Answer. This is a late-normal gross-motor attainment on the information given, not developmental delay established by the 12-month expected age. Continue developmental surveillance and reassess the trajectory. Crossing the WHO **17.6-month** boundary would be a screening signal, while regression or abnormal neurological findings would change the action immediately.

23.2 Deviation call: a sourced communication red flag

Presentation. A child at 24 months of age uses several meaningful single words and obtains desired objects by pointing, but never produces spontaneous two-word phrases. Motor development has progressed, and the caregiver reports no loss of previously acquired skills.

Reasoning. The finding is not judged from vocabulary size alone. Berk gives 18 to 24 months as an expected range for combining two words, but the clinically decisive statement is separate:

Developmental Psychopathology identifies absence of spontaneous two-word phrases at **24 months** as a **red flag** warranting further developmental evaluation. Here the age is explicitly a referral signal, not merely an expected attainment or median.

The threshold does not diagnose a language disorder or autism. The next step is to define the profile.

- Confirm whether phrases are truly spontaneous rather than imitated.
- Assess receptive language, gesture, joint attention, reciprocity and symbolic play.
- Establish language exposure across all languages used by the child.
- Examine hearing and oral-motor function.

Map gross motor, fine motor, cognitive or adaptive, and social-emotional development rather than inferring global delay from one communication finding. The absence of regression is reassuring about the course but does not cancel the hard red flag.

Answer. This is a developmental red flag requiring prompt hearing and developmental evaluation. It is a referral decision, not a diagnosis, and a reassuring motor history or a negative brief screen would not overrule it.

23.3 Attachment-pattern identification: contact sought but not used effectively

Presentation. During a Strange Situation observation, a young child explores little, becomes intensely distressed when the caregiver leaves, and hurries toward the caregiver on reunion. Once picked up, the child arches away, pushes against the caregiver, continues crying and does not settle sufficiently to return to play.

Reasoning. Separation distress alone cannot classify attachment. The reunion is more informative because it shows whether renewed access to the caregiver restores regulation. A secure child can seek and use comfort, then resume exploration. An avoidant child minimises overt proximity-seeking and may turn away or ignore the caregiver. This child instead maximises attachment signals: proximity is urgently sought, yet contact is mixed with anger or resistance and does not readily produce settling.

The simultaneous approach and resistance form a coherent strategy and therefore differ from **disorganisation**, where behaviour lacks a stable strategy and may include contradictory, apprehensive, frozen or disoriented sequences. The observed pattern describes this child's relationship with this caregiver in this procedure. It is not a diagnosis, a fixed personality trait or proof of inconsistent caregiving.

Answer. The pattern is ambivalent or resistant attachment, pattern C. The decisive clues are intensified proximity-seeking, resistance during contact and failure to use reunion effectively for soothing and renewed exploration.

23.4 Temperament and goodness of fit: the setting creates the difficulty

Presentation. A child watches an unfamiliar classroom activity from the edge, declines the teacher's first invitation and speaks little while everyone is watching. With advance warning, time

to observe and a quiet invitation from a familiar teacher, the child joins, completes the task and later participates comfortably. At home and with familiar peers, communication and play are fluent. The school has responded by calling the child stubborn and demanding immediate public participation.

Reasoning. Thomas and Chess directs attention to behavioural style before disorder. The initial response to novelty is withdrawal; the later capacity to join after familiarisation concerns adaptability. Because the response is cautious, improves gradually and is not described as an intense, persistently negative or globally dysregulated pattern, it is closer to a slow-to-warm-up temperament than to the classic difficult constellation.

The impairment is transactional. Immediate public performance and criticism create a poor **goodness of fit** by punishing the child's characteristic entry into novelty. Preparation, observation, phased entry and a familiar adult improve the fit without declaring that the child can avoid every demand. The cross-setting strengths argue against treating silence in the unfamiliar classroom as proof of a fixed language deficit, but persistent distress or impairment would still require assessment for anxiety, communication difficulty and other alternatives.

Answer. This is a **slow-to-warm-up temperamental pattern** made problematic by poor goodness of fit. Management should alter the transaction through predictable preparation and graduated participation while separately assessing any sustained impairment.

23.5 Theory application: supported performance is not a new Vygotskian stage

Presentation. A child cannot independently sort a mixed set of objects by a changing rule. The examiner first restates the goal, then models the opening move and finally uses a visual cue. With that graduated help, the child completes the remaining items and applies the same strategy to a related task. A static score records only the initial failure.

Reasoning. Vygotsky asks how performance changes through social mediation. The lower boundary is what the child can do independently; the upper boundary is what becomes possible with appropriate assistance. The distance between them is the zone of proximal development. The examiner's calibrated prompts are scaffolding: support is increased only as needed, then should be faded as responsibility transfers to the learner. Improvement that generalises to a related task is stronger evidence of an emerging strategy than completion achieved by simply giving the answer.

Piaget would ask what structure of classification or rule use the task requires, and a static assessment would describe unaided performance. Those are different questions. The assisted success must not be reported as independent mastery, but neither should the unaided failure be treated as the child's full learning capacity. Most importantly, Vygotsky proposed no age-linked stages, so the case cannot be labelled as entry into a new Vygotskian stage.

Answer. The case demonstrates performance within the **zone of proximal development**, revealed through **scaffolding** and dynamic assessment. Record independent and assisted

performance separately, fade help, test transfer and investigate language, attention, memory or task mismatch if well-calibrated support does not unlock performance.

- **RULE** **An expected age describes typical attainment; it is not a referral threshold.** A typical age cannot be promoted into a referral rule.
- **TRAP** **Do not report assisted success as independent mastery.** The assisted success must not be reported as independent mastery, but neither should the unaided failure be treated as the child's full learning capacity.

12 months TYPICAL INDEPENDENT WALKING AGE	9 to 15 months KAPLAN AND SADOCK EXPECTED RANGE	8.2 to 17.6 months WHO WALKING WINDOW (1ST TO 99TH CENTILE)	24 months TWO-WORD-PHRASE RED FLAG
--	--	--	---

HOLD THIS
A correct label without the intervening reasoning is fragile in a viva and unsafe in practice.

24 . Consolidation

This closing topic turns developmental psychiatry into a retrievable clinical map. The aim is not to recite every date in isolation. It is to recover **three** linked structures under examination pressure: which question each **developmental theory** asks, which observable functions belong to each **developmental domain**, and how the meaning of an age determines whether the doctor continues surveillance, screens or refers.

How to use this page. Learn the master mnemonic, cover the page and work through the answer-free prompts, then redraw the theory and milestone tracks from memory. On checking the redraw, be strict about construct.

- **TRAP** **A typical age is not a deadline.** A WHO **99th-centile** boundary is a screening signal rather than a diagnosis, and only an explicitly sourced red flag supports an age-linked referral call.

three LINKED STRUCTURES	five DEVELOPMENTAL DOMAINS	1st-to-99th-centile WHO GROSS-MOTOR CONSTRUCT
-----------------------------------	--------------------------------------	---




Fig 20.21 The whole-chapter synthesis map, drawn to be redrawn from memory. Theory lanes follow Berk, Mahler follows Wolberg, and Kohlberg sits in descriptive bands because no numeric ages are sourced for him. Vygotsky is drawn across the axis rather than on it, since his theory is not stage-based. Gross-motor windows are the WHO Motor Development Study, 1st to 99th centile; the language and gesture red flags are Developmental Psychopathology, stated for autism developmental evaluation. The plate carries one anchor per domain rather than the milestone tables, and names the domains for which no red-flag age is sourced.

24.1 The retrieval method: rebuild relationships, not isolated lists

Retrieve the map in a fixed clinical order. Start with the explanatory theories, move to observable development across domains, attach every age to its normative construct, and finish with the clinical action. This order prevents three common collapses: a theory stage becoming a milestone, an expected attainment becoming a referral threshold, and a positive screen becoming a diagnosis.

The theories answer different questions about the same developing person.

THEORY	QUESTION
Piaget	how cognition is organised
Erikson	which psychosocial conflict is foregrounded
Freud	how drive and conflict are organised
Mahler	how psychological separateness develops within relationship
Kohlberg	how moral decisions are justified
Vygotsky	what learning becomes possible through social mediation

HOLD THIS
Their temporal alignment is a recall aid, not evidence that the constructs are interchangeable.

24.2 Master mnemonic: MAP, MARK, MATCH, MOVE

MNEMONIC

MAP - MARK - MATCH - MOVE

MAP the theory lanes: cognition, psychosocial conflict, psychosexual organisation, separation-individuation and moral reasoning, with sociocultural mediation running across them. **MARK** observable development in the **five** domains: gross motor, fine motor, language and speech, cognitive or adaptive, and social and emotional. **MATCH** each age to its **normative construct**: median, expected attainment, expected range, upper limit, red flag or a source-stated age whose normative role remains unclear. **MOVE** from observation to action: describe the trajectory and quality, look for loss, asymmetry and multi-domain involvement, use screening as a risk filter, and refer for regression, a sourced red flag or a concerning neurological or sensory pattern.

The mnemonic contains the chapter's division of labour. The theory map explains; the milestone map describes; the construct check interprets; and the final verb commits the doctor to an action. If the answer jumps directly from an age to a diagnosis, one of the middle steps has been skipped.

24.3 Answer-free active recall

Cover the rest of the page and answer each prompt aloud or on paper. Check only after completing the retrieval attempt.

1. What distinct developmental question is asked by Piaget, Erikson, Freud, Mahler, Kohlberg and Vygotsky?
How would you reconstruct the source-stated age lanes for Piaget, Erikson and Freud without turning theoretical brackets into clinical thresholds?
What is the sequence of Mahler's separation-individuation account, and which two similarly named cognitive and relational constructs must not be confused?
How are Kohlberg's levels and stages ordered, and what exactly is scored in a moral dilemma?

Where should Kohlberg and Vygotsky sit on a theories-by-age map, and why must neither be given an invented numeric row?

2. Name the five developmental domains and describe how a limitation in one can alter performance in the others.

What do cephalocaudal and proximodistal organisation explain, and what do they not date?

Reproduce the WHO gross-motor achievement windows from the milestone plate and state the clinical meaning of the upper boundary.

Why is absence of hands-and-knees crawling alone an unsafe referral rule?

Trace the communication sequence from prelinguistic exchange through words, combinations, grammar and intelligibility, preserving the construct attached to each anchor.

Which communication findings are explicit age-linked red flags, and how would you evaluate a child who meets one?

3. How do **developmental delay**, **dissociation**, **deviance**, **plateau** and **regression** differ?

How do **surveillance**, **screening** and **formal developmental assessment** differ in purpose and output?

Which reunion behaviours distinguish secure, avoidant, ambivalent or resistant, and disorganised attachment?

Name the Thomas-and-Chess dimensions, distinguish their three constellations, and formulate one example of goodness of fit.

How do responsiveness and demandingness generate the parenting-style patterns, and why are outcome associations not deterministic?

4. How would you separate gender identity, gender expression, sexual orientation and sexual behaviour in a developmental assessment?

Why must Tanner stage be assigned from morphology rather than chronological age or psychological maturity?

How do dual-systems and maturational-imbalance accounts explain a difference between calm reasoning and behaviour in a highly charged peer context?

Which features of deprivation, later care and protective relationships determine whether risk becomes persistent impairment or recovery?

24.4 The synthesis redraw: theories above, milestones below, action at the base

Redraw the synthesis as aligned horizontal tracks. Across the top, place the age axis from infancy through adulthood. Give Piaget, Erikson and Freud separate lanes using the source-stated stage bands, and give Mahler a shorter early relational lane using only its sourced phase timings. Place Kohlberg in ordered descriptive bands rather than numeric ages. Draw Vygotsky beside the entire axis as a cross-cutting process of mediation, assisted performance, internalisation and self-regulation. The alignment should make temporal overlap visible while keeping each theory's explanatory target separate.

Below the theory lanes, draw the five developmental domains. In the gross-motor lane, reconstruct the WHO windows from sitting without support through walking alone and label the **1st-to-99th-centile** construct. In the fine-motor lane, recover the sequence from grasp and

transfer through pincer control, construction, scribbling and copied forms. In the language lane, move from vocalisation and gesture through meaningful words, combinations, grammar and intelligibility. In the cognitive or adaptive lane, place symbolic representation, problem-solving and theory-of-mind anchors. In the social and emotional lane, move from reciprocity and selective attachment through shared attention, symbolic play, peer engagement and regulation. Use the existing milestone and reflex plates to check the redraw rather than recreating their source-labelled tables in prose.

Add a separate pubertal branch rather than forcing it onto the childhood milestone ladder. Record Tanner morphology by component, not by age, and place identity formation, relationships, gender and psychosexual development beside the adolescent body and brain changes. This preserves the central distinction: physical maturation, cognitive capacity, moral reasoning, identity and consent are related but are not interchangeable measures.

Finally, draw the clinical action rail beneath every domain. Its sequence is *trajectory, pattern, construct, action*. Mark regression, plateau, asymmetry, abnormal quality, sensory or neurological findings and unevenness across domains before checking a date. Then label the date honestly as expected, a range, an upper limit, a red flag or unclear. End with the appropriate move:

- continue surveillance or apply a validated screen
- arrange comprehensive developmental evaluation

■ **RULE** **Developmental psychiatry is not the memorisation of ages, but the disciplined interpretation of change over time.** The completed redraw should make one principle unavoidable.

Previously asked

This chapter is the one place in the series where the honest answer about past questions is that the examiner has not asked them here. Normal development is examined, and examined often, but it is examined as *psychology* under Paper I rather than as child psychiatry under Paper III. Every developmental question on record in the catalogue sits in the basic sciences and psychology section: Erikson's psychosocial stages has appeared four times, most recently in 2021 and before that in 2017 and twice in 2012, always as a ten-mark question; Jean Piaget was asked in 2011; and psychological development in adolescence, paired with its relation to psychiatric disorders in adolescence, came up in 2018 and again in 2020. Those questions and their model answers live where they were asked, in the General and Developmental Psychology notes, and they are not repeated here.

That distribution is worth reading rather than shrugging at, because it tells you how the material is tested. The theories are examined as theories, on their own terms, in a psychology paper: name the stages, get the order right, get the count right, say what each stage is *for*. What this chapter adds is the part the examiner does not ask directly and the clinic asks constantly, which is the deviation call. No question on record asks for a milestone table. Plenty of clinical practice asks

whether this particular child, at this particular age, is outside the normal window, and that judgement rests on knowing which of the ages you memorised is a typical attainment and which is a screening boundary. A candidate who can recite Piaget's four stages and still cannot say whether a fourteen-month-old who is not walking needs referring has learned the examinable half and missed the useful half.

ON THE THEORIES, EXAMINED UNDER PAPER I PSYCHOLOGY

Erikson and his psychosocial stages is the highest-yield developmental question in the catalogue, at four appearances for ten marks. Piaget follows. Both reward the same discipline: the stages in order, at their exact count, each with its central conflict or operation and its one clinical use. The age bands are the least reliable part of both answers and should be given as the approximate, source-dependent ranges they are. MUHS 2011, 2012, 2017, 2021

ON ADOLESCENCE, EXAMINED UNDER PAPER I PSYCHOLOGY

Psychological development in adolescence is asked with its psychiatric relation attached, which is the examiner signalling that normal adolescent development is the setup and the disorders are the payoff. The normal half is owned here: identity formation, the neurodevelopment of adolescence, the imbalance between a fast-maturing reward system and a slow-maturing control system, and why risk-taking is developmentally expected rather than pathological. The disorder half belongs to the disorder chapters. MUHS 2018, 2020

WHAT HAS NEVER BEEN ASKED, AND WHAT TO DO ABOUT IT

There is no question on record on the milestone tables, on the primitive reflexes, on attachment classification, on temperament, or on the developmental screening instruments. Two readings are available and both are worth holding. The first is that these are low-yield for the written paper and should be learned to a clinical rather than an examination standard. The second is that they are exactly the material a viva or a clinical case can turn on without warning, because they are what a child psychiatrist is assumed to know rather than asked to prove. Prepare the theories for the paper. Prepare the deviation call for the patient.

Quick review

DEVELOPMENTAL PSYCHIATRY: THEORIES, MILESTONES AND THE DEVIATION CALL	
THE INTERPRETIVE RULE	
CONSTRUCT BEFORE ACTION	Median means half have attained the skill; expected means usually attained; expected range is a normal window; upper limit is the outer normal bound; and red flag warrants further evaluation. If the source leaves the construct unclear, repeat only its own wording. Never turn it into a referral threshold.
WHAT EACH THEORY ASKS	
PIAGET	How is cognition organised? Sensorimotor action becomes preoperational representation, concrete operations bring reversible logic for tangible material, and formal operations extend reasoning to propositions and possibilities. The stage bands are theoretical descriptions, not milestone norms or referral ages.
ERIKSON	Which psychosocial conflict is foregrounded? Trust versus mistrust, autonomy versus shame and doubt, initiative versus guilt, industry versus inferiority, identity versus role confusion, intimacy versus isolation, generativity versus stagnation, and integrity versus despair. Use the conflict and strength for formulation, not diagnosis.
FREUD	How are drive, bodily focus and conflict organised? Oral, anal, phallic, latency and genital. Fixation is an enduring investment in an earlier mode; regression is movement back toward it under stress. These are historical psychosexual mappings, not developmental screening stages or Tanner stages.
MAHLER	How does psychological separateness develop within relationship? Distinguish separation, "I am not you," from individuation, "I am me." Object constancy is the affective capacity to retain a whole loved person across absence and frustration; it is not Piagetian object permanence.
KOHLBERG	How is a moral judgement justified? Preconventional reasoning moves from punishment to exchange; conventional reasoning from approval to social order; postconventional reasoning evaluates law through rights and principles. Score the reason, not the verdict. Kaplan and Sadock gives only descriptive bands: before adolescence, school-aged children, some young adolescents, during adolescence, later in adolescence, and late adolescence or early adulthood. There are no sourced numeric ages.

DEVELOPMENTAL PSYCHIATRY: THEORIES, MILESTONES AND THE DEVIATION CALL	
VYGOTSKY	What becomes possible through social mediation? The zone of proximal development lies between independent and assisted performance; scaffolding is contingent support that is faded; social speech becomes private and inner speech. Vygotsky proposed no stages, so he belongs beside an age axis as a cross-cutting mechanism.
LANGUAGE ACQUISITION	Skinner: operant contingencies shape verbal behaviour. Chomsky: innate grammatical preparedness supports generative language. Vygotsky: shared speech is socially mediated and internalised. Consequence, grammar and mediation answer different parts of the same problem.
ATTACHMENT	Bowlby asks how proximity to a familiar caregiver provides safe haven and secure base; internal working models carry expectations of self, others and help. In Ainsworth's Strange Situation, reunion distinguishes secure use of comfort, avoidant minimisation, ambivalent or resistant approach with resistance, and disorganised absence of a coherent strategy. These are relationship patterns, not diagnoses.
TEMPERAMENT AND PARENTING	Thomas and Chess ask how activity, rhythmicity, approach or withdrawal, adaptability, threshold, intensity, mood, distractibility, and attention span or persistence fit the environment. Easy, difficult and slow-to-warm-up are constellations; outcome depends on goodness of fit. Baumrind maps responsiveness and demandingness: authoritative is high-high, authoritarian low-high, permissive high-low, and uninvolved low-low.
ADOLESCENCE AND ADVERSITY	Adolescence is asynchronous change in body, reward, control, identity and relationships. Dual-systems and maturational-imbalance models explain context-sensitive risk, not global irrationality. Marcia separates exploration from commitment. Deprivation removes expected input; threat adds danger. Resilience is better-than-expected adaptation produced by dynamic child-context processes, especially stable responsive care, not an invulnerable trait.
SOURCED MILESTONE ANCHORS	
WHO GROSS MOTOR WINDOWS	The WHO 1st-to-99th-centile windows are: sitting without support 3.8 to 9.2 months; standing with assistance 4.8 to 11.4 months; hands-and-knees crawling 5.2 to 13.5 months; walking with assistance 5.9 to 13.7 months; standing alone 6.9 to 16.9 months; walking alone 8.2 to 17.6 months. Beyond an upper boundary signals the need for appropriate screening. It is not a diagnosis or an independently sourced referral threshold.

DEVELOPMENTAL PSYCHIATRY: THEORIES, MILESTONES AND THE DEVIATION CALL	
FINE MOTOR	Berk expects a well-coordinated pincer grasp by the end of the first year , vigorous scribbling between 10 and 21 months , and a tower of two cubes between 10 and 19 months . These are expected attainments or ranges, not referral thresholds. Observe vision, posture, symmetry, tone, comprehension and opportunity.
LANGUAGE AND SPEECH	Kaplan and Sadock expects cooing at about 3 to 4 months , meaningful words at about 12 months , and word combinations at 18 months . Speech is expected to be about 25 percent intelligible to strangers at 18 months , 50 percent at 24 months , 75 percent at 36 months , and nearly 100 percent by 4 years . These are expected values, not referral cuts.
SOCIAL AND COGNITIVE	Berk gives the social-smile window as 6 to 10 weeks . Joint attention and pointing are expected around 12 months in the New Oxford Textbook of Psychiatry; Dulcan expects symbolic play at 18 months ; explicit false-belief performance is expected at 4 or 5 years in the cited account. None is an automatic referral threshold.
PUBERTY AND TANNER	Tanner stage is morphological: record breast or genital development and pubic hair separately, never infer stage from age or psychological maturity. Berk's expected ranges include breast budding at 8 to 13 years , testicular enlargement at 9 to 13.5 years , menarche at 10.5 to 14 years , and spermatarche at 12 to 16 years . These are expected ranges, not clinical cut-offs.
SCREENING SIGNALS, RED FLAGS AND REFLEXES	
WHO SCREENING SIGNAL	The gross-motor windows describe variation and signal screening when a child appears late. About 4.3 percent of children in the WHO study never exhibited hands-and-knees crawling, so absence of crawling alone is not a red flag. Judge progression, transitions, symmetry, tone and movement quality.
COMMUNICATION RED FLAGS	Arrange prompt developmental evaluation for no babbling or gesturing at 12 months , no single words at 16 months , no spontaneous two-word phrases at 24 months , or loss of language or social skills at any age . These Developmental Psychopathology thresholds are referral signals, not diagnoses.
EXPECTED REFLEX DISAPPEARANCE	Keep the source attached: rooting at 3 weeks in Berk; sucking after 4 months in Berk; palmar grasp at 3 to 4 months in Berk; stepping at 2 months in Hutchison's Clinical Methods; asymmetric tonic neck reflex at 4 months in Berk; Moro at 12 weeks in Tasman but 6 months in Berk and Hutchison. Do not average disagreement.

DEVELOPMENTAL PSYCHIATRY: THEORIES, MILESTONES AND THE DEVIATION CALL	
BABINSKI AND REFLEX RED FLAGS	Berk gives Babinski disappearance at 8 to 12 months ; Hutchison gives the extensor plantar response an upper limit of 18 months and calls persistence beyond 2 years a red flag. Hutchison also flags absence of sucking beyond the 36th week of gestation and gives persistence of primitive reflexes beyond the fourth month of life as a general warning. Reflex-specific expectations, symmetry, tone and the neurological examination still govern interpretation.
THE DEVIATION CALL	
NAME THE TRAJECTORY	Delay is slow acquisition with broadly preserved sequence; dissociation is uneven development between domains; deviance is an unusual sequence or quality; plateau is failure to acquire new skills; regression is loss of a convincingly acquired skill. Also act on marked asymmetry, abnormal tone or movement, sensory concern, neurological findings and convergent multi-domain difficulty.
SURVEILLANCE TO ASSESSMENT	Surveillance repeatedly integrates concerns, history, observation, examination and trajectory. Screening applies a standardised risk filter and does not diagnose. Assessment characterises strengths and impairments across domains and investigates cause. A negative screen never overrules regression, a sourced red flag, abnormal examination or persistent concern.
INDIAN PATHWAY	The Trivandrum Development Screening Chart covers birth to 6 years ; one item delay is the positive screen termed TDSC delay. DASII yields Motor DQ and Mental DQ; Indian follow-up practice uses DQ below 85 as a delay marker. A positive screen or quotient is not a diagnosis.
FINAL ACTION	Use trajectory, pattern, construct, action . Continue surveillance for a coherent progressing course within an expected window; screen when a WHO upper boundary is crossed; arrange comprehensive evaluation for regression, an explicit red flag, marked asymmetry, abnormal quality, neurological or sensory findings, multi-domain delay, or persistent clinical concern.

References

Normal development is a chapter where the sources disagree with each other more than they admit, and the reference list is built to show that rather than hide it. Every age, stage boundary, reflex age and instrument property in these notes was taken from a named source and checked against it; where two standard texts give different bands for the same stage, both are reported and neither is presented as the answer. The depth bar for the theories is Berk's *Development Through the Lifespan*, which is the source that aligns Freud, Erikson and Piaget on shared boundaries and therefore the source the alignment plate follows. Tasman and Kaplan and Sadock carry the

psychiatric framing, Lewis and Cicchetti's *Developmental Psychopathology* the child-psychiatry depth and the language red flags, Wolberg the Mahler subphases, Hutchison's *Clinical Methods* the primitive reflexes and their persistence, Schunk's *Learning Theories* the Vygotskian material, and Kaufman's *Clinical Neurology for Psychiatrists* the early-handedness sign. The Indian frame comes from the Indian Psychiatric Society child and adolescent guideline and from the primary validation literature for the Indian instruments.

Two sources carry more weight than the rest and deserve naming for what they are. The WHO Motor Development Study is the only source in this chapter that states the normative role of its own numbers: its windows are bounded by the 1st and 99th centiles, it followed children in Ghana, India, Norway, Oman and the USA, and it says in its own words that the windows exist to signal the need for screening rather than to diagnose. Every gross-motor boundary taught here comes from it, and comes labelled as what it is. The Trivandrum validation is the only Indian screening instrument in this chapter with published psychometrics, and its sensitivity, specificity and predictive values are reported from the primary paper rather than from a secondary summary.

Three absences are load-bearing and are stated rather than papered over. Nelson's *Textbook of Pediatrics*, the source a paediatric milestone chapter would normally lean on, was not available; the milestone ages here come from the developmental psychology and psychiatry texts instead. No Indian Academy of Pediatrics milestone chart was available, so no IAP milestone age is stated anywhere in this chapter. The DASII manual was not available, so its administration range, item structure and norming sample are not stated; only its two developmental quotients and the delay marker used in Indian follow-up practice are given, from the applied literature. Where a value could not be sourced, it was left out.

SOURCE HIERARCHY

1. Berk LE. *Development Through the Lifespan*. The alignment of Freud, Erikson and Piaget on shared age boundaries, the milestone ages, the pubertal windows and the social-emotional sequence.
2. Berk LE. *Child Development*, 8th edition. Supporting developmental detail.
3. Tasman A, et al. *Psychiatry*. The psychiatric framing of the theories, the Tanner staging tables and the alternative stage bands that disagree with Berk.
4. Sadock BJ, Sadock VA, Ruiz P. *Kaplan and Sadock's Comprehensive Textbook of Psychiatry*, 2024. Kohlberg's levels as descriptive bands, Mahler, the intelligibility gradient and the motor-delay allowance.
5. Cicchetti D, ed. *Developmental Psychopathology*. The sourced language and social red flags and their referral ages.
6. Martin A, Volkmar FR, Bloch M, eds. *Lewis's Child and Adolescent Psychiatry*. Child-psychiatric development and the separation-anxiety window.
7. Dulcan MK, ed. *Dulcan's Textbook of Child and Adolescent Psychiatry*. Developmental surveillance framing.
8. Rutter M, et al. *Rutter's Child and Adolescent Psychiatry*, 6th edition.
9. Wolberg LR. *The Technique of Psychotherapy*. Mahler's separation-individuation subphases and their ages.
10. Schunk DH. *Learning Theories: An Educational Perspective*, 6th edition, 2011. Vygotsky's sociocultural theory, the zone of proximal development, scaffolding and private speech.
11. Hutchison R, Glynn M, eds. *Hutchison's Clinical Methods*. The primitive reflexes and the ages beyond which persistence is a warning sign.
12. Kaufman DM, Milstein MJ. *Kaufman's Clinical Neurology for Psychiatrists*. Early unequivocal handedness as a warning sign in infancy.
13. American Psychiatric Association. *DSM-5-TR*, 2022. Speech intelligibility and the late-acquired sound allowance.

14. Indian Psychiatric Society. *Clinical Practice Guidelines for Child and Adolescent Psychiatry*. The Indian assessment and screening landscape.
15. Ciccarelli SK, White JN. *Psychology*, 5th edition. Supporting developmental psychology.
16. Gelder M, Andreasen N, Lopez-Ibor J, Geddes J, eds. *New Oxford Textbook of Psychiatry*.

PRIMARY PAPERS, EVERY IDENTIFIER VERIFIED

1. WHO Multicentre Growth Reference Study Group. WHO Motor Development Study: windows of achievement for six gross motor development milestones. *Acta Paediatr Suppl* 2006;450:86-95. PMID 16817682. The 1st and 99th centile windows for the six gross motor milestones, from 816 children in Ghana, India, Norway, Oman and the USA. The only construct-explicit motor source in this chapter.
2. Nair MKC, Harikumar Nair GS, George B, Suma N, Neethu C, Leena ML, Russell PSS. Development and validation of Trivandrum Development Screening Chart for children aged 0-6 years. *Indian J Pediatr* 2013;80(Suppl 2):S248-55. PMID 24014206. Fifty-one items, validated against the Denver test in 1,183 children, with its sensitivity, specificity, predictive values and reliability.
3. Shekhawat DS, Gupta T, Singh P, Sharma P, Singh K. Monitoring tools for early identification of children with developmental delay in India: an update. *Child Neuropsychol* 2022;28(6):814-830. PMID 35722754. The Indian screening landscape, the Baroda test as the first Indian tool in 1991, and the estimated prevalence of developmental delay in India.
4. Kadam AS, Nayyar SA, Kadam SS, Patni BC, Khole MC, Pandit AN, Kabra NS. General Movement Assessment in babies born preterm: Motor Optimality Score-Revised, trajectory, and neurodevelopmental outcomes at one year. *J Pediatr X* 2022;8:100084. PMID 37334030. The DASII motor and mental developmental quotients in Indian preterm follow-up.
5. Shukla A, Bandyopadhyay T, Vallamkonda N, Maria A. Long-term neurodevelopmental outcomes of neonatal chikungunya: follow-up of a series of cases till one year. *J Trop Pediatr* 2021;67(3). PMID 32756979. The DASII delay marker in Indian practice.

PART II



Intellectual Disability and Autism Spectrum Disorder

Intellectual disability and autism: how each is defined and assessed, what lies beneath them, and what can actually be done.

-
- 01 Intellectual disability: the concept and its measurement
 - 02 The causes of intellectual disability
 - 03 Living with intellectual disability
 - 04 Autism: concept, causes and criteria
 - 05 The clinical picture of autism and how it is recognised
 - 06 Managing autism and its outcome
 - 07 Discriminate, apply and consolidate

PAPER III

Intellectual disability & autism spectrum disorder

Six bands . one complete notes document

Intellectual disability: the concept and its measurement

1 · The neurodevelopmental frame

Begin with the person, not the label. Intellectual disability and autism spectrum disorder are developmental presentations that demand careful formulation rather than rapid categorisation. The marks lie in showing that one can move from a concern raised by a family or school to a coherent account of abilities, everyday functioning, communication, behaviour, context and need. A diagnosis may organise that account, but it does not replace it.

This chapter is best read as a clinical sequence. It first establishes the language and measurement needed to describe intellectual disability, then considers causes, associated psychiatric and practical concerns, and care across settings. It then develops autism spectrum disorder from its diagnostic framework through recognition, assessment, intervention and outcome. The final task is discrimination: deciding which explanation best fits the observed developmental pattern and what the distinction changes for the person and family.

ID CONCEPT AND MEASUREMENT	ASD RECOGNITION AND FORMULATION	PERSON FUNCTION BEFORE LABEL
--------------------------------------	---	--




Fig 21.1 The neurodevelopmental frame. The chapter's advance organiser: age runs across, the axis of impairment runs down, and every entity the chapter teaches has a place on both. Global developmental delay is drawn with a dashed border because it is a placeholder, not a diagnosis, and it forks on reassessment; intellectual developmental disorder and autism spectrum disorder sit in different lanes and are joined by a dashed rule because they are diagnosed together rather than ranked; borderline intellectual functioning sits on the same global axis as ID but above the diagnostic threshold, where DSM-5-TR calls it a condition and not a disorder. The age axis is ordered rather than scaled, and carries the single boundary the sources give.

1.1 The organising clinical question

A **developmental formulation** is a working explanation of how a person has developed, what they can do in ordinary settings, what creates difficulty, and what may help. It is deliberately

broader than a diagnostic label. It prevents a clinician from mistaking a test result, a conspicuous behaviour, or a single historical detail for the whole clinical picture.

■ **RULE** **Formulate functioning in context before using a diagnostic label to organise the next step.** A label becomes clinically useful only when it clarifies the pattern of difficulty, the questions still to be answered, and the support that may be needed.

At the outset, hold several strands together:

- the **presenting concern** in the words of the person, family, school or referrer;
- a **longitudinal history** that asks how the current pattern emerged and changed;
- the person’s **functional profile** across their actual environments; and
- the **contextual formulation** of relationships, opportunity, stress, culture and available support.

Developmental expectations belong in *the Normal child development and milestones notes*. A full child psychiatric assessment, service response and safeguarding approach belong in *the Child psychiatry: assessment, treatment, services and protection notes*. Those cross-references matter because a developmental formulation is only as good as the history and context on which it rests.

1.2 A map of the chapter

The chapter separates related questions so that each can be answered precisely. The following map is a reading aid, not a diagnostic algorithm.

CLINICAL QUESTION	CHAPTER FOCUS	WHY THE DISTINCTION MATTERS
How should intellectual ability and everyday functioning be described?	Concept, criteria, assessment and adaptive behaviour	Description needs interpretation in the person’s life context.
What may account for the developmental presentation?	Causes and syndrome-specific discussions	Explanation can alter investigation, anticipatory care and counselling.
Is autism spectrum disorder the best account of the observed developmental pattern?	Diagnostic framework, clinical picture and assessment	Recognition requires a pattern-based judgement rather than an isolated feature.
What should change in the person’s everyday life?	Interventions, health review, family support and prognosis	A support plan translates formulation into feasible action.

The questions interact, but they should not be collapsed. A person may need more than one explanatory frame, and the clinician must decide what each frame adds. Assessment methods, severity language, aetiological work-up and intervention are therefore developed in their own sections rather than treated as interchangeable.

■ **TRAP** **Examination trap:** do not let an early label replace description of the observed problem. Establish the pattern first, then use the relevant framework to explain it.

1.3 Reading findings without reductionism

GOOD TO REMEMBER

Interpret clinical information together: history may reveal a developmental trajectory; direct observation may show communication and response to change; assessment may characterise ability or practical skill; and collateral may reveal demands invisible in the consulting room. Check **ecological validity**: the account should fit the person's life outside the clinic as well as the clinician's impression inside it.

Use the chapter to ask progressively sharper questions: What is the pattern of strength, difficulty and participation? What information would materially change the formulation? Which co-occurring concerns require active recognition rather than attribution to the developmental diagnosis? What support is needed now, and what should be reviewed as circumstances change?

This approach resists **diagnostic reductionism**: the error of allowing one diagnosis to explain every difficulty, behaviour or need. It also makes room for the person's preferences and the family's realities. Good developmental psychiatry is exact about constructs while remaining practical about lives.

1.4 Master mnemonic: MAP

A compact way to retain the chapter. Use **MAP** as a sequencing aid when approaching a developmental presentation. It is not a substitute for assessment; it is a safeguard against beginning in the wrong place.

MNEMONIC

MAP

Map the developmental and functional story.

Ask which framework best explains the pattern.

Plan support around the person, family and participation.

The first step keeps the account anchored in lived function. The second directs the clinician to the chapter's conceptual and diagnostic material without forcing a premature conclusion. The final step prevents assessment from becoming an endpoint: the relevant outcome is a more useful, humane and workable response.

The durable answer in developmental psychiatry is a formulation that explains function, recognises pattern and directs support.

2 · Global developmental delay concept

Global developmental delay is a clinical description for an early developmental presentation, **not a shortcut to a settled prognosis**. It is used when a child has not acquired expected developmental abilities across several areas, but the clinician cannot yet make a reliable statement about the severity of intellectual impairment. The label is valuable precisely because it

preserves that uncertainty. It allows the record to acknowledge a real developmental concern, organise the next clinical steps, and communicate need without pretending that a premature measurement has answered the longer-term question.

The topic is frequently examined because it tests clinical reasoning at the boundary between observation and diagnosis. Candidates must distinguish a broad developmental delay from a definitive intellectual-developmental diagnosis, explain why early assessment can be limited, and avoid converting a description of current functioning into a prediction about lifelong ability. In DSM-5-TR, global developmental delay is recorded with the code **F88**. The code should serve the formulation, not replace it.

F88 DSM-5-TR GDD CODE	below 5 years DSM-5-TR AGE USE	between 3 months and 5 years INDIAN GUIDELINE WINDOW	at least 2 domains OPERATIONAL THRESHOLD
---------------------------------	--	--	--

2.1 What the label is for

Global developmental delay is reserved in DSM-5-TR for a child **below 5 years** when the clinical severity of intellectual impairment cannot yet be assessed reliably. This is not merely a matter of a child being young. The label is appropriate when the child has failed to achieve expected milestones in several areas of intellectual functioning and cannot undergo a systematic assessment that would permit a dependable severity judgement. Developmental variation, limited cooperation, sensory or physical difficulties, and the practical limits of formal assessment can all make an apparently precise conclusion unsound.

The key idea is **assessability**. A clinician may have persuasive evidence that development is not proceeding as expected while lacking a valid way to quantify the enduring intellectual and adaptive picture. The correct response is neither to deny the concern nor to force a severity label. Global developmental delay creates an honest interim formulation: there is a clinically meaningful developmental problem, its pattern must be described, and the formulation must remain open to revision.

■ **RULE** **Defining principle.** Use global developmental delay when delay is evident but a reliable developmental severity judgement cannot yet be made; make reassessment part of the diagnosis from the outset.

GOOD TO REMEMBER

Families often arrive seeking a definitive answer after a milestone concern, a referral from an educational setting, or an earlier clinical opinion. The psychiatrist must hold uncertainty without sounding evasive. Explain what has been observed, what remains difficult to judge, and what information will make the formulation more robust. A diagnostic label should reduce confusion, not offer a false certainty that later requires withdrawal.

Developmental expectations and their usual sequence belong in *the Normal child development and milestones notes*. The full approach to history-taking, examination, assessment planning, services and protection belongs in *the Child psychiatry: assessment, treatment, services and protection notes*. This section focuses on the conceptual work: when to use the label, what it means, and what it does not mean.

2.2 Why early developmental uncertainty is clinically real

Early childhood is a period of rapid acquisition, uneven performance and changing demands. A child may demonstrate an ability in a setting but not another, or show an emerging skill that cannot yet be relied upon in everyday life. Conversely, a striking delay may be clear before the clinician can determine whether it reflects a broad and persistent intellectual-developmental difficulty, a transient lag, a limitation of the available assessment, or a combination of these. The purpose of global developmental delay is to represent this clinical state accurately.

IN INDIA

The Indian guideline makes the same practical point: although intellectual disability may be recognisable early in development, it is often difficult to diagnose it accurately before 5 years of age. This is why the guideline uses global developmental delay as a **surrogate marker between 3 months and 5 years**. “Surrogate” should be read carefully. It means the term can organise recognition and follow-up during a period when the final intellectual-developmental formulation is not yet reliable. It does not mean that the two labels are interchangeable.

Diagnostic uncertainty is not diagnostic inaction. A useful **developmental formulation** states the abilities that are emerging, the activities that remain difficult, the contexts in which difficulty is most evident, and the collateral information needed to clarify the pattern. It also records uncertainty explicitly. This protects against a common clinical drift in which a provisional descriptive term is treated as though it had already settled questions about intellectual ability, adaptive functioning, cause or prognosis.

A child who cannot complete a systematic assessment should not be described as “untestable” and then left outside a formulation. The more accurate question is what has prevented a valid assessment on this occasion and how the clinician can obtain a better account of function over time. That may involve observation in familiar settings, informed collateral accounts, repeated contact, and attention to communication, motor, sensory and behavioural barriers. The methods and interpretation of intelligence testing are dealt with in the chapter’s assessment material; the present point is that a score is not required before a developmental concern can be named responsibly.

■ **TRAP Examination trap:** do not write that global developmental delay is simply “intellectual disability in a young child”. It is the appropriate label when the developmental concern is clear but the enduring degree of intellectual impairment cannot yet be judged reliably.

2.3 Operationalising a broad developmental delay

The Indian guideline cites **Shevell et al. (2008)** for an operational definition. In that approach, significant delay must be present in **at least 2** developmental domains. This is an operational framework for describing global delay, not a substitute for the diagnostic rationale about why a reliable severity assessment cannot yet be made. In an answer, it is good practice to name the framework being used rather than silently combining criteria from different systems.

The listed areas form a **developmental profile**, not a checklist to be ticked without context. A delay in a domain may affect performance in another. For example, limited motor opportunity can alter what is observed in self-care, and reduced communication can obscure social participation. The clinician’s job is to decide whether the observed pattern reflects broad developmental delay rather than merely adding isolated concerns together.

DEVELOPMENTAL AREA	QUESTION FOR THE CLINICAL FORMULATION	COMMON REASONING ERROR TO AVOID
Gross and fine motor	What movement and hand-use abilities are established in ordinary routines?	Assuming that limited opportunity and intrinsic delay are automatically the same.
Speech and language	How does the child understand, express and use communication in context?	Equating a language difficulty with global delay without reviewing the broader profile.
Social and personal functioning	How does the child participate with familiar people and manage age-expected interactions?	Explaining every social difficulty through a single diagnostic label.
Cognition	How does the child learn, solve everyday problems and generalise an acquired ability?	Treating an isolated observed success or failure as the whole cognitive account.
Activities of daily living	What can the child do in dressing, feeding, routines and other everyday tasks?	Ignoring environmental support or opportunity when judging independence.

The table is deliberately descriptive. It does not convert the domains into a severity scale, an intelligence test, or a statutory disability assessment. Those are different constructs and are treated elsewhere in the chapter. For this topic, the useful result is a coherent account of which developmental abilities have not emerged as expected, which strengths are already present, and why the available evidence supports a broad developmental concern.

- Describe abilities in the child’s actual routines rather than using only a global label.
- Identify whether delays are spread across the developmental profile or confined to a more limited concern.
- Record factors that may make observed performance underestimate the child’s capacity on that occasion.
- State what should be reviewed when more reliable developmental information becomes available.

2.4 A provisional formulation, not a forecast

Global developmental delay is often clinically adjacent to intellectual developmental disorder, but adjacency is not identity. DSM-5-TR notes that some children younger than 5 years whose later presentation meets criteria for intellectual developmental disorder first meet criteria for global developmental delay. This is an account of a possible developmental pathway, not a rule that every child with global delay will later receive that diagnosis.

The Indian guideline is explicit about **prognostic non-equivalence**: global developmental delay is not an inevitable precursor to intellectual disability. Some children described as having global delay will not have cognitive deficits that persist as intellectual disability. This caution is more than a prognostic nicety. If the early label is treated as a permanent verdict, the family may be given an unnecessarily narrowed view of the child's possibilities, while the clinician may stop looking for change, emerging skills or a more differentiated explanation.

Equally, the opposite error is harmful. Because some developmental delays may change over time, clinicians should not minimise a broad delay in the hope that it will resolve without review. A longitudinal formulation balances both risks: it recognises current impairment and need, avoids premature permanence, and specifies the questions that subsequent assessment must answer. It is a way of being precise about the present without claiming ownership of the future.

MNEMONIC

PAUSE when the early developmental picture feels uncertain

- Profile the child's functioning in ordinary settings.
- Ask whether current assessment is genuinely valid.
- Use uncertainty honestly rather than forcing a severity label.
- See whether delay extends across developmental areas.
- Evaluate again as the profile becomes clearer.

This is not a substitute for assessment. It is a guard against the familiar tendency to make the label carry more certainty than the evidence can support. The careful clinician does not wait passively for certainty; the clinician makes the uncertainty visible, develops a plan to reduce it, and documents the child's current needs in the meantime.

2.5 Reassessment is part of the diagnosis

The DSM-5-TR conditions of use for global developmental delay include **reassessment** after time has passed. That is not an administrative afterthought. The diagnosis has been chosen partly because present methods cannot support a reliable severity judgement; repeating the same unsupported conclusion at later contacts defeats its purpose. Reassessment asks whether the developmental trajectory has clarified, whether systematic assessment has become feasible, and whether the child's everyday functioning now supports a more definite formulation.

Good documentation makes this possible. It separates what was directly observed from what was reported, identifies the developmental areas of concern, records strengths and contextual supports, and states why the assessment could not yet be definitive. It also avoids language that

freezes a child into an early label. “Current presentation is consistent with global developmental delay; reassessment is planned as developmental information becomes more reliable” is clinically more useful than either an unsupported categorical diagnosis or a vague note that offers no plan. The review is not solely a search for a new diagnostic name. It should ask whether the formulation has become more specific, whether the child’s functional needs have changed, and whether the initial pattern needs to be reconsidered. A broad developmental concern can coexist with communication, motor, sensory, medical or contextual contributors that alter what is seen. The fuller developmental and psychiatric assessment belongs in *the Child psychiatry: assessment, treatment, services and protection notes*; this label should prompt that work rather than close it down.

■ **TRAP** **Examination trap:** “provisional” does not mean “wait and see without a plan”. It means document the current developmental concern, address present need, and arrange the reassessment that the provisional formulation requires.

2.6 Coding boundaries and classification caution

Classification systems do not always provide a direct vocabulary correspondence for the same clinical presentation. DSM-5-TR supplies global developmental delay as F88 for the early-childhood situation described above. By contrast, the **ICD-11 CDDR** contains no category named global developmental delay. The safe clinical conclusion is not to invent a correspondence. Describe the child’s developmental presentation clearly, use the classification framework required in the setting, and record why diagnostic certainty is limited.

A related DSM-5-TR category is **Unspecified Intellectual Developmental Disorder**, coded **F79**. It is reserved for exceptional circumstances in an individual above 5 years when locally available procedures cannot establish the degree of intellectual developmental disorder because associated impairments or severe problem behaviours make this difficult or impossible. It also requires reassessment. This is a different problem from the expected early-childhood uncertainty represented by global developmental delay, and the age boundary should not be reversed in an examination answer.

CLINICAL SITUATION	LABELING LOGIC	ESSENTIAL CAUTION
Early childhood with broad developmental concerns and unreliable severity assessment	DSM-5-TR global developmental delay, F88, for a child below 5 years.	It is a provisional developmental formulation that needs review.
Later assessment made impracticable by associated impairments or severe problem behaviours	DSM-5-TR unspecified intellectual developmental disorder, F79, in exceptional circumstances above 5 years.	Do not use it as a routine substitute for a proper assessment.
ICD-11 terminology	The CDDR has no category named global developmental delay.	Do not assert an unverified direct counterpart.

Kaplan and Sadock's summary describes an ICD-11 provisional disorder-of-intellectual-development category for a child below 4 years when there is evidence of disorder but it is difficult to know whether observed impairments may be transient. This is a useful reminder of the underlying clinical problem, but it does not resolve the naming discrepancy in ICD-11. The relevant **provisional category** reflects uncertainty about whether early impairment will persist, rather than a licence to treat every early delay as a definite disorder of intellectual development.

2.7 Clinical communication and the referral-to-support link

The value of recognition is practical. Identifying global developmental delay gives the clinician a shared language for referral, follow-up and communication with caregivers. It tells other professionals that developmental concerns extend beyond an isolated complaint and that the child needs an organised account of function. It also preserves the imperative to review the formulation as the child becomes more assessable. That combination of recognition and openness is the clinical strength of the term.

Conversations with families should be clear about what is known and what is not. Avoid implying that a descriptive early label proves a fixed lifelong outcome. Avoid the opposite reassurance that a broad delay needs no active response because the future is uncertain. Explain that the current pattern warrants attention, that abilities and needs should be described in everyday terms, and that later reassessment can refine the understanding. The family's observations of what the child can do, what needs assistance, and which settings make participation harder are central clinical data, not decorative collateral.

Why identifying global developmental delay matters for referral and **early intervention** is developed in the management section of this chapter. The present diagnostic task is more limited: make the developmental concern visible early enough that appropriate support is not postponed while diagnostic certainty is still evolving. The service model, its evidence and its Indian delivery context belong to that owner section and should not be inferred from the label alone.

- Communicate the present developmental profile in plain, functional language.
- State clearly that global developmental delay is an interim formulation rather than a prediction of a fixed outcome.
- Document the reason a definitive severity judgement is not yet reliable.
- Build reassessment into the plan and record what information should clarify the next formulation.

Global developmental delay names a genuine broad developmental concern while protecting the child from a severity judgement that the available evidence cannot yet support.

3 · Definition and diagnostic criteria of intellectual disability (DSM-5 and ICD-11)

The diagnostic task is to establish a developmental disorder of functioning, not to attach a label to a low test score. Intellectual disability is diagnosed when limitations in intellectual

functioning, adaptive functioning, and developmental onset form the same clinical picture. The criteria are deliberately conjunctive: each protects against a different error. Intellectual testing without functional assessment can medicalise a score; functional difficulty without developmental onset may represent an acquired disorder; and a developmental history without evidence of current limitation does not itself establish the diagnosis.

For examination answers and clinical records, the useful sequence is therefore simple: establish the nature of the intellectual limitation, establish its consequences for everyday independence and social responsibility in the person's sociocultural setting, and establish that the pattern began in development. The DSM formulation calls the condition **intellectual developmental disorder**, with intellectual disability retained in parentheses; the ICD formulation uses the equivalent term disorders of intellectual development. The difference in wording should not obscure their shared clinical core.




Fig 21.2 The three-criterion structure of intellectual disability. Criteria A, B and C are joined by a conjunction, not a menu: DSM-5-TR requires all three, and the two internal rules that decide A and B are examined more often than the criteria themselves. The ICD-11 6A00 strip reads straight down from each DSM column so the one real divergence stands out.

3.1 What the diagnosis is designed to capture

The diagnosis is about a persistent limitation in the person’s capacity to understand, learn, reason and manage the ordinary demands of life, where that limitation has developmental origins. It is

neither a synonym for educational underachievement nor a shorthand for disadvantage, illness, sensory disability, behavioural disturbance, or inadequate opportunity. Each of these may complicate assessment, and some may impair performance in a testing situation, but none can substitute for the full diagnostic formulation.

DSM expresses this through **three** criteria that must all be met. Their structure is clinically elegant. The intellectual criterion asks whether the relevant cognitive functions are impaired. The adaptive criterion asks whether that impairment matters in real life. The onset criterion prevents an acquired cognitive disorder from being placed in a developmental category. The diagnosis consequently rests on convergent evidence, not on one encounter or one instrument.

■ **RULE** **Diagnostic principle:** diagnose intellectual disability only when intellectual limitation, adaptive limitation and developmental onset are demonstrated together.

This formulation is especially important where a family requests a certificate, school accommodation, or diagnostic clarification after a long period of difficulty. The clinician should keep the diagnostic question separate from the administrative question of whether services are known to the family or available locally. Service contact may reveal need, but it does not create the disorder. Conversely, absence of prior contact does not negate a developmental disability.

3.2 DSM diagnostic criteria: the diagnostic formulation

Criterion A concerns deficits in intellectual functions. In practice, the clinician looks for difficulty in reasoning, problem solving, planning, abstract thinking, judgment, academic learning, or learning from experience, and confirms the finding through both clinical assessment and individually administered standardised intelligence testing. The word “both” matters. A formal result gives disciplined comparison with an appropriate normative group, whereas the clinical assessment asks whether the result is coherent with the history, observed performance, language, education, sensory status and opportunities to learn.

Criterion B concerns **adaptive functioning**. The relevant question is whether deficits lead to failure to meet developmental and sociocultural expectations for personal independence and social responsibility. Without continuing support, the deficits must limit activity in everyday life across multiple settings. This moves assessment from an abstract estimate of ability to the person’s actual functioning at home, in education or work, and in the community. The criterion can be met where there is sufficient impairment in at least one adaptive domain; the domains are conceptual, social and practical.

Criterion C requires onset of both intellectual and adaptive deficits during the **developmental period**. This is a historical criterion, so it requires collateral. Developmental records, school reports, informant accounts and prior assessments may be more informative than a patient’s retrospective account alone. The question is not whether the person was ever labelled, but whether the characteristic pattern was present in development.

- Ask whether the cognitive difficulty is broad enough to affect reasoning and learning rather than reflecting a circumscribed academic problem.

- Ask what the person can do independently, what they can do only with prompting, and what remains unsafe or unachievable despite ordinary opportunities.
- Seek a developmental history from someone who knows the person's earlier functioning, while recognising that documentation may be incomplete.

The criteria are not a checklist to complete mechanically. A person may perform poorly because of an acute mental state, language mismatch, sensory limitation, poor engagement or an unsuitable test. Equally, a person may show apparently acceptable performance in a familiar, highly supported routine while being unable to generalise skills or assume responsibility in less structured settings. Clinical synthesis is what decides whether the three criteria describe one developmental condition.

3.3 Interpreting the intellectual-functioning criterion

Criterion A is met when intellectual functioning falls **two or more standard deviations below the population mean**. That is a diagnostic threshold, not a severity grade, and it must not be turned into one: this chapter grades severity by adaptive functioning, and the reasons are set out above. What an IQ score actually is, how far it can be trusted, and what makes an obtained score misleading are the business of the clinical-assessment section, which takes the normative distribution, the error margin around a score, the **Flynn effect**, practice effects and the instruments that cannot carry a diagnostic opinion in turn. Read it before you act on any number.

■ **TRAP Examination trap:** an IQ score is evidence for Criterion A, not a stand-alone diagnosis and not a substitute for adaptive assessment.

One consequence belongs here rather than there, because it is a statement about the criterion itself and not about the instrument. Discordance between intellectual testing and everyday performance deserves active thought rather than reflexive exclusion: DSM recognises that a person whose score lies somewhat above the conventional Criterion A band may nonetheless have adaptive problems so substantial that their actual functioning resembles that of people with lower scores. The response is not to lower a threshold by intuition. It is to review test validity, collateral information, the quality of adaptive evidence, and the developmental course before reaching a defensible formulation. The criterion is a floor for evidence, not a verdict delivered by a number.

3.4 Adaptive functioning and onset: turning scores into a diagnosis

Adaptive functioning is the bridge between ability and lived consequence. It asks whether the person can meet the expectations of independence and social responsibility that are reasonable for their developmental stage and sociocultural context. It is assessed through functioning, not inferred from social class, literacy, diagnosis already recorded in a file, or the wishes of a caregiver. The clinician should distinguish lack of opportunity from inability, and a task performed under extensive supervision from a task performed independently and reliably.

GOOD TO REMEMBER

Adaptive functioning asks what the person can do in everyday life across settings without ongoing support. Assess the quality and consistency of functioning, the scaffolding supplied by relatives or institutions, and what happens when that scaffolding is withdrawn. Measurement of adaptive behaviour belongs alongside, rather than in place of, this clinical enquiry. The framework specifies three domains: conceptual, social and practical.

Developmental onset has a second purpose beyond classification. It forces a chronological formulation. If the observed cognitive and functional decline is new, fluctuating, or clearly follows a later medical or psychiatric event, the clinician must consider an acquired cause rather than assume an intellectual developmental disorder. If early difficulties were present but were never assessed, the absence of a historic score should not itself defeat the diagnosis. What matters is whether reliable developmental evidence can establish the required pattern.

MNEMONIC**Think “mind, mastery, maturation”**

Intellectual functions describe the mind; adaptive functioning demonstrates mastery of everyday demands; developmental onset establishes maturation as the time-course. The elements must point in the same direction.

In documentation, state the evidence for each criterion separately before writing the diagnosis. This prevents a report from using one broad phrase, such as “poor functioning,” to conceal which element has actually been demonstrated. It also makes the reasoning reviewable when informants disagree, records are sparse, or testing conditions were imperfect.

3.5 ICD diagnostic framework

ICD-11 groups the condition under **6A00 Disorders of intellectual development**. It describes the construct as a **metasyndrome** occurring in the developmental period: a clinically useful reminder that the diagnosis identifies a pattern of developmental functional impairment, not a single aetiology. The aetiological formulation must therefore remain open and be pursued separately when indicated.

ICD specifies three essential features: significant limitation in intellectual functioning, significant limitation in adaptive behaviour, and onset during the developmental period. Its intellectual-functioning feature explicitly includes perceptual reasoning, working memory, processing speed and verbal comprehension. These named areas should not be substituted for the DSM Criterion A list, which is framed differently. The two systems overlap clinically but phrase their intellectual evidence in distinct ways.

DIAGNOSTIC QUESTION	DSM FORMULATION	ICD FORMULATION
Intellectual limitation	Deficits in intellectual functions, supported by clinical assessment and individual standardised testing.	Significant limitation in intellectual functioning, including the specified cognitive areas.
Functional consequence	Adaptive deficits that compromise expected independence and social responsibility.	Significant limitation in adaptive behaviour.
Time-course	Intellectual and adaptive deficits begin during development.	Both limitations have developmental onset.
Use of IQ	Interpret in clinical context; never use as the sole basis.	Treat as a proxy for intellectual limitation, alongside comprehensive adaptive evaluation.

ICD indicates significant limitation in intellectual functioning at approximately **two or more standard deviations below the mean, or below the 2.3rd percentile**. It uses the same statistical convention for a total adaptive-behaviour score. These thresholds describe the boundary with expected functioning; they do not remove the requirement for comprehensive clinical interpretation.

3.6 Assessment at the boundary: reliability, culture and unavailable tests

Both systems reject the seductive but unsafe idea that IQ alone separates disorder from normality. ICD states that IQ is a proxy measure rather than an isolated diagnostic requirement at this boundary, and diagnosis must include a comprehensive evaluation of adaptive behaviour. This protects against the false certainty produced when a numerical score is detached from language, culture, education, health, developmental opportunity and everyday functioning.

Scores may vary substantially across development. ICD consequently advises testing on repeated occasions across the developmental trajectory when a reliable estimate is required. A child can appear to meet diagnostic requirements at one assessment and not another; the clinician should interpret change before declaring either recovery or error. **Reassessment** is particularly valuable when the initial assessment was constrained by unfamiliarity, illness, poor engagement, communication difficulty, or an unstable environment.

Where appropriately normed standardised tests are unavailable, ICD places greater weight on **clinical judgement**, including behavioural indicators supplied in its diagnostic guidance. This is highly relevant to heterogeneous linguistic and educational contexts. Clinical judgement here means explicit, accountable reasoning from history, observation and functional evidence. It does not mean replacing assessment with a casual impression, assuming disability from poverty, or assigning a label merely because a formal test cannot be obtained.

- Record why a particular test result is credible, limited, or unavailable.
- State whose account supports developmental onset and what setting each adaptive observation represents.

- Describe the language and sociocultural context relevant to interpretation, rather than treating them as background decoration.

When standardised testing is difficult or impossible, such as in the presence of sensory impairment or severe problem behaviour, DSM permits the diagnosis of **unspecified intellectual developmental disorder**. This is a category for genuine diagnostic limitation, not a shortcut around careful clinical work. The report should make clear what cannot currently be established, what evidence is present, and what reassessment could clarify.

3.7 Terminology, prevalence and the distinction from service counts

Terminology has changed because diagnostic language affects how people are understood in families, services and law. DSM uses intellectual developmental disorder to align with ICD-11, while retaining intellectual disability because it remains widely used by professionals, advocacy groups and the public. In a clinical record, use respectful current terminology and describe the person’s functioning and support context rather than allowing a historical label to stand in for a formulation.

Prevalence needs a denominator before it can be interpreted. DSM estimates overall general-population prevalence at approximately **10 per 1,000**, but figures derived from registers, schools or service rosters answer a different question from population studies. The concept of **administrative prevalence** concerns the people for whom a community would need services, and in practice often those whose needs are known to providers. It is useful for planning, but it should not be presented as an intrinsic biological rate.

10 per 1,000 DSM OVERALL ESTIMATE	16 per 1,000 DSM MIDDLE-INCOME ESTIMATE	9 per 1,000 DSM HIGH-INCOME ESTIMATE
---	---	--

IN INDIA

Reported Indian point prevalence lies between 1/1,000 and 32/1,000. A cited meta-analytic estimate is 10.37 per 1,000. In high-income settings, reported figures are 9.2 per 1,000 across all ages, 18.3 per 1,000 in child and young-person populations, and 4.9 per 1,000 in adult-only populations. The breadth of such estimates, including the reported Indian range, should make the reader ask how cases were defined, what population was sampled, and whether ascertainment relied on services, schools or household assessment. This is not pedantry. A prevalence figure used without its case definition and denominator can distort planning, obscure unmet need, or create a misleading comparison between settings.

Must remember: a low IQ score can support the diagnosis, but intellectual disability requires a developmental history and demonstrable adaptive limitation as well.

3.8 A defensible diagnostic conclusion

A high-quality conclusion names the diagnostic framework used, then shows how the evidence meets each required element. It identifies the source and limitations of intellectual assessment, gives a function-based account of adaptive impairment in relevant settings, and states the

developmental evidence. It should also acknowledge uncertainty when the evidence is incomplete. This is more clinically useful than a conclusion that merely reproduces a test result.

Do not use the diagnostic threshold to assign a severity category. The grading of intellectual disability is based on adaptive functioning and is addressed in the severity classification section. Do not confuse the diagnosis with borderline intellectual functioning, which is considered in the differentiation section. If the broader assessment has not yet been completed, say what is provisional and what collateral, functional assessment or repeat testing is still needed. The full assessment framework is covered in the Child psychiatry: assessment, treatment, services and protection notes.

4 · Severity classification (mild, moderate, severe, profound) by IQ and adaptive functioning

Severity is a description of lived support need. In intellectual disability, a severity label should help the clinician communicate what is difficult in everyday life, anticipate the breadth of assistance required, and organise a realistic plan with the person and family. It is not a shorthand for worth, potential, behaviour, or prognosis. The same label can conceal very different strengths, vulnerabilities, opportunities and barriers if it is treated as an identity rather than a clinical formulation.




Fig 21.3 Contemporary severity grading in intellectual disability is based on adaptive functioning across conceptual, social and practical domains. IQ contributes evidence but does not by itself determine the level of support a person needs.

The familiar grades run across **four** levels: mild, moderate, severe and profound. Their enduring examination relevance is that older systems linked them closely to IQ bands, whereas contemporary diagnostic practice grades severity through functioning in ordinary settings. Candidates need to be able to state both schemes, distinguish their purposes, and explain why the second is clinically more useful.

4 RECOGNISED SEVERITY GRADES	Adaptive BASIS FOR CONTEMPORARY GRADING	IQ SUPPORTING CLINICAL EVIDENCE	Supports PRACTICAL CONSEQUENCE OF THE GRADE
--	---	---	---

4.1 What a severity classification is for

A **severity classification** is a communication tool, not a substitute for assessment. It gives a shared language for describing the extent of impairment, but the clinical account must still say what the person can do independently, what they can do with prompting or teaching, and where

risk or dependence becomes important. A label that is not linked to concrete functional information has little value to the family, school, employer or care team.

Severity must therefore be used at the level of the individual. It should be informed by functioning across conceptual, social and practical domains. Conceptual functioning includes the use of knowledge in everyday demands. Social functioning concerns interpersonal judgement, communication in relationships and the ability to understand social situations. Practical functioning concerns self-care, routine, safety, work-related tasks and participation in community life. These domains may be uneven. A person may show a relative strength in one setting and substantial dependence in another, particularly when a familiar and supportive environment masks difficulty that becomes apparent with changing demands.

The clinician's task is to translate this profile into a defensible grade without reducing the profile to the grade. Good documentation makes clear which examples support the judgement: what was directly observed, what reliable informants describe, and what the person manages when environmental scaffolding is withdrawn or new demands arise. It also separates lack of opportunity, sensory or motor limitations, language mismatch, physical illness and psychiatric symptoms from limitations attributable to intellectual disability. Otherwise severity can be overstated by deprivation or understated by unusually protective care.

■ **RULE** **Grade severity by the person’s adaptive functioning, then use the IQ result as supporting clinical evidence.**

GOOD TO REMEMBER

This rule protects against a common distortion. A psychometric result can be highly informative, but it does not by itself show how the person manages money, relationships, self-care, safety, travel, routine, communication or unexpected change. Conversely, an apparently competent routine may depend on extensive family organisation and may not generalise beyond a familiar context. Severity is therefore a clinical judgement grounded in the person’s actual life, not merely an arithmetic location on a test distribution.

4.2 **The contemporary adaptive-functioning approach**

Adaptive functioning is the organising principle of contemporary severity grading. The DSM-5-TR severity table describes each grade through conceptual, social and practical functioning rather than through an IQ column. This makes the table clinically legible: severity is not simply a description of test performance but a description of the gap between the person's abilities and the demands of their usual environment.

The reason for this shift is pragmatic as well as conceptual. Support needs are determined by what the person can and cannot accomplish safely and consistently in daily life. An IQ score contributes evidence about intellectual functioning, but it does not directly specify supervision, teaching, environmental adaptation, decision support or the degree to which another person must take responsibility for safety. In this approach, the grade predicts the nature of help that has to be planned, not a fixed ceiling on learning.

Adaptive information should be interpreted in context. The same skill may be performed independently in a familiar home and not be available in a more demanding environment. Families may compensate so effectively that important dependence is missed, while social exclusion or inadequate education may produce apparent deficits that are not a stable reflection of underlying capacity. The assessor should therefore ask about performance across settings, response to ordinary change, consistency over time and the amount of cueing, monitoring or physical assistance required. This is why collateral history is central rather than decorative.

The grading rule also prevents false precision. Two people with similar test results may differ materially in social judgement, practical safety, communication, co-occurring illness, sensory impairment, motor disability or access to support. Conversely, a person whose score is difficult to interpret because of language, behaviour or physical limitations may still have a clear adaptive profile. The clinical formulation should acknowledge uncertainty where it exists, rather than force a spurious numerical conclusion.

■ **TRAP Examination trap:** do not write that a contemporary severity grade is “based on IQ.” State that severity is based on adaptive functioning, while IQ informs the wider assessment.

For DSM severity terminology, the crucial distinction is that the grade is decided from adaptive deficits. This is a useful sentence to carry into a long answer because it explains why the classification has clinical purpose. It also guards against a mechanical answer in which a candidate recites IQ ranges without demonstrating that they understand the modern framework.

4.3 DSM severity labels and coding language

The diagnostic labels are familiar, but they should be treated as severity specifiers attached to an individualised functional account. The codes below are useful for records and examinations. They do not replace a narrative description of the person’s functioning, current supports, vulnerabilities or strengths.

The linked coding language is F70 for mild, F71 for moderate, F72 for severe, and F73 for profound. Pair the code and label with the adaptive profile and the help required in real settings. The code identifies a category; the formulation tells another clinician how to care for the person.

In clinical writing, the label should lead to action. A bare diagnosis may be sufficient for a register, but it is insufficient for care planning. The formulation should say which tasks need teaching, which require environmental adaptation, which need supervision, and which are currently managed with assistance. It should also identify how communication, choice and autonomy will be protected. Severity is most useful when it improves coordination between psychiatry, family, education, rehabilitation and community services.

Do not infer a person's adaptive profile from the label alone. Higher-grade disability can coexist with preferences, learning and meaningful participation, and lower-grade disability can be accompanied by marked risks in social judgement, exploitation or practical safety. The label alerts the team to likely breadth of need; the assessment tells the team what that need actually is.

4.4 IQ-based severity bands: their place and their limits

IQ-based severity bands belong to the older classification logic and remain examinable because they appear in legacy records, teaching material and administrative language. They are best presented as a historical mapping, then immediately qualified: a score range is not a contemporary severity judgement. The IQ construct, its interpretation and the choice of intelligence testing are addressed in the clinical assessment section; here the concern is only the mapping of an IQ result to a grade.

LEGACY GRADE	IQ BAND IN ONE REPORTED ICD-10 SCHEME	CLINICAL READING
Mild	50-69	Use as a historical IQ band, not as the sole current severity determinant.
Moderate	35-49	Interpret alongside the person's actual adaptive functioning.
Severe	20-34	Do not assume practical abilities from the band alone.
Profound	below 20	Functional observation remains indispensable, especially when testing has limits.

The legacy statement that the IQ cut-off is 70 is often quoted alongside the claim that ICD-10 categorised the grades by IQ. That statement accurately captures the older scheme, but should not be carried forward uncritically into contemporary diagnostic phrasing. The meaningful clinical question is not only where a score lies, but whether the individual can meet culture-appropriate everyday demands without assistance.

A source discrepancy deserves explicit caution. Another textbook presentation places the upper end of the mild band at 50-70, rather than at the upper boundary shown in the preceding table. This is not a reason to blend the two into an invented compromise. In an answer, name the classification framework being used, give its band accurately, and then state that contemporary severity is determined from adaptive functioning. That is safer than presenting a disputed edge value as universal fact.

The deeper lesson is that score bands create an appearance of sharp boundaries where clinical reality is continuous and context-dependent. Borderline intellectual functioning is a separate boundary concept and should not be folded casually into intellectual disability; its definition belongs in the differential-diagnosis discussion. Likewise, a test result affected by language, behaviour, fatigue, sensory limitations or unfamiliarity with the testing context should prompt careful interpretation, not an automatic severity label.

4.5 ICD-11: severity across ability and adaptive behaviour

The **ICD-11 approach** assigns severity according to the level in which the majority of the person's intellectual ability and adaptive-behaviour skills fall across conceptual, social and practical domains. This is an important refinement of single-score thinking. It accepts that a

profile can be uneven, yet asks the clinician to identify the level that best represents the predominant pattern rather than choosing the most impaired isolated skill or the most successful isolated activity.

The numerical bands in ICD-11 are expressed as distance from the mean and corresponding percentile information. They should be read as psychometric context, not as a replacement for the adaptive judgement. The accompanying table is deliberately dense because the codes and bands are examination material; in practice, the functional formulation remains the more clinically useful document.

ICD-11 GRADE	PSYCHOMETRIC BAND	CODE	KEY IMPLICATION
Mild	2-3 SD below the mean; approximately 0.1-2.3 percentile	6A00.0	Assign the grade with the adaptive profile in view.
Moderate	3-4 SD below the mean; approximately 0.003-0.1 percentile	6A00.1	Do not let the numerical band eclipse functional variation.
Severe	4 or more SD below the mean; less than approximately the 0.003rd percentile	6A00.2	Distinguish this grade through adaptive behaviour.
Profound	4 or more SD below the mean; approximately less than the 0.003rd percentile	6A00.3	Distinguish this grade through adaptive behaviour.

The apparent overlap between the highest-grade psychometric bands is deliberate and clinically instructive. Severe and profound are separated exclusively by adaptive-behaviour differences, because available standardised intelligence tests cannot reliably and validly distinguish people at the extreme lower end of intellectual functioning. This is not a defect to be repaired by making the score seem more exact. It is a recognition that the most meaningful discrimination at this level concerns what the person is able to do and the support required for comfort, safety, communication and participation.

ICD-11 also provides 6A00.4 for a provisional grade and 6A00.Z for an unspecified grade. These should be used thoughtfully when a definitive grade cannot yet be assigned, rather than guessing from incomplete information. A provisional label preserves diagnostic honesty while assessment continues. An unspecified label should prompt the clinician to record what information is missing and what will be needed to clarify the person’s profile.

For longitudinal records that need a crosswalk to legacy coding, the mappings are 6A00.0 to F70, 6A00.1 to F71, 6A00.2 to F72, 6A00.3 to F73, 6A00.4 to F78, and 6A00.Z to F79. A crosswalk translates administrative language; it does not change the adaptive-functioning basis on which the current clinical grade is reached.

4.6 Reconciling discordant evidence in Indian practice

Apparent disagreement between intellectual testing and social-maturity findings is not unusual. It should not be treated as a nuisance to be averaged away. First check whether either result is invalid or unrepresentative because of communication barriers, behaviour, physical limitations, fatigue, unfamiliarity or an unsuitable testing context. Then return to the central clinical question: how does the person meet everyday demands in their own cultural and environmental setting?

IN INDIA

Where IQ and social-maturity results indicate different grades, the Indian guideline directs the clinician to decide in favour of the **social-maturity result**. The rationale is clinically coherent: social maturity reflects the degree to which the person meets culture-appropriate demands of daily life and generally better represents severity in ordinary circumstances. This is not an invitation to dismiss intellectual assessment. It is a reminder that the outcome of classification should describe functioning, not reward a deceptively precise test number.

The assessment method, the score it yields and its limitations belong in the adaptive-behaviour assessment section. Statutory disability quantification is a separate legal process and must not be confused with clinical severity or with adaptive functioning. In a clinical report, make the distinction explicit: describe the diagnostic severity formulation, describe the evidence used to reach it, and deal with certification through the relevant disability-certification pathway.

- Check that the intellectual and functional information reflects the person's usual, not merely best or worst, performance.
- Explain discordance rather than concealing it behind a single label.
- Prefer the evidence that best captures everyday functioning when a severity judgement is required.
- Record strengths, communication preferences and environmental supports alongside limitations.

4.7 From severity label to support profile

A grade becomes clinically useful only when it changes the plan. It should lead the team to ask what support is needed to enable learning, choice, health care, family participation, education, work, relationships and safety. The answer should be revisited as circumstances change. Severity is relatively stable as a description of pervasive need, but support can be improved through accessible communication, structured teaching, environmental adaptation and respect for the person's established competencies.

The **AAIDD** classification links the grades to a **profile of supports** instead, and is multidimensional rather than categorical. Its four intensities of support, in ascending order, are **intermittent, limited, extensive and pervasive**. Two qualifications keep that list honest. It belongs to the ninth edition of the AAIDD system and was not carried into the later editions, which found it lacked the standardised data to support graded intensities, so it is a historical

scheme that remains examinable rather than a current classification. And it is a description of what a person needs, not a fifth way of grading them: the four intensities are often lined up against mild, moderate, severe and profound, but the mapping is a convenience and the whole point of a supports model is that two people at the same severity grade may need very different support. Reproduce the four-word list as written: one widely used textbook prints extreme supports at the severe tier, but the term in general use, and the one an examiner expects, is extensive. These words should not be used as a rigid prescription for any one person. They are a reminder to specify intensity, consistency and context of help rather than to leave the plan at the level of a diagnostic adjective.

A support profile is also more humane than deterministic language. It turns the clinical focus from what the person lacks to the conditions under which the person can participate. It permits planning around autonomy as well as risk: who should provide information in an accessible form, how decisions can be supported, what routines promote competence, what supervision is proportionate, and how family burden can be recognised without erasing the person's voice.

MNEMONIC

GRADE

Ground the decision in daily functioning; Reconcile discrepant evidence; Avoid IQ-only classification; Describe supports; Explain the functional basis in the record.

The defensible severity statement is the one that explains the person's adaptive functioning and consequent support needs, not merely the one that reports an IQ range.

4.8 A disciplined answer and documentation approach

For a short-answer question, begin with the names of the grades, state the modern principle that adaptive functioning determines severity, and then contrast this with the legacy IQ-linked scheme. If asked for a classification table, present the requested framework cleanly. If the question is clinical, move beyond the table: identify the adaptive domains affected, state the basis for the grade, and outline the supports that follow. This sequence shows the examiner that the candidate understands both taxonomy and clinical reasoning.

For a case record, avoid language that makes the severity label sound final in isolation. State the diagnosis, the current grade, the evidence from functioning, the degree and type of assistance required, relevant contextual modifiers, and the plan to address identified needs. Be cautious when information comes only from one informant or one setting. A carefully qualified assessment is stronger than a confident but unsupported label.

The same discipline protects against two opposite errors. One is overmedicalisation, in which a score becomes the whole person. The other is under-recognition, in which a person's dependence is minimised because they have learned familiar routines or because relatives quietly perform the demanding parts of daily life. Both errors lead to mismatched care. Severity classification is

valuable precisely because, done well, it keeps clinical language anchored to the reality of support.

5 · Clinical assessment and IQ testing in ID

Assessment is a clinical act, not a score-collecting exercise. In suspected intellectual disability, the psychiatrist has to decide whether an observed difficulty represents a broad limitation in intellectual functioning, a barrier to showing ability in the assessment setting, a circumscribed learning problem, or a mixture of these. The task is therefore to obtain a valid account of how the person reasons and learns, and then to set that account beside history, observation, collateral information and adaptive-behaviour assessment. The numerical result matters, but it is never self-interpreting.

The marks in this area are usually won by showing disciplined interpretation. A candidate should be able to explain what an IQ score represents, why it is reported as an interval rather than treated as an infallible point, how a poor-quality administration can mislead, and why a patterned profile may be more clinically useful than an overall score. The assessment must also stay within its proper boundary: severity grading and the formal measurement of adaptive behaviour are separate tasks, addressed elsewhere in this chapter.




Fig 21.4 Choosing an ability instrument by age and by language. Two branch points, not a catalogue: whether an IQ test applies at all, and whether a verbal test can be given in the language the child actually speaks. Each of the three routes is named with its instruments and their sourced age bands, and each is drawn with the limit its own source states, because every escape from the language problem has a floor and the floor sits where intellectual disability lives.

5.1 The clinical question before testing

Begin with the referral question. “What is the IQ?” is usually an impoverished question. The useful question is: what decision will a careful estimate of intellectual functioning help the team make? The answer may concern the nature of a broad learning difficulty, the credibility of an apparent decline, the meaning of an uneven school record, the person’s capacity to engage with an explanation, or the supports that should be considered. Stating the question first protects the assessment from becoming an administrative ritual.

Intelligence can be understood clinically as a broad capacity to act with purpose, reason coherently and deal effectively with the environment. An intelligence test samples that capacity under defined conditions; it does not sample every ability a person possesses, every opportunity they have had, or every demand made by ordinary life. The psychiatrist should therefore ask what the test performance means in this person's lived context rather than asking the score to speak beyond its remit.

The first interview establishes whether testing is likely to answer the question. Clarify the presenting concern in the informant's own language, the course of learning and functioning, the educational and occupational setting, sensory or motor barriers, communication style, behaviour during unfamiliar tasks, and the person's current mental and physical state. Review available records for consistency rather than assuming that a label in a previous document is a conclusion. A recent deterioration in performance, acute distress, reduced alertness or inability to engage may change the meaning of a low obtained score. The full developmental history and child psychiatric assessment framework belong in the *Child psychiatry: assessment, treatment, services and protection notes*; here, their relevance is that they determine whether a test session is interpretable.

■ **RULE** Use psychometry to refine clinical judgment, not to replace it. A score becomes useful only when the referral question, the testing conditions and the person's real-world functioning have been considered together.

- Formulate the decision that testing is meant to inform before selecting the assessment.
- Establish whether communication, sensory, motor, emotional or situational factors could distort performance.
- Obtain collateral evidence sufficient to judge whether the test result fits the longitudinal clinical picture.
- Keep the estimation of intellectual functioning distinct from adaptive-behaviour measurement and from severity grading.

5.2 What an IQ score is, and what it is not

An IQ is a standardised comparison of an individual's performance with that of the reference group used by the test. Its clinical value lies in making the comparison explicit and reproducible. It is not a direct measure of worth, potential, character, family effort or entitlement to care. Nor does an overall score reveal the route by which a person reached it. People with a similar overall result may have quite different cognitive strengths, educational experiences, communication demands and support needs.

Older teaching may describe **ratio IQ** using the expression $\text{IQ} = (\text{mental age} / \text{chronological age}) \times 100$. This formulation is historically useful because it explains the term "IQ", but it should not be casually substituted for the standard-score framework used in contemporary interpretation. A clinician should state which metric the report provides rather than sliding between different meanings of the same abbreviation.

For the test metric assumed when deriving the diagnostic convention, the mean is 100 and the standard deviation is 15. The phrase “standard deviation” expresses the spread of scores around the mean in the reference population. It allows a clinician to describe the distance of a result from the population average without converting that distance into a moral or deterministic statement about the person.

100 ASSUMED TEST MEAN	15 ASSUMED STANDARD DEVIATION	≥2 SD below population mean CLINICAL CONVENTION	65-75 SCORE INTERVAL REQUIRING INTERPRETATION
---------------------------------	--	---	---

The intellectual-functioning convention lies **approximately 2 standard deviations or more below the population mean**. This is a convention for interpreting a result in relation to a population distribution, not permission to diagnose from arithmetic alone. The clinician still asks whether the obtained result is valid, whether the reference group is appropriate, and whether the person’s presentation is consistent with the assessment as a whole.

A report should name the metric before interpreting it. This is especially important because a composite score and a subtest score are not interchangeable. The overall number is a summary; it should not erase clinically meaningful variation within the assessment. Equally, a striking isolated weakness is not automatically evidence of a broad intellectual impairment. Interpretation requires pattern recognition, corroboration and restraint.

5.3 Measurement error and the discipline of reporting a range

An obtained score is an estimate. Every psychometric result contains uncertainty arising from the instrument, the testing occasion and the person’s performance on that occasion. The relevant question is therefore not “what is the person’s true IQ?” but “what range of intellectual performance is reasonably supported by this assessment, given its quality and context?” This language is more accurate and prevents a falsely precise clinical conclusion.

The generally applicable margin around an obtained full-scale score is **plus or minus 5 IQ points**. Thus, once measurement error is considered, the score interval associated with the intellectual-functioning criterion is **65-75**. The range must be read as an interval requiring clinical interpretation, not as a new single cut-off and not as a device for negotiating a preferred label.

GOOD TO REMEMBER

A result near the relevant convention calls for particularly careful attention to the quality of administration, the person’s engagement, collateral evidence and adaptive-behaviour assessment. An isolated point score does not settle a complex clinical question: explain how the assessment was conducted and what may have affected it.

- **TRAP Examination trap:** treating an IQ result as exact. A technically sound answer states the obtained score, acknowledges measurement error, interprets the interval, and then integrates the wider assessment.

INTERPRETIVE QUESTION	WHAT THE ASSESSMENT CAN CONTRIBUTE	WHAT STILL REQUIRES CLINICAL JUDGMENT
Is the score low?	It locates performance relative to relevant norms.	Whether the administration and norms make that comparison valid.
Is the result stable enough to use?	It provides an estimate with a stated margin of uncertainty.	Whether state factors or testing conditions make the estimate misleading.
Is there a broad difficulty?	It summarises performance across sampled tasks.	Whether the pattern fits history and everyday functioning.
Does a single total tell the whole story?	It offers a convenient summary score.	Whether uneven abilities make the profile more informative than the summary.
What support is needed?	It may identify cognitive demands likely to be difficult.	Adaptive-behaviour assessment and the real-world support context.

5.4 Selecting and conducting a valid assessment

Choice of assessment follows the person, not the convenience of the clinic. Select an individually administered, appropriately standardised measure that can address the referral question and can be administered meaningfully to the person in front of you. Language, communication access, sensory and motor capacity, familiarity with formal testing, educational opportunity and current state all affect the evidentiary weight that can be placed on an obtained result.

Norming for sociocultural background and native language is a requirement, not a courtesy. A score derived from unsuitable norms risks measuring distance from the test’s expected experience rather than the intellectual ability the clinician intends to estimate. This is particularly important when the assessment language is not the person’s strongest language or when the testing situation makes culturally specific knowledge carry disproportionate weight. The report should make any limitation transparent rather than concealing it behind a number.

Standardisation means that the conditions of administration, scoring and comparison are controlled as far as possible. In clinical practice, that does not mean rigidly ignoring a person’s needs. It means recording adaptations, interruptions and barriers, then judging whether they preserve or weaken the interpretability of the score. Rapport, a quiet setting, comprehensible instructions and adequate opportunity to respond are not cosmetic features. They are part of the validity of the encounter.

Observe the process as carefully as the final result. Does the person understand the task? Do they persist, become overwhelmed, guess rapidly, seek repeated reassurance, lose the instruction, or show a discrepancy between verbal explanation and non-verbal problem solving? Such

observations can clarify a profile, expose a threat to validity or indicate the need to defer a conclusion. They should be documented in clinically intelligible language, not reduced to a vague statement that the person was “cooperative”.

Assessment should be paced to protect effort without coaching the answer. The examiner may clarify the task only within standardised rules, attend to physical comfort, and recognise when fatigue or distress has made further performance unrepresentative. If a session is not interpretable, the responsible clinical conclusion may be that the assessment has not answered the question yet. A forced total score is not more scientific than an honest limitation.

- Explain the purpose of the assessment in language the person and family can understand.
- Record the assessment language, relevant accommodations and any barriers to participation.
- Describe behavioural observations that bear on validity, rather than using generic reassurance.
- State clearly when a result should be treated as provisional or when re-assessment is needed.

5.5 Composite scores, subtests and cognitive profiles

Do not confuse the ruler with its markings. A composite or index score uses a different metric from an individual subtest scaled score. **Wechsler composite and index scores: mean 100, SD 15; Wechsler subtest scaled scores: mean 10, SD 3**. Mixing these metrics is a basic interpretive error: a subtest value should not be read as though it were an IQ, and a composite should not be treated as though it describes a single task.

A **cognitive profile** is the pattern of relative strengths and weaknesses across the assessment, interpreted cautiously in context. It directs attention to how the person approaches demands rather than merely where the overall score falls. Profiles are most useful when they explain an observed clinical discrepancy, guide further assessment or suggest a practical way to communicate and support learning. They are least useful when they are turned into speculative labels unsupported by history or observation.

The preference for individual cognitive profiles over a single IQ is reflected in **cross-battery intellectual assessment**, which is more useful for understanding intellectual abilities than a single overall score alone. This does not mean that every patient needs an elaborate battery. It means that the breadth of assessment should match the clinical ambiguity. Where the total score and the observed person tell the same coherent story, a focused assessment may suffice. Where they diverge, the divergence is data, not an inconvenience to be averaged away.

Large differences across tasks require interpretation before the full-scale result is treated as a stable global estimate. Highly discrepant individual subtest scores may make an overall IQ score invalid. The key word is “may”: discrepancy prompts clinical scrutiny rather than an automatic rejection of the entire assessment. Consider whether the pattern is understandable from language demands, attention, motivation, sensory or motor factors, emotional state, educational experience or a specific cognitive weakness. The conclusion should describe what is known and what remains uncertain.

This distinction prevents a common diagnostic shortcut. Specific learning disorder may show a verbal-performance discrepancy, usually with performance IQ higher, and an ACID pattern of relative weakness in Arithmetic, Coding, Information and Digit-span. It may also show weaknesses in verbal comprehension, working memory and processing speed. These findings concern an uneven learning profile; they must not be converted mechanically into a severity statement about intellectual disability. The broader formulation belongs with the educational and communication assessment described in the *ADHD and specific learning/communication disorders notes*.

5.6 Threats to validity and misleadingly high or low results

Flynn effect refers to the risk of spuriously high IQ scores when test norms are out of date. It is a norms problem, not a personal change in ability. **Practice effects** are different: repeated testing can yield spuriously high scores because the person has learned from prior exposure. The distinction matters because the remedy is different. In one situation the concern is the currency of the reference norms; in the other it is the interpretive impact of repeated familiarity with the material.

Before accepting an unexpectedly high or low score, ask whether the test was suitable and whether the score was generated in a valid way. A brief intelligence screening test or group test may produce an invalid IQ score. Such tools can be useful for triage or as one source of information, but they do not automatically carry the evidentiary weight of a comprehensive individual assessment. The report should identify the method used and resist upgrading a screen into a definitive estimate simply because it has yielded a number.

Other validity threats are often visible in the room. A person who cannot hear or understand the instruction, whose motor difficulty prevents the intended response, or whose distress and fatigue collapse effort may obtain a result that reflects the barrier more than the target ability. The same caution applies when language mismatch or unfamiliarity with the testing format changes what is being sampled. The clinical response is not to “correct” the score by intuition. It is to document the limitation, seek appropriate assessment conditions and integrate other evidence.

Validity is not an all-or-none property. Some results provide useful minimum estimates, some are broadly representative but qualified, and some should not be interpreted as an IQ estimate at all. This graded approach is more honest than either rejecting every imperfect assessment or accepting every printed total. It also makes the report usable to colleagues: they need to know how much confidence to place in the result and why.

An IQ score is clinically meaningful only when the test, norms, administration and interpretation are all appropriate for the individual.

5.7 Integrating and communicating the result

The final account should read like a clinical formulation, not a laboratory slip. Start with the reason for referral and identify the assessment method and conditions. State the obtained result as an estimate, include its uncertainty where relevant, describe important profile findings, and

explain the factors that strengthen or limit confidence. Then integrate this with the history, observation, collateral information and adaptive-behaviour assessment. This sequence allows another clinician to see how the conclusion was reached and where its boundaries lie.

Families commonly hear an IQ result as a verdict. The psychiatrist's language can either amplify that harm or restore accuracy. Explain that the assessment describes performance on a defined set of tasks compared with relevant norms. Discuss strengths as well as difficulties, avoid presenting a score as a ceiling on development, and translate findings into practical implications. Do not use technical uncertainty as evasiveness: it can be explained plainly as the ordinary fact that a single testing occasion estimates rather than perfectly captures ability.

When a low score appears inconsistent with everyday functioning, do not decide in advance which source is wrong. Re-examine the referral question, test validity, language and norming, profile variability, collateral quality and the possibility that a different condition is influencing performance. When everyday difficulties seem greater than the score would predict, the same discipline applies. Intelligence testing and adaptive-behaviour assessment answer related but non-identical questions; neither should be used as a shortcut for the other.

Finally, communicate the scope of the opinion. A psychometric assessment may inform diagnosis and care planning, but it does not replace medical evaluation, aetiological work-up or adaptive-behaviour measurement. The laboratory investigation of possible causes is addressed in the etiological work-up section of this chapter, and formal adaptive-behaviour instruments are addressed in the adaptive behaviour assessment section. Keeping these strands separate while integrating their conclusions is the hallmark of a defensible intellectual-disability assessment.

5.8 Choosing intelligence instruments in Indian practice

IN INDIA

Selection is a clinical discrimination: fit the instrument to age, language and schooling, not departmental familiarity. A verbal age scale may be technically well administered yet answer the wrong question when the child is being tested outside the language of its norms. Conversely, a performance measure is not a lesser substitute for a "real" intelligence test when language is the obstacle: it samples ability through a different route. Before choosing a battery, decide whether the task requires a broad verbal and performance profile, a brief performance estimate, or a nonverbal reasoning measure, and be explicit about what the chosen instrument actually reports.

-
- **RULE** Choose the instrument by construct, age and language before choosing it by familiarity. The score is interpretable only when the test's norms and task demands fit the child in front of you.
-

INSTRUMENT	AGE BAND AND FORM	WHAT IT YIELDS	WHEN IT IS THE BETTER FIT - AND ITS LIMIT
Binet-Kamat Test of Intelligence	3 years to adulthood; a predominantly verbal age scale.	Age-scale estimate, not a language-free performance measure.	Use only when the verbal demands and language are appropriate. Its verbal items are unavailable in vernacular languages other than Kannada and Marathi.
Stanford-Binet, Kulshreshta Indian adaptation	3 years to adulthood; predominantly verbal age scale.	Age-scale assessment.	It is more suitable for a Hindi-speaking population; its norms did not include people with low intelligence.
Malin's Intelligence Scale for Indian Children	6-16 years; Indian adaptation of the original Wechsler child scale.	Performance IQ, Verbal IQ and Full-scale IQ.	Useful when a Wechsler-style profile is needed, but some verbal scales depend on formal education.
Bhatia's Battery of Intelligence	standardised on Indian boys aged 11 to 16, though the Indian guideline prints the range as 11 years and above; indigenous performance battery.	Performance-based intelligence assessment.	Choose it when verbal load is the principal barrier. It measures IQ only above 70 , so it is unsuitable for establishing lower-range intellectual functioning.
Seguin Form Board	Reliable at 3-11 years; valid across age groups of people with ID.	Performance test and quick measure of general intelligence.	A valuable low-verbal-demand option. Above 11 years , it increasingly reflects visuo-motor speed rather than global intelligence.
Raven's Standard Progressive Matrices	11 years to adulthood; nonverbal matrices.	Does not yield an IQ score.	Assesses general ability through form comparison and analytical reasoning. It is not suitable for illiterates or people at the lower end of the ID spectrum.
Raven's Coloured Progressive Matrices	5-11 years.	Percentile ranks, not an IQ.	Use when a developmentally younger child needs a matrix task, while reporting the actual metric rather than relabelling it as IQ.

INSTRUMENT	AGE BAND AND FORM	WHAT IT YIELDS	WHEN IT IS THE BETTER FIT - AND ITS LIMIT
WISC-IV India	6 years to 16 years 11 months ; a full child battery.	A full-scale score with an index profile, reported in this edition as verbal comprehension, working memory and processing speed alongside perceptual reasoning.	Use where age and language permit a full child battery. Do not describe this edition's output as a verbal/performance split: that is the older Wechsler structure, and some Indian teaching sources still print it. Do not wrongly transfer Bhatia's stated above-70 floor to WISC-IV India.

The **Binet-Kamat** and the **Kulshreshta Stanford-Binet adaptation** are the classic age-scale choices, but both are predominantly verbal. That is their clinical cost. If the child cannot comprehend the assessment language with the same ease as the norm group, a low result may partly index linguistic distance, schooling or test familiarity rather than the intended construct. Binet-Kamat's language restriction makes this concern concrete. The Hindi orientation of the Kulshreshta adaptation is similarly a selection rule, not a minor reporting footnote. Neither should be used merely because it is available in the department.

MISIC is often the most useful familiar-scale option for a school-age child because it derives from the Wechsler child scale and separates performance, verbal and full-scale results. The separation matters clinically: an uneven verbal-performance pattern can signal that the overall total conceals how the child is managing the tasks. It does not remove the education dependence of some verbal scales. A child with limited formal schooling should not be made to look globally less able because a verbally loaded test has been treated as culture-free.

The performance instruments address this problem directly. **Bhatia's Battery** includes many indigenous subscales and is the performance route for an older child, within its norms. Two limits govern its use and both are examinable. Its stated floor prevents it from being the decisive measure when lower-range intellectual functioning is suspected. And although the Indian guideline prints its range as 11 years and above, the battery was standardised on Indian boys aged 11 to 16 in the 1950s, so reading an adult against it applies sixteen-year-old norms to a grown patient. That is an extrapolation beyond the instrument, not normative adult use, and the open-ended phrasing of the guideline is exactly what invites it. The **Seguin Form Board** is a quick performance measure, useful where verbal demands obstruct assessment, but its changing emphasis after the relevant age means that fast visuo-motor work must not be misread as a global estimate of intelligence.

■ **TRAP** “**Raven's gives an IQ**” is wrong. **Standard Progressive Matrices does not output an IQ**. Do not convert a percentile or a matrix result into an IQ merely because the test is familiar.

Raven’s Standard Progressive Matrices has Indian norms and is culture-fair to a large extent, but that does not make it universally suitable or make it an intelligence quotient. It evaluates form comparison and analytical reasoning, and its limits in illiteracy and more severe intellectual impairment are clinically important. **Coloured Progressive Matrices** supplies percentile ranks for younger children. State the result in the metric provided. Finally, **WISC-IV India** offers a full child battery across its specified age range, reporting a full-scale score with an index profile rather than the verbal/performance split of the older Wechsler editions. It may be the appropriate comprehensive choice only after language, schooling and tolerance of formal testing have been considered. Selection is a defensible match between construct and patient.

6 · Adaptive behaviour assessment (Vineland, DST, VSMS, Gesell)

The question is not merely what the person can do in a testing room. Adaptive-behaviour assessment asks how a person manages the ordinary demands of life in the setting in which they live. It therefore supplies information that a performance score cannot supply: whether an observed difficulty is translated into dependence in self-care, communication, social participation, safety, or practical daily routines. For intellectual disability, this is clinically decisive because support planning must follow real-world functioning, not the elegance of a single test result.

The instruments often grouped together in examination answers do not measure one thing. A developmental screen estimates developmental progress; a social-maturity scale yields a social quotient; an adaptive-behaviour scale documents functioning through an informant; and a drawing screen provides only a screening impression. The candidate who first identifies the construct, the respondent, and the output is much less likely to call a developmental quotient an IQ score or to use a social-maturity result as though it were a statutory disability percentage.

0-15 DST AGE SPAN, YEARS	0-15 VSMS AGE SPAN, YEARS	0-90 VINELAND-3 AGE SPAN, YEARS
------------------------------------	-------------------------------------	---




Fig 21.5 Four constructs, four instruments. Intelligence, development, social maturity and adaptive behaviour are four separate constructs with four separate instrument sets and four different output metrics, and no score in one column converts into a score in another. The plate also prints the two places the corpus contradicts itself, rather than resolving them: whether the VSMS is an intelligence test or the adaptive-behaviour instrument, and whether an IQ is a ratio of mental to chronological age or a deviation score with a mean of 100 and a standard deviation of 15. Choosing an ability instrument by age and language is the previous plate's job; this one is only about what each column measures and what it reports.

6.1 Begin with the construct, not the familiar name

Adaptive-behaviour measurement is a structured account of habitual functioning. It is not an abbreviated intelligence test. The best information comes from a respondent who has seen the person across ordinary situations and can distinguish what is done independently, what is done with prompting, and what is not yet part of the person's routine. The assessor then integrates this account with direct observation, history, opportunities for learning, physical or sensory

limitations, and the quality of the informant’s knowledge. A score may be useful, but the clinical formulation is stronger when it retains the profile behind the score.

This distinction matters especially when opportunity and capacity have been confused. A person may fail to perform a task because it has never been taught, because the setting does not permit it, because the task is inaccessible, or because of a primary developmental limitation. Conversely, an apparently competent answer from a family member may describe what the family does for the person rather than what the person can actually do. Adaptive assessment should therefore examine typical performance, not an idealised report and not a one-off best performance in clinic.

- **RULE** **Rule:** treat an adaptive score as an organised summary of everyday functioning, then return to the underlying profile whenever the score and the clinical picture do not fit.

The terminology itself provides a useful safeguard. **Developmental quotient** is the output of developmental testing and is interpreted in the same metric logic as an IQ score, but it describes developmental assessment rather than a conventional intelligence battery. **Social quotient** is the output of the Indian social-maturity scale. Neither label should be casually substituted for the other. A developmental result, a social-maturity result, and a formal ability result answer overlapping but non-identical clinical questions.

CLINICAL QUESTION	APPROPRIATE CONSTRUCT	ILLUSTRATIVE OUTPUT	WHAT IT MUST NOT BE CALLED
Has the child’s overall development pro-gressed as expected?	Developmental screening	DQ	An IQ result
How does the person manage socially and practically in their milieu?	Social maturity and adaptive profile	SQ with domain profile	A developmental quotient
What is the person’s functioning across adaptive domains by informant report?	Adaptive behaviour	Adaptive profile	A direct performance battery
Is a rapid developmen-tal screen needed be-fore fuller assessment?	Performance screening	Screening impression	A definitive diagnostic quotient

6.2 The Indian social-maturity scale in practice

The **Vineland Social Maturity Scale**, in its Indian adaptation, is the central practical instrument for this purpose. The adaptation is attributed to Malin and was later expanded by Bharatraj. It assesses culturally appropriate adaptive behavioural skills across **eight** important domains and yields an SQ. The domain profile is often more clinically useful than a compressed global statement because it shows where the person is independent, where cueing is enough, and where sustained assistance is needed.

IN INDIA

In the cited guideline, the VSMS has a unique place among standardised adaptive-behaviour measures in India. This gives it an established role in Indian assessment, without implying that every clinical population, language group, or contemporary setting is equally represented.

Document the sociocultural context of responses: functioning is enacted in a household, school, neighbourhood, and set of expectations.

GOOD TO REMEMBER

Administration is not a mechanical interview. Identify who knows the person's usual functioning best, and ask for concrete examples. Caregiver disagreement may reveal different opportunities, supervision, or settings, or an informant who has not observed the task. Test implausibly high or low responses with observation and collateral information rather than forcing premature consensus.

- Ask what the person does on an ordinary occasion, not what they have done once under exceptional help.
- Clarify whether initiation, sequencing, supervision, and completion are independent.
- Separate lack of opportunity from inability, while recording both as clinically relevant.
- Interpret a scattered profile as a starting point for formulation and support planning, not as a nuisance to be averaged away.

The VSMS is an adaptive-behaviour instrument: it yields a social quotient plus a profile of 8 domains of adaptive behaviour, never an IQ. Naming it, the DST or Gesell among intelligence tests is the single commonest error in this topic, and where a social quotient and an IQ disagree the severity decision is made on adaptive grounds, as the severity discussion sets out.

This is a measurement rule, not a licence to ignore contradictory evidence. Discordance should prompt a review of language demands, test engagement, motor or sensory barriers, educational opportunity, and the reliability of the informant account. It should also prompt a return to the person's actual support needs. Grading intellectual disability is dealt with separately; the point here is narrower but essential: everyday adaptation has priority when the two numerical summaries pull in different directions.

If the scale cannot be administered, the recommended fallback is to ask socioculturally relevant questions and use DSM-5 severity specifiers as an interim clinical reference, pending ICD-11 guidance. This fallback must be documented as a clinical assessment rather than presented as a scored scale. It is particularly important to make explicit who supplied the history, which settings were represented, and which ordinary activities could not be meaningfully assessed.

6.3 Vineland scales: distinguish the scale family from the Indian VSMS

The **Vineland Adaptive Behavior Scales** are distinct from the Indian VSMS despite their similar names. The scale family is associated with Sparrow, Balla and Cicchetti; later editions are separately attributed in the source. The earlier social-maturity tradition is associated with Edgar Doll. In a viva, naming “Vineland” without specifying which instrument invites avoidable ambiguity about the form, the respondent, the norms, and the result being discussed.

The Vineland scales assess adaptive capacities for self-sufficiency across domains that include **communication, daily living skills, socialization, and motor skills**. They can be administered as a survey interview, expanded interview, informant-administered rating form, or teacher rating form. This range is clinically useful because the person’s everyday capacities are often expressed differently at home and in an educational setting. It also creates an interpretive obligation: a discrepancy between parent and teacher reports should be explored as a context effect, not dismissed as an error.

Vineland-3 provides an age-normed framework for adaptive functioning and has self-report, parent-report, and teacher-rated youth forms. Its reported age span runs from **birth to age 90**. The strikingly broad span is a reminder that adaptive behaviour is not a childhood-only construct. Nevertheless, a score gains meaning only when the clinician considers the person’s communication style, culture, opportunities, environmental supports, and the informant’s frame of comparison.

Informant scales can compare a child with peers of the same age, but their parent-report basis lacks the objectivity of standardised child-based assessment. This is not a reason to discard informant data. It is a reason to use it intelligently. Direct testing samples what the person can show under examination conditions; an informant scale samples what is habitually enacted across time. These evidence streams should be reconciled, not forced into a false competition.

■ **TRAP Examination trap:** do not use VSMS and VABS as interchangeable labels. They are distinct instruments with different stated adaptations, formats, and assessment frameworks.

6.4 Developmental assessment: where the DST and DQ belong

The **Developmental Screening Test** was developed by Bharat Raj. It uses developmental tasks to assess global development across the stated childhood age range. The result is developmental screening, not an IQ test. This distinction is not semantic fussiness. A developmental screen is appropriate when the clinical question is whether development is globally delayed and whether fuller assessment is needed; it should not be made to carry a conclusion that its construct does not support.

Developmental scales are used when formal IQ tests are inapplicable, including very young age, sensory-motor issues, and gross understimulation. The Indian instruments named for this role are the Developmental Screening Test and the Developmental Assessment Scales for Indian Infants. Such selection is clinical reasoning, not a lesser substitute for formal testing. If a child cannot access the demands of an ability test, the clinician should choose a measure that captures the available developmental evidence and record why the conventional approach was not suitable.

The DST is highly loaded with speech and language items. It therefore requires cautious interpretation in cerebral palsy, autism, speech impairment, and hearing impairment. A low score in these circumstances may reflect the route through which the test asks for competence rather than the global developmental capacity one is trying to understand. The report should make the limitation visible and integrate nonverbal observation, history, and relevant specialist assessment. It should not quietly convert a language-loaded result into a global intellectual conclusion.

SITUATION	ASSESSMENT IMPLICATION	CLINICAL SAFEGUARD
Very young child, sensory-motor difficulty, or gross understimulation	Developmental scales can replace inaccessible IQ testing.	State why a conventional ability test could not answer the question.
Marked speech or language limitation	DST results need guarded interpretation.	Do not equate verbal loading with global developmental capacity.
Need for an adaptive account in an Indian setting	VSMS supplies the adaptive component.	Keep its SQ and profile separate from an IQ result.
Motor deficit affects an ability-test performance	Binet-Kamat profile analysis can separate a motor contribution.	Refer detailed IQ-instrument selection to the clinical assessment and IQ testing discussion.

6.5 Gesell: developmental model, drawing screen, and infancy

The **Gesellian model** starts with the newborn and tracks normal developmental trends across distinct but interrelated domains, rather than reducing infancy to a single factor of intellectual ability. This is a conceptual advance worth retaining in practice. Infancy is a period in which posture, movement, communication, social engagement, regulation, and emerging problem solving may develop unevenly. A developmental account should preserve that pattern rather than overstate the precision of an infant intelligence label.

The older infant-IQ model did not successfully extend conventional IQ assessment into infancy and has largely fallen out of use. The practical lesson is modest: report developmental findings with appropriate caution about what they predict, and explain the functional significance of the observed pattern. Prediction from infant intelligence testing is limited, whereas preschool testing is generally more informative. This is why longitudinal clinical review and repeated observation often matter more than attaching undue certainty to an early numerical label.

The **Gesell’s Drawing Test** is an Indian screening instrument adapted by Verma and colleagues and later revalidated by Venkatesan. It is a performance screen for mental development in the stated age span. It obtains a rough estimate of intelligence from the child’s ability to draw age-expected shapes. That word “rough” is the safeguard: the task can orient an assessment, but it cannot carry the authority of a comprehensive diagnostic conclusion.

Drawing performance is inseparable from opportunity and motor access. The test is not valid for children with no school attendance, no experience with a pencil, or specific finger-dexterity problems. A poor drawing response in such circumstances should be treated as a limitation of the

method, not as evidence of global inability. The same discipline applies to all brief screens: a screen should direct the next clinical question, never close it.

In the broader infant-assessment lineage, Nancy Bayley drew heavily on Gesell's developmental tasks and techniques. The Bayley-III assesses **five** domains: cognitive, language, motor, adaptive behaviour, and social-emotional functioning. It is reported for children from **1 to 42 months**. This breadth explains why infant assessment cannot be compressed into the question, "What is the IQ?" The clinician must communicate which developmental dimensions were sampled and which require follow-up.

6.6 Selecting and reporting measures when access is uneven

Test choice should begin with access to the task. Vision, hearing, motor control, expressive language, familiarity with materials, schooling, and cooperation can all change what a score represents. For screening in intellectual disability alone or with visual impairment, the stated pairing is DST with VSMS. With hearing impairment, the guideline pairs Gesell's Drawing Test and Seguin Form Board with VSMS; with cerebral palsy or locomotor difficulty, it pairs DST and Gesell's Drawing Test with VSMS. These are selection aids, not substitutes for clinical judgement about whether the child could genuinely engage with each task.

The Binet-Kamat Test may be unsuitable where its verbal and language demands dominate the child's route to a response. Detailed selection among IQ instruments belongs with formal clinical and IQ assessment; the important point here is to resist an instrument's familiar name when its route of administration is mismatched to the child. A defensible report says what was selected, why it was selected, what barriers remained, and how the barriers temper the conclusion.

A good report makes uncertainty useful. State the construct assessed, the source of information, the settings represented, any access barriers, and the practical implications of the profile. Avoid converting a screening estimate into a diagnostic certainty, an informant account into an objective observation, or an adaptive score into a legal percentage. The final formulation should identify present abilities, difficulties, supports, and the next assessment step. It should also state which conclusions are well supported, which remain provisional, and what observation or collateral history would reduce the uncertainty. That is more clinically valuable than a crowded list of scales.

■ **TRAP** **Examination trap:** DST, VSMS, and Gesell's Drawing Test are not interchangeable IQ tests. They respectively address developmental screening, social maturity and adaptive profile, and developmental screening.

The causes of intellectual disability

7 · Etiology of intellectual disability (genetic, prenatal, perinatal, postnatal)

Aetiology is clinical work, not a label-collecting exercise. Intellectual disability (ID) is the final common developmental outcome of heterogeneous influences on the developing brain. The

useful question is not merely “what diagnosis does this person have?”, but whether the history, examination and targeted investigations can identify a cause that changes care, anticipatory guidance, recurrence discussion, or surveillance. A causal formulation should preserve timing, mechanism and certainty. It should also distinguish a demonstrated cause from an associated finding and from an unexplained presentation.

For the examination candidate, a timing-based account gives a dependable spine: prenatal influences, events around birth, and postnatal acquired influences. This is not a claim that every presentation fits a clean compartment. Developmental vulnerability may begin before birth and become clinically apparent later; genetic susceptibility and environmental exposure can coexist. The classification is useful because it structures the history, directs investigation, and keeps preventable acquired causes visible without reducing a complex developmental history to a single event.

50-70% ESTIMATED PRENATAL CONTRIBUTION	10-20% ESTIMATED PERINATAL CONTRIBUTION	5-10% ESTIMATED POSTNATAL CONTRIBUTION	25-50% GENETIC CAUSES OVERALL
---	--	---	--




Fig 21.6 The aetiological timing map of intellectual disability. The causes of ID sorted by when the insult falls, and the reason a single answer does not exist: three sources cut the same axis with different category boundaries, different populations and different denominators, so their proportions are reported lane by lane and never merged. The plate teaches three things at once. Genetic is a mechanism that sits inside the prenatal column, not a fourth time slot. Severity shifts the prenatal and genetic share upward without licensing the conclusion that a cause is found more often. And the fraction with no mechanism in it rivals or exceeds every named mechanism in both schemes that quantify it. Shorter Oxford quantifies no unknown fraction, and none is inferred for it here.




Fig 21.7 The aetiological workup in intellectual disability. The sequence a clinician actually runs: clinical assessment first, then a single gate that asks whether the aetiology is already obvious, then first-line testing, where the Western and the Indian guideline diverge and both orderings are drawn without either being called correct. Each test carries its own sourced diagnostic yield, and the yields are deliberately shown to be small. The workup ends not at a threshold but at a referral, because no source used here supplies a numeric stopping rule, and because an aetiology is found in only about half of the people tested.

7.1 A timing-based causal framework

Prenatal aetiology includes genetic causes, congenital malformations and adverse intrauterine exposures. Across all severities, prenatal factors are estimated to account for 50-70% of cases. The figure is best read as an orientation, not as a prediction for an individual patient. It combines genetic and environmental pathways, and its apparent size will vary with ascertainment, availability of neonatal care, diagnostic methods and the population being studied.

Perinatal aetiology refers to influences operating around birth. It is estimated to contribute 10-20% of ID overall. The central clinical discipline is temporal: obtain the obstetric and neonatal narrative carefully, but do not convert a difficult delivery into a causal conclusion without corroborating developmental, neurological and investigation findings. A birth complication may be relevant, incidental, or one part of a broader developmental disorder.

Postnatal aetiology comprises acquired influences after birth that affect brain development. It is estimated to account for 5-10% of cases. In practice, the history asks whether there was a previously acquired skill that was lost, a period of severe illness or injury, or evidence that development changed trajectory. Such questions protect against prematurely calling a presentation genetic or unexplained.

Causal language should match the evidence. A documented exposure or event can be part of the history without being sufficient to explain the developmental phenotype. Concordance matters: the proposed timing, neurological findings, developmental course and investigation results should tell a coherent story. Where they do not, retain alternative explanations and revisit the formulation as new records or findings become available. This avoids both diagnostic anchoring and the equally unhelpful practice of listing every historical event as a cause.

■ **RULE** **Timing is a map for enquiry, not proof of causation.** The history proposes a mechanism; examination and appropriately chosen tests decide how securely it can be attributed.

The categories also overlap in a meaningful way. A congenital brain malformation is prenatal in origin even if recognised only after delivery. A genetic condition may make a child more vulnerable to an intercurrent insult. Conversely, an exposure can cause a developmental phenotype without yielding a pathognomonic laboratory result. Good formulation therefore states the strongest supported explanation and records residual uncertainty honestly.

7.2 Reading frequencies without misreading the patient

Published distributions are not interchangeable. In a broad classification, prenatal genetic disorders were estimated at **4-28%**, congenital malformations at 7-17%, and prenatal exposure at 5-13%. The wide ranges are themselves the lesson: aetiological proportions are shaped by what a service can recognise and whom it sees. The same classification estimated perinatal causes at 2-10%, postnatal causes at 3-12%, and unknown causes at 30-50%.

Severity also changes the diagnostic landscape. Prenatal aetiology has been reported in **59-73%** of severe ID and in 23-43% of mild ID. This does not make mild ID “non-biological”, nor does it authorise an assumption that a severe presentation must be genetic. It means the threshold for

pursuing a careful aetiological workup should remain high across presentations, while the pattern of findings and the probable yield may differ.

CLINICAL QUESTION	WHAT IT PROTECTS AGAINST	HOW TO USE THE ANSWER
When did concern first become apparent?	Assigning timing from a single memorable event	Place the concern beside pregnancy, birth and developmental history.
Was development static, delayed, or did skills diminish?	Missing an acquired or progressive process	Use the trajectory to decide how urgently neurological and metabolic causes need consideration.
Are there family, physical or neurological clues?	Calling a pattern idiopathic before examining it	Integrate inheritance, morphology and examination into the testing plan.
Is a cause demonstrated, probable, or still unknown?	Overstating causality to families	Document the degree of certainty and keep a route for review.

Unknown cause is not synonymous with inadequate care. It can reflect extreme genetic heterogeneity, limitations of the available test, incomplete historical information, or an interaction of influences that cannot be disentangled. In tertiary care, an aetiology is identifiable in 50% of people presenting with ID, yet about 50% of cases may lack a molecular diagnosis. These statements refer to different denominators and endpoints. One is a clinically identifiable cause; the other is a molecular answer. They should never be treated as competing estimates of the same thing.

7.3 Genetic architecture: why no single test tells the whole story

Genetic does not mean simple inheritance. Genetic causes are thought to account for 25-50% of ID overall, and family and twin studies suggest heritability between 30-90%, depending on diagnostic category and study design. These broad estimates should not be turned into an individual recurrence estimate. They establish that inherited and newly occurring genomic changes deserve active consideration, particularly when the phenotype, family history or examination points in that direction.

Copy number variation refers to a deletion or duplication of genomic material. Cumulatively, CNVs are described as the commonest genetic abnormality class causing ID. Array-based CNV analysis identifies microdeletions or duplications in approximately 15-20% of tested patients with ID. This is why a genomic test that can detect dosage imbalance belongs early in the assessment rather than being reserved only for those with striking dysmorphism.

Single-gene heterogeneity is equally important. Mutations in more than 700 genes are estimated to be capable of causing ID. In severe sporadic non-syndromic ID, trio-based exome sequencing has identified de novo heterozygous variants or small indels in 13-55%. The range should prevent opposite errors: neither assume that an apparently isolated case lacks a genetic explanation, nor promise that sequencing will always deliver a definitive answer.

X-linked intellectual disability provides a reason to elicit family history with attention to affected males and the maternal lineage. Mutations in known X-linked genes account for **10-12%** of males with ID, and yet no single one of those X-linked genes explains more than **0.1%** of intellectual disability. This is locus heterogeneity in practical form: a recognisable inheritance pattern does not identify the molecular diagnosis by itself. The syndrome-specific genetics and recurrence counselling belong with the relevant syndrome discussions, not with a generic workup.

Autosomal recessive intellectual disability deserves particular attention when the pedigree suggests shared ancestry or recurrence among siblings. In outbred populations it may account for 10-20% of ID; autosomal recessive disorders are reported to be up to threefold more frequent among inbred than non-inbred cases. The history must be asked respectfully and documented neutrally. Its purpose is to guide diagnosis and counselling, never to assign blame.

7.4 From phenotype to an ordered genetic evaluation

The diagnostic interview begins before the requisition form. Obtain a pregnancy and birth history, a developmental trajectory, a pedigree, previous records and the parents' account of strengths as well as difficulties. On examination, look for growth pattern, head size, minor anomalies, skin findings and focal neurological signs. The presence of at least four minor physical anomalies should alert the clinician to a probable genetic cause. The value of this threshold is not that it diagnoses a syndrome; it is that it changes the prior probability and supports a more deliberate genetic examination.

In a person without an obvious syndrome or clear extrinsic cause, **first-line genetic testing** comprises chromosomal microarray and fragile X repeat-length testing.

GOOD TO REMEMBER

These tests answer different questions and are complementary: microarray looks for chromosomal copy-number abnormalities, whereas a separate fragile X assay addresses a repeat-length abnormality. Do not assume that one result substitutes for the other merely because both are called genetic tests.

Karyotyping is no longer the appropriate routine test when chromosomal microarray is available. It remains indicated when a specific aneuploidy or a balanced translocation is suspected. This is a test-selection distinction, not a hierarchy of “old” and “new” medicine: the suspected abnormality determines which technique can answer the question.

- **TRAP Examination trap:** a normal sequencing result does not exclude every genetic mechanism. Currently available sequencing tests do not detect **repeat expansions and imprinting disorders**.

After initial testing, a negative result should trigger re-phenotyping and **expert clinical genetics consultation**, rather than indiscriminate laboratory panels and imaging. A genetics assessment can refine the phenotype, determine whether a different genomic method is appropriate, revisit

the pedigree and frame the residual uncertainty for the family. This is also the right setting for generic counselling: explain what is known, what remains unresolved, and why recurrence discussion must be based on the established or suspected mechanism rather than on a generic population figure.

Test ordering should remain hypothesis-led. A phenotype can make a particular mechanism more plausible, but it should not be used as a gate that excludes broad genetic evaluation solely because dysmorphism is subtle or family history is unrevealing. Conversely, a long list of investigations is not a substitute for a better history and examination. Reassessment after each result is often more informative than simply adding tests: ask whether the new finding explains the developmental picture, whether it is concordant with the family history, and whether it creates an actionable surveillance or counselling question. This is how the workup stays both clinically rigorous and proportionate.

A negative initial genetic screen ends neither the clinical search nor the duty to explain uncertainty clearly.

7.5 Metabolic and neuroimaging pathways

Inborn errors of metabolism are individually uncommon but collectively affect 1-1.5% of the general population. Their importance in ID is disproportionate to their frequency because some are potentially modifiable and because clinical clues may be subtle. First-pass metabolic screening accounts for up to approximately 2.5% of screened patients with ID. A low yield is not an argument for abandoning clinical reasoning; it is an argument against indiscriminate testing in the absence of a phenotype-driven question.

When broad screening has already been undertaken, second-line metabolic testing identified an IEM in **14%** of those tested, many of which were treatable. Progressive change, loss of psychomotor skills, episodic deterioration, unusual neurological features or a family pattern should therefore prompt reconsideration of metabolic disease. The detailed disorders and their treatment are addressed in the section on inborn errors of metabolism; here, the point is to keep an aetiology that may alter outcome within the diagnostic field of view.

Neuroimaging is clinically indicated with microcephaly or macrocephaly, abnormal head circumference, seizures, focal neurological signs, progressive cognitive impairment, or loss of psychomotor skills. MRI is generally superior to CT, except when acute haemorrhage or calcification is suspected. Imaging must answer a clinical question. Subtle nonspecific findings may support altered neurodevelopment without establishing a diagnosis, whereas a structural abnormality may reorganise the entire workup.

Guidelines can differ by setting and resource pathway. The candidate should state the local recommendation accurately while retaining the principle that testing is most useful when integrated with phenotype and clinical trajectory.

IN INDIA

Where obvious clinical aetiologies are not found, the Indian Psychiatric Society guideline designates **brain MRI with MRS and advanced metabolic tests using GCMS and TMS** as mandatory investigations.

7.6 Prenatal, perinatal and acquired causes in the clinical narrative

After the genetic and structural possibilities have been considered, return to the developmental history and ask what may have acted before birth, around birth or after birth. Prenatal exposure categories include maternal infection, teratogen exposure and toxemia. The purpose of asking is to establish timing and biological plausibility, not to retrospectively place moral responsibility on a mother or family. The same non-judgemental stance applies to histories of substance exposure, obstetric difficulty, neonatal illness and later acquired insults.

Perinatal history should be reconstructed from records wherever possible. The more remote the event and the less specific the account, the more cautious the attribution should be. The examiner's answer becomes stronger when it separates a reported event from evidence of its neurological consequence. Likewise, a postnatal history should seek changes in development, neurological symptoms and the sequence between illness and altered functioning. Aetiological formulation can then state whether the acquired explanation is established, plausible or insufficiently supported.

- Start with timing, then ask what mechanism could link the event to altered development.
- Seek corroboration from records, examination and the developmental trajectory before naming causation.
- Keep genetic and acquired explanations open together when the evidence supports neither as exclusive.

This approach matters especially when families arrive with a single explanatory story, such as a difficult delivery or a relative with developmental problems. Both may be important. Neither should stop the clinician from completing the rest of the formulation. The respectful task is to test the story, preserve what is supported, and correct what is overconfident without making the family feel dismissed.

7.7 Aetiological formulation and communication

A useful written formulation has a layered structure: developmental phenotype, best-supported timing category, demonstrated mechanism where available, associated comorbid findings, and a statement of certainty. It is better to write "ID with a probable prenatal genetic aetiology under evaluation" than to attach a speculative syndrome name. This language guides follow-up and is safer for families, schools and other clinicians who may read the record later.

Explain the purpose of testing before ordering it. Families should understand that testing may identify a diagnosis, clarify future health needs, inform recurrence discussion, or remain non-diagnostic. They should also be prepared for findings whose significance is uncertain. Counselling is a process of interpretation and decision support, not simply the handing over of a laboratory

report. When a family pattern is suspected, referral allows the family to discuss who else may benefit from assessment without converting a probability into a certainty.

The clinical payoff is precision with humility. Aetiology helps prevent a treatable contributor from being missed, prevents unnecessary investigations from accumulating, and makes recurrence counselling more honest. Yet the person's current needs do not wait for molecular closure. Diagnostic workup should proceed alongside support planning and management, with each service informed by the level of certainty rather than delayed by it.

8 · Down syndrome and other chromosomal causes

Why this cause deserves its own clinical frame. Down syndrome is the chromosomal condition most likely to turn a general discussion of intellectual disability into a real consultation: the clinician has to recognise a developmental profile, separate appearance from diagnosis, look for treatable contributors to disability or behavioural change, and explain why the cytogenetic result matters to a family. It is therefore not enough to say that the person has an extra chromosome. The mechanism determines the meaning of the result for relatives and future pregnancies.

For the psychiatry candidate, the useful mental model is a sequence: establish the syndrome and its chromosome mechanism; understand the pattern of associated health needs; then interpret new symptoms against the person's usual functioning. Intellectual disability is part of the presentation, but its definition, assessment and severity grading belong in their respective sections. Equally, a behavioural change should not be dismissed as an inevitable consequence of the syndrome. It may be the presentation of an associated medical disorder, sensory loss, or a co-occurring psychiatric condition.




Fig 21.8 The three cytogenetic mechanisms of Down syndrome. Free trisomy, translocation and mosaicism read across from the karyotype to the only thing that changes management, which is what you tell the parents about the next pregnancy. The plate is built so the three columns diverge exactly where the counselling diverges: only the translocation column earns a parental karyotype, and only that column ends in recurrence figures the sources actually supply. The other two columns end in a named gap, because this chapter's corpus states no recurrence risk for free trisomy 21 and none for mosaicism, and the plate declines to supply from memory the figure a viva asks for. The maternal-age effect is shown last and bound to one mechanism only.

8.1 Epidemiological position and clinical significance

Down syndrome is a leading chromosomal cause of intellectual disability and is clinically important precisely because it is common enough to be encountered across child, adult and liaison settings. Textbook birth estimates differ: Shorter Oxford gives about **1 in 650** live births, whereas Tasman gives **1 in 800** live births. These are not competing bedside tests. They illustrate that a quoted frequency depends on the source population and the outcome counted. A figure

expressed per pregnancies must not be treated as interchangeable with a live-birth figure: a cited source estimates trisomy 21 at 1.5 in 1000 pregnancies.

Within the population of people with intellectual disability, Down syndrome accounts for between **6-8%**. That burden explains why psychiatrists need fluency in its medical context, not only recognition of its phenotype. The consultation often occurs at a transition point - concern about development, a new behavioural presentation, family anxiety about recurrence, or declining function in an adult - when an apparently familiar label can obscure the actual clinical question.

95% FREE TRISOMY 21	about 4% TRANSLOCATION	about 1% MOSAICISM
-------------------------------	----------------------------------	------------------------------

8.2 Cytogenetic architecture: the mechanism is the counselling clue

■ **RULE** State the chromosome mechanism before discussing recurrence: the phenotype may be similar, but the family implication is not.

Down syndrome results when there is extra chromosome 21 material. In **full trisomy 21**, the additional chromosome is present as a separate chromosome. This is the usual arrangement and, in the source cited here, applies to nearly 95% of people with Down syndrome. The extra material arises through a chromosome-separation error, conventionally termed **nondisjunction**. The molecular event makes it essential to distinguish this arrangement from a structural rearrangement identified in a parent.

In **translocation Down syndrome**, the extra chromosome 21 material is attached to another chromosome, usually chromosome 14, designated t(14;21). The total count can still be 46 chromosomes, but chromosomal material from 47 is present. This is the exam discriminator: counting chromosomes alone does not tell the whole genetic story. The relevant question is whether the chromosome complement is balanced in a parent or unbalanced in the child. The answer changes the recurrence discussion and is why a cytogenetic report must be read rather than merely filed.

Mosaicism means that different cell lines are present. Its presence should prompt precision in description rather than casual prediction from appearance or a presumed developmental outcome. The safe clinical teaching point is not to manufacture a hierarchy of functioning from a mechanism alone. Mechanism informs counselling; it does not replace individual assessment of development, communication, adaptive functioning and health.

CHROMOSOME ARRANGEMENT	WHAT THE REPORT MEANS CLINICALLY	WHY PSYCHIATRY SHOULD CARE
Full trisomy	An extra chromosome 21 is present as a separate chromosome.	It frames the syndrome but does not predict an individual's support needs.
Unbalanced translocation	Extra chromosome material is attached to another chromosome.	Parental chromosome studies become central to the family discussion.
Mosaicism	Different cell lines are present.	Do not infer clinical ability from the label alone; assess the person.

The accompanying figure is useful as a memory aid because it puts these arrangements beside their counselling consequence. It should prevent a common error: treating every Down syndrome result as if it carried the same familial implication.

Read the report in a clinically disciplined way. **Cytogenetic interpretation** starts by identifying whether it describes a separate extra chromosome, attached extra material, or differing cell lines. Next ask whether the wording establishes a structural rearrangement that could also be present in a parent. Finally, distinguish what the report establishes from what still needs direct clinical assessment. A chromosome result explains the syndrome's mechanism; it does not by itself describe communication, capacity, family resources, psychiatric symptoms or the cause of a new behavioural change. This prevents genetic language from becoming a shortcut around formulation.

The same distinction helps when communicating with other clinicians. “Down syndrome confirmed” may be adequate as a diagnostic headline but is incomplete as a handover when the mechanism has not been stated. Record the arrangement in words, note whether parental studies are relevant, and separate known information from questions that remain open. This makes later counselling more coherent and stops a vague family-history statement from acquiring unwarranted certainty as it passes between services.

8.3 Origin of the extra material and maternal-age reasoning

Most molecularly studied trisomy 21 cases have an extra chromosome of maternal origin, reported as 92%. Maternal meiotic error is therefore the dominant framework, but a candidate should avoid turning that into an indiscriminate statement about every pathway. Around 75% of nondisjunction errors are reported at the first maternal meiotic division, although the cited source does not make the denominator sufficiently explicit to claim more than that. In contrast, 60% of paternal meiotic errors occur at the second meiotic division, and mitotic error is estimated in around 5% of trisomy 21 cases.

The maternal-age association belongs mainly to the maternal meiotic pathway. **Paternal and mitotic errors do not show an age effect, and Robertsonian formation is not related to parental age.** This distinction matters because a risk factor should be linked to the biological pathway it plausibly modifies, not applied as a universal explanation after the event.

Age-specific risk rises substantially. Shorter Oxford reports about 1 in 2000 live births among mothers aged 20-25, compared with 1 in 30 among those aged 45 years or over. Another table gives 1 in 2000 at maternal age 30 and 1 in 100 at age 40. The direction is the high-yield point, not the temptation to merge differing source anchors into a fabricated universal rate. Women in their forties have at least a 15-fold increase in risk compared with women in their twenties.

■ **TRAP** Do not confuse high age-specific risk with where most affected children are born.

Up to 80% of children with Down syndrome are born to mothers under 35. This is not a paradox. Population impact depends on both individual risk and the number of births in each age group. It is the cleanest way to explain why counselling cannot be reduced to stereotypes about maternal age.

8.4 Phenotype, development and associated health needs

The physical phenotype is a recognition aid, not a diagnostic algorithm. Individual signs vary, and no individual sign is universal. The most consistent physical feature, upslanting palpebral fissures, is still restricted to 80% of people with Down syndrome. A transverse palmar crease occurs in 53%, while a protruding tongue is present in fewer than 50%. The practical implication is simple: absence of a familiar sign should not overrule a chromosome result, and presence of an isolated sign should not be treated as proof of the syndrome.

Intellectual ability varies considerably. Tasman reports a mean IQ of approximately 50, but the number is not a prognostic verdict for an individual and must not be used to grade intellectual disability. The clinician should describe present skills, communication, daily functioning and the support environment. The sections on clinical assessment, IQ testing and adaptive behaviour explain how to do that without mistaking a test score for the person.

ASSOCIATED FINDING	REPORTED FREQUENCY	CLINICAL CONSEQUENCE IN A PSYCHIATRIC ASSESSMENT
Congenital heart disease	40-50%	Medical history and exertional symptoms shape both assessment and treatment planning.
Gastrointestinal atresias	12%	Past surgical and nutritional history can clarify the developmental context.
Hearing loss	75%	Often conductive and related to otitis media; it can mimic inattention, withdrawal or a language problem.
Ocular findings	approximately 50%	Visual difficulty may alter engagement, learning and apparent behaviour.
Gluten intolerance	4-20%	Gastrointestinal symptoms should not be reduced to a behavioural complaint.
Autoimmune thyroid dysfunction	up to 20%	In adolescents and adults, consider it when behaviour, mood or cognition changes.

GOOD TO REMEMBER

These are not decorative associations. Hearing and visual difficulty alter how a person is read in the clinic. Pain, discomfort or endocrine disturbance may appear as irritability, reduced participation, slowing, sleep change or loss of previously reliable skills. In particular, thyroid dysfunction is a crucial differential when behavioural disturbance, depression or an apparent dementia presentation emerges. Start with a change from baseline and ask what has changed in the body, the sensory environment and the person's routines before assigning a psychiatric explanation.

Some children with Down syndrome also meet criteria for autism spectrum disorder, estimated at approximately 7%. Autism should be assessed using its own diagnostic framework. The diagnostic criteria, clinical features and differential diagnosis of autism are dealt with in the autism sections of this chapter. Here the point is narrower: a known chromosomal diagnosis does not make an additional neurodevelopmental diagnosis impossible, nor does it make assessment dispensable.

8.5 A psychiatry assessment that does not miss the treatable problem

■ **RULE** A new symptom is a clinical question, not an automatic feature of Down syndrome.

Begin by identifying the change in observable terms: what is different from the person's own baseline, where it occurs, and what has altered in eating, sleep, activity, communication, comfort or social participation. A **baseline comparison** prevents the interview from confusing a lifelong trait with a fresh symptom. Establish the time course from people who know the person well, but

do not let collateral description substitute for direct engagement. A quieter person may be depressed, unwell, hearing less well, overwhelmed, or responding to a changed environment. These possibilities require clinical discrimination.

- Review hearing and vision when apparent inattention, language difficulty or withdrawal is new or worsening.
- Ask about physical discomfort and gastrointestinal symptoms when behaviour becomes irritable or avoidant.
- Consider thyroid dysfunction when there is behavioural disturbance, depressive symptomatology or cognitive decline.
- Keep autism assessment criterion-based rather than treating the chromosomal diagnosis as either proof or exclusion.

This approach guards against diagnostic overshadowing, discussed fully in the dual-diagnosis section. The syndrome-specific safeguard is especially important here because associated health conditions can change behaviour without announcing themselves in conventional psychiatric language. The clinical record should preserve the observed baseline, the reported change, the relevant physical and sensory questions, and the working explanation. Doing so makes it possible for the next clinician to see whether the proposed explanation accounted for the change or merely accompanied it.

MNEMONIC

CHROME

: **C**hromosome mechanism; **H**earing and vision; **R**eview change from baseline; **O**rgan-system contributors; **M**ood and mental-state assessment; **E**xplain the recurrence implication. Use it as an interview sequence, not as a substitute for examination.

When behaviour changes in Down syndrome, look for a change in health, sensory input or environment before attributing it to the syndrome.

8.6 Family implications of translocation Down syndrome

The family conversation becomes most specific when a translocation is identified. In about half of translocation Down syndrome cases, a parent has a balanced translocation; a related source estimates that around 45% of Robertsonian translocation Down syndrome cases are inherited from a balanced carrier parent. The practical message is not that every parent is a carrier, but that parental chromosome status is material to that particular family's recurrence discussion.

For a balanced **t(14;21) translocation**, the cited recurrence risk for a subsequent child with Down syndrome is **1 in 10** when the mother carries the translocation and **1 in 20** when the father carries it. These are mechanism-specific figures. They must not be offered as the risk after ordinary full trisomy, nor copied into a discussion before the parental studies have established the relevant arrangement.

A familial balanced **rob(21;21) fusion** has a qualitatively different implication: all offspring will have Down syndrome (100%). This is exactly why wording matters. “Down syndrome runs in the family” is too imprecise to guide care. The chromosome report, the parent who carries any balanced rearrangement, and the specific rearrangement together determine what can responsibly be said.

Refer families for genetic counselling and interpret recurrence as a syndrome-specific issue, not as a generic estimate. The general approach to aetiological investigation and counselling referral is covered in the etiology of intellectual disability section. The psychiatrist's contribution is to recognise when the mechanism makes that referral urgent, to communicate uncertainty honestly, and to ensure that distress does not prevent the family from receiving a clear explanation.

8.7 Examination synthesis: how to write the answer safely

A strong answer begins with Down syndrome as a chromosomal cause of intellectual disability, then moves from cytogenetics to clinical care. Describe full trisomy, translocation and mosaicism as different chromosome arrangements; explain that translocation has particular implications for parental studies and recurrence; then show why sensory, endocrine, gastrointestinal and cardiac associations matter to psychiatric formulation. Finish with the warning against diagnostic overshadowing. This structure demonstrates reasoning rather than reciting a facial-feature list.

■ **TRAP** The classic examination mistake is to give a single recurrence figure for every form of Down syndrome.

That answer is unsafe because it erases the distinction that the cytogenetic result is meant to reveal. A further mistake is to treat the physical phenotype as diagnostic while ignoring the associated health review. The better answer states that phenotype prompts recognition, cytogenetics establishes mechanism, and longitudinal assessment identifies the problem requiring action now.

Other chromosomal causes of intellectual disability need syndrome-specific evidence on their phenotype, mechanism and recurrence patterns. This fragment deliberately does not generalise Down syndrome data to them. Where a person has a chromosomal finding without the characteristic Down syndrome arrangement, the clinical task is to read the individual report and pursue the aetiological pathway rather than forcing it into a familiar template.

9 · Fragile X syndrome

Why this syndrome deserves a deliberate search. Fragile X syndrome is a recognisable cause of intellectual disability, but it cannot safely be reduced to a facial gestalt or a single childhood phenotype. It is the most common inherited cause of intellectual disability, while among all specific causes it is described as second to Down syndrome. Its importance extends beyond the proband: the same locus can produce a developmental disorder in one family member and late-life neurological or reproductive disorders in relatives who carry a premutation.

The useful clinical stance is therefore syndromic and family-based. Recognise the developmental and behavioural pattern, examine without over-relying on dysmorphology, confirm the molecular finding with an appropriate repeat-expansion assay, and explain that molecular category and parent of origin determine the implications. The accompanying figure can be used to hold the relationship between repeat expansion, gene silencing and the distinct premutation disorders in mind.




Fig 21.9 The FMR1 CGG repeat ladder. The three repeat classes this corpus supports, what each one carries, and the fact that neither boundary between them can be stated as a single number.

9.1 Molecular lesion and its consequences

Fragile X syndrome results from loss of function of **FMR1 (fragile X mental retardation gene 1)**, located at **Xq27.3**. The unstable sequence is a **CGG (cytosine-guanine-guanine)** repeat in the promoter-adjacent 5' untranslated region. It is worth retaining the anatomical wording rather than insisting on a single exon label, because the sources differ at that finer level while agreeing that the repeat is non-coding and close to the promoter.

Repeat expansion accounts for **more than 95%** of affected cases. In the full-mutation state, the expanded sequence and neighbouring regulatory region become hypermethylated; histone deacetylation and repressive chromatin then prevent transcription. Thus the mutation is not simply a long piece of DNA. It converts a transcriptionally active locus into one that is functionally silent. The extent of methylation is directly related to the loss of functional protein, which helps explain why repeat size alone is not a complete clinical predictor.

The protein product, **FMRP, an RNA-binding protein**, normally binds polyribosomes in dendrites and synaptic spines and largely restrains local translation of synaptic proteins. Loss of that restraint provides a coherent bridge from molecular change to altered synaptic maturation, learning, attention and behaviour. It also prevents a simplistic conclusion that every clinical feature is fixed at birth: the disorder concerns regulation of developing neural systems and its expression changes with development.

■ **RULE** Full mutation means loss of FMRP through epigenetic silencing; premutation disorders arise through excess FMR1 messenger RNA, not a mild form of the same process.

This distinction is the organising principle of the chapter. In a premutation, the locus is not silenced in the same way. Instead, **RNA-mediated toxicity from overproduction of FMR1 mRNA** is implicated. Calling a premutation carrier “mildly affected with fragile X syndrome” is therefore biologically misleading and clinically unhelpful. One state is principally a protein-loss developmental disorder; the other is an RNA gain-of-function state with its own age-related and reproductive associations.

9.2 Repeat categories, instability and anticipation

Repeat-length bands are useful in laboratory interpretation, but their lower boundaries vary modestly across reference texts. A normal stable allele is reported in ranges such as 5-55 CGG repeats, while a commonly used premutation band is **55-200 repeats**. The clinically robust distinction is that a full mutation is generally defined as more than 200 repeats, although a cited source uses a higher threshold. A report must be interpreted using the laboratory's stated classification rather than by treating a disputed boundary as a universal biological cliff.

Normal alleles tend to be stable. **AGG interruptions** interspersed within the repeat are proposed to stabilise the sequence by reducing secondary structures during replication. Long uninterrupted stretches are more prone to expansion. This is why molecular counselling cannot be reduced to a family member's label alone: the repeat architecture and its transmission context matter.

The inheritance is **X-linked**, but it does not behave like a simple static X-linked mutation. Expansion from a premutation to a full mutation occurs through the female germline; male carriers usually transmit without further expansion and may show contraction in sperm. This parent-of-origin effect explains the apparently paradoxical pedigree in which an unaffected transmitting man has daughters who are carriers, yet the more severely affected descendants emerge through subsequent maternal transmission.

■ **TRAP** Do not infer full-mutation risk from “X-linked” alone. The relevant question is whether the premutation is being transmitted through a woman, because that is the route in which expansion to a full mutation occurs.

The historical pattern of worsening expression in successive generations is called the **Sherman paradox**. It was an observation before it had a molecular explanation. The repeat-expansion mechanism now turns that observation into a counselling problem: a pedigree should be drawn across generations, noting cognitive and developmental histories as well as tremor, ataxia and early ovarian failure in relatives. Recurrence counselling should be syndrome-specific and undertaken with clinical genetics support; the generic framework belongs in the etiological assessment discussion.

- Ask which side of the family carries the molecular result and whether transmission has been maternal or paternal.
- Obtain an extended pedigree history rather than stopping at developmental delay in the child.
- Explain that a premutation and a full mutation have different clinical implications.
- Offer cascade evaluation to relatives through an appropriately supported genetics service.

9.3 Population burden and the diagnostic place of fragile X

Published prevalence estimates differ because ascertainment and molecular methods have changed. A sensible examination answer is to state the estimate with its denominator: fragile X syndrome occurs in about 1 in 4,000 males and around 1 in 8,000 to 1 in 9,000 women. Female prevalence is described as approximately one-half of the male rate. These figures refer to the syndrome, not to premutation carriage, which is much more frequent and has a different sex distribution.

1 in 4,000 MALES WITH SYNDROME	1 in 8,000-9,000 WOMEN WITH SYNDROME	~40% MALE PREMUTATION CARRIERS WITH FXTAS	~20% FEMALE PREMUTATION CARRIERS WITH POF
--	--	---	---

The distinction between affected syndrome and carrier state changes the conversation with families. Premutation carriage is reported at 1 in 150-200 females and 1 in 400-450 males in the general population. It should not be quoted as the prevalence of fragile X syndrome, nor should syndrome prevalence be used to dismiss the relevance of a premutation in an extended family. Fragile X is also clinically relevant in autism assessment. It is described as the most commonly known single genetic cause of autism spectrum disorder. The direction of the statistic is the trap:

the proportion of people with fragile X who meet autism criteria is not the same as the proportion of autism attributable to fragile X. Reported rates of autism-spectrum diagnosis within fragile X vary substantially with the diagnostic construct and method, so they should be read as evidence of overlap rather than as a single invariant percentage. The medical and genetic workup of autism is addressed in the autism workup section.

9.4 Recognising the phenotype without overcalling it

The physical phenotype is developmental and variable. In adults, the familiar pattern includes an elongated face, prominent or cupped ears, a high arched palate and connective-tissue features.

Large anteverted ears, smooth velvet-like skin, hyperextensible finger joints, flat feet and mitral valve prolapse are the useful connective-tissue cluster. Macro-orchidism becomes relevant after pubertal development and should be asked about and examined for sensitively, with appropriate consent and privacy.

ADULT PHYSICAL FEATURE	APPROXIMATE FREQUENCY AMONG ADULT CASES	CLINICAL USE
Macro-orchidism	~90%	Most evident after puberty
Elongated face	~80%	Supports, but never establishes, suspicion
Prominent or cupped ears	~65%	Look for it in the broader phenotype
High arched palate	~60%	Examine as part of a general dysmorphology review
Flat feet	~60%	Connective-tissue clue rather than a diagnostic sign

GOOD TO REMEMBER

These are group frequencies, not a checklist diagnosis. Up to 20% of affected children may have no obvious physical features, while individual facial or connective-tissue traits occur in many people without the syndrome. Developmental history, family history and a coherent behavioural phenotype raise suspicion; molecular testing establishes the diagnosis.

Associated physical morbidity deserves ordinary clinical attention. Seizure disorder occurs in 20% of individuals with fragile X syndrome. Visual and hearing assessment, cardiac review when indicated, and review of neurological symptoms should be integrated into care rather than deferred because the presenting concern is developmental. The specific diagnostic approach to epilepsy in autism belongs to the autism medical-workup section, but a seizure history in fragile X is directly relevant to this syndrome review.

9.5 Cognitive, language and behavioural profile

In males with a full mutation, intellectual disability is often moderate, with an average IQ of **about 40**; individual functioning is nevertheless broad and must not be assumed from a molecular result. In females with a full mutation, random X inactivation offers partial protection,

and cognitive impact is correspondingly variable. The assessment of IQ, adaptive behaviour and intellectual-disability severity is covered elsewhere in this chapter; here, the key point is that the fragile X profile is uneven, not merely globally delayed.

Short-term memory, visuomotor coordination, arithmetic and sequential processing are relative weaknesses, while simultaneous processing and long-term storage or retrieval may be relative strengths. This matters clinically. A child may do better when information is presented as a whole visual structure than when it is delivered as a long serial verbal sequence. It also matters in the consultation: apparently inconsistent performance can be a profile effect rather than evidence of poor effort or oppositionality.

The behavioural phenotype has a distinctive social-anxious quality. **Gaze aversion or reduced eye contact, shyness and social anxiety, hand-flapping and repetitive mannerisms, hyperactivity with short attention span, and poor peer relations** can co-occur. These features invite diagnostic shadowing in both directions: gaze avoidance and repetitive behaviour can be assumed to establish autism, while an established genetic diagnosis can lead the clinician to overlook a co-occurring autism-spectrum condition. Evaluate the person's actual social communication, restricted behaviour and functional needs rather than using the syndrome label as a shortcut.

Disruptive behaviour is reported to peak in preschool years and to decline progressively with age, although attentional difficulties may remain. This course is useful when setting family expectations: a difficult early behavioural phase does not predict a static trajectory. ADHD itself and its formal assessment are dealt with in the ADHD and specific learning/communication disorders notes. In fragile X, practical supports should take account of attention, anxiety, sensory responses, communication and the relative cognitive profile rather than treating "behaviour" as a single symptom.

MNEMONIC

FRAME the consultation

Family transmission, Repeat category, Appearance across development, Mind and behavioural profile, Extended-family premutation effects. This is a clinical sequence, not a diagnostic score.

9.6 Females, mosaicism and the premutation family phenotype

Female presentation requires particular care because it is often quieter and more variable.

Random X inactivation (lyonization) protects many full-mutation females from full clinical expression. About half of females with the full mutation show cognitive deficits. A clinician should therefore neither assume that all female relatives are unaffected nor equate absence of intellectual disability with absence of a meaningful molecular result.

Mosaicism is described in 15-20% of individuals with FMR1 mutations. Different cell lines can carry different molecular profiles, which is one reason phenotype, repeat length and methylation pattern may not align in a simple linear fashion. The clinical response to an apparently

discordant result is not to reject it, but to interpret it with the reporting laboratory and genetics team in the context of the person's functioning and family history.

Premutation-associated disorders are not peripheral footnotes. **FXTAS (fragile X-associated tremor/ataxia syndrome)** is a late-onset neurodegenerative syndrome in premutation carriers, characterised by intention tremor and cerebellar ataxia, with later neuropathy, autonomic symptoms and cognitive or behavioural change in some individuals. Its median onset is 60 years. When an adult relative of a child with fragile X reports these symptoms, the family history has become a neurological referral issue, not merely background information.

Female premutation carriers may also develop premature ovarian failure, also termed fragile X-associated primary ovarian insufficiency. Around **20%** are reported to develop premature ovarian failure. The risk does not rise in a simple straight line with repeat size, so counselling should avoid a crude "more repeats always means more ovarian risk" message. Reproductive, neurological and developmental implications should be held together in family counselling while respecting the different needs of each relative.

9.7 Confirmation, interpretation and management

Confirmation is molecular, not morphological. The diagnostic method of choice is **DNA-based fragile X repeat-length testing, using PCR and/or Southern blot methods including repeat-primed PCR**. Cytogenetic fragile-site methods are historical: culturing lymphocytes without sufficient folic acid made an apparent X-chromosome breakpoint visible, but that laboratory phenomenon is not a treatment rationale. In particular, folic acid should not be prescribed as therapy for the primary disorder on the basis of the old cytogenetic technique.

Testing is especially appropriate when intellectual disability is unexplained, when the developmental and behavioural pattern is suggestive, or when the pedigree contains compatible developmental, neurological or reproductive histories. The broader etiological workup for intellectual disability is covered in the causes-and-workup discussion; this section's responsibility is to ensure that the fragile X assay is a dedicated repeat-expansion test. Routine sequencing alone may not detect repeat expansions, so the test request must match the suspected mechanism.

The written report should be read as a molecular category, not merely as a positive or negative result. A full mutation directs attention to the developmental phenotype and its supports; a premutation redirects the family conversation toward transmission, reproductive implications and later neurological symptoms. Mosaic findings require correlation rather than dismissal. This interpretive step protects against two opposite errors: making the molecular result the whole explanation for a person's needs, or treating a clinically relevant finding as incidental because the current phenotype is subtle.

■ **TRAP** A normal-looking child does not exclude fragile X, and a standard sequencing result is not automatically a fragile X test. Suspect from the phenotype and pedigree; confirm with the repeat-expansion assay.

There is no established treatment for the primary disorder; management is of associated conditions. This is not therapeutic nihilism. It means that care is organised around developmental assessment, communication support, educational adaptation, treatment of target symptoms and associated medical conditions, and genetic counselling for the family. Developmental interventions and educational planning belong to the management section of this chapter. A syndrome diagnosis should improve the precision of that plan, not replace person-centred assessment.

Must remember: Fragile X is an X-linked repeat-expansion disorder in which maternal transmission can convert a premutation into a full mutation; full mutation silences FMR1 and reduces FMRP, whereas premutation disorders are mediated by toxic excess messenger RNA.

10 · Inborn errors of metabolism causing ID (PKU and others)

Why this group earns disproportionate attention. Inborn errors of metabolism are individually uncommon causes of intellectual disability, but they matter because a metabolic diagnosis may change the trajectory of a child and the advice offered to the family. The useful clinical question is not whether every person with developmental difficulty has a metabolic disorder. It is whether the pattern contains enough biological coherence to make a potentially modifiable cause worth pursuing. For the psychiatrist, these disorders are also a reminder that intellectual disability can be the long-term expression of a disturbance in a biochemical pathway rather than a fixed label detached from mechanism.

Inborn errors of metabolism are best approached as disorders in which a missing or ineffective metabolic step has consequences for the developing brain and, often, for other organs. Their psychiatric relevance lies in the combination of developmental phenotype, neurological signs, behaviour, systemic clues and the possibility of pathway-directed treatment. A good answer therefore does more than list syndromes: it links substrate, enzyme, clinical pattern and the therapeutic implication.

1 in 10,000 to 1 in 20,000 PKU BIRTH INCIDENCE	first 3 months LATEST WINDOW TO BEGIN PKU DIETARY RESTRICTION	below 20% ADULTS WITH PKU PERHAPS MAINTAINING DIETARY CONTROL
--	--	--

10.1 Clinical frame: think pathway, phenotype and treatability

Metabolic disorders should not be reduced to an examination catalogue. They are a way of thinking about causation. A child may have global developmental difficulty or established intellectual disability as one part of a wider disorder, while another person may present later to psychiatry because attentional, behavioural or affective symptoms have become more conspicuous than the developmental history. The history and examination should therefore remain alert to discordance: a developmental presentation that sits alongside altered growth, pigment, eyes, skeleton, movement, episodic illness or a distinctive form of self-injury deserves a broader explanatory frame.

The pattern matters more than any isolated sign. A physical clue is not merely a dysmorphology spotter's flourish; it can identify the organ system in which the metabolic disturbance is declaring itself. Similarly, an apparently psychiatric symptom may be a direct consequence of altered neurochemistry, a response to neurological disability, or both. This is why a formulation should preserve the medical diagnosis, the developmental phenotype and the current mental-state problem as separate but connected layers.

■ **RULE** Define the problem by the disrupted pathway, then ask what can still be modified. The same developmental endpoint can arise from very different biochemical mechanisms, and treatment must follow the mechanism rather than the behavioural surface.

There is an important boundary here. The detailed sequence of aetiological tests and the general genetic-counselling process belong to the wider intellectual-disability assessment. This section instead explains the recognised metabolic conditions and the clinical reasoning that makes their recognition consequential. When a metabolic disorder is known or suspected, psychiatric care works alongside paediatrics and metabolic medicine, rather than attempting to translate a complex behavioural presentation into a purely psychiatric diagnosis.

10.2 Phenylketonuria: the prototype of preventable metabolic neurodevelopmental injury

Phenylketonuria (PKU) is an autosomal recessive disorder and remains the classic example of a biochemical defect in which early recognition changes neurodevelopmental outcome. It is described as the commonest inborn error of metabolism. Textbook estimates give an incidence between **1 in 10,000 and 1 in 20,000 births**; a further estimate is 1 in 10,000 live births. These figures are useful as scale, but the clinical lesson is larger: rarity at the level of an individual clinic does not make a reversible diagnosis unimportant.

The primary defect is deficiency of **phenylalanine hydroxylase**, a hepatic enzyme. In ordinary biochemical terms, this enzyme mediates conversion of phenylalanine to tyrosine. The relevant gene is the PAH gene, located on chromosome 12, with a reported locus at 12q22-24.1. This is not genetic detail for its own sake. It explains why the disorder is familial in pattern and why a diagnosis has implications beyond the presenting child.

When this metabolic conversion fails, the brain is affected through more than one route. Low tyrosine contributes to depletion of central monoamines, including dopamine and serotonin. Reduced availability of large neutral amino acids also compromises synaptogenesis and myelination. That dual mechanism makes sense of a phenotype that crosses developmental cognition, attention, executive control and behaviour. It also prevents a simplistic conclusion that a child who survives the early developmental period without profound disability has necessarily escaped neurocognitive consequences.

MNEMONIC**PKU clinical spine: pathway, prevention, persistence**

Start with the failed phenylalanine-to-tyrosine step; connect it to early dietary treatment; then remember that later executive, attentional and affective symptoms still deserve active review.

10.3 PKU phenotype: untreated disease and the treated survivor are different clinical questions

Untreated PKU is associated with intellectual disability and developmental delay. The recognisable untreated phenotype may include reduced pigment with fair or blond hair and blue eyes, microcephaly, eczema, epilepsy, growth retardation and hyperactivity. These features should be understood as clinical clues within a syndrome, not as a diagnostic shortcut. No one sign establishes the diagnosis, and a psychiatric assessment should not turn a physical sign into a substitute for biochemical confirmation.

The name itself reflects the secondary pathway. **Phenylketones**, produced through phenylpyruvic acid in the secondary pathway, are excreted in urine. This is a compact but useful examination link between the disease name and its biochemistry. It also illustrates a broader metabolic principle: once the usual route is blocked, diversion products can become clinically informative.

The early-treated adult or adolescent requires a different formulation from the untreated child described in older accounts. Even with early treatment, there can be subtle reductions in IQ with executive and attentional impairment. The psychiatrist should therefore ask concrete functional questions about planning, sustained attention, error monitoring, academic or occupational organisation, adherence to a demanding diet and the emotional burden of living with a lifelong disorder. This avoids two equal errors: attributing every difficulty to PKU, or dismissing cognitive and emotional concerns because gross intellectual disability has been prevented.

■ **TRAP Examination trap:** do not present early-treated PKU as synonymous with absence of neuropsychiatric morbidity. Early treatment changes the expected phenotype; it does not make follow-up unnecessary.

There is evidence of elevated depression, anxiety disorders, particularly agoraphobia, and attention-deficit disorder in early-treated adolescents with PKU compared with matched healthy individuals. In practice, this means that a symptom review needs ordinary psychiatric discipline. Establish the symptom's time course and impairment, consider medical and psychosocial contributors, and do not write off distress as an inevitable accompaniment of a metabolic label.

10.4 PKU screening and treatment: timing is part of the treatment

PKU illustrates why a metabolic diagnosis must be made early enough for the intervention to matter. Neonatal screening uses blood or urine. For prevention of severe, irreversible cognitive deficit, dietary phenylalanine restriction must begin within the **first 3 months of life**. The phrasing is deliberately temporal: a correct diet begun too late cannot be assumed to restore an already altered developmental pathway.

A low-phenylalanine diet maintained for at least the **first decade of life** is associated with an essentially normal cognitive outcome. The diet is not a casual instruction to avoid a food group. It is a sustained medical treatment whose implementation, monitoring and acceptability need specialist support.

IN INDIA

In the Indian Psychiatric Society clinical practice guidance, tetrahydrobiopterin with a low-phenylalanine diet is listed as a small-molecule intervention for PKU.

Longitudinal care needs to acknowledge that adherence can become difficult when parents no longer organise every meal and social life broadens. Perhaps **below 20%** of adults with PKU maintain dietary control throughout adulthood. This is not a reason to moralise about non-adherence. It is a reason to explore practical constraints, autonomy, cognition, family support, mood and the person's own understanding of risk. The psychiatrist's role includes making these barriers discussable while ensuring that metabolic management remains with the specialist team.

The broader lesson is that screening, diagnosis and treatment cannot be separated into unrelated facts. In PKU, the specimen identifies the condition, the biochemical pathway explains the diet, and the time window explains why developmental prevention is possible. That integrated chain is what candidates should reproduce rather than a bare list of enzyme and symptoms.

10.5 Homocystinuria: systemic clues should change the psychiatric formulation

Homocystinuria is an important contrast to PKU because the developmental presentation sits alongside striking extra-central-nervous-system findings. These include ocular lens dislocation, pectus excavatum, and a tall, thin, long-limbed Marfan-like stature. It is also associated with strokes in young and middle-aged adults.

GOOD TO REMEMBER

In a psychiatric setting, this combination should prevent diagnostic compartmentalisation: intellectual disability, compulsive symptoms or altered personality should not obscure systemic vascular and physical risk.

Intellectual disability occurs in almost all cases, and the behavioural phenotype can include obsessive-compulsive symptoms and personality disorders. These associations do not permit a psychiatrist to infer homocystinuria from an obsessive-compulsive presentation. They do, however, show why apparently ordinary phenomenology becomes diagnostically different when it appears with developmental and somatic clues.

Treatment may include pyridoxine, or vitamin B6, which increases homocysteine metabolism and reduces methionine and homocysteine. The Indian guidance also lists pyridoxine with vitamin B12 and folate among small-molecule approaches. Pyridoxine reduces these biochemical levels in approximately **50%** of patients. This is a treatment-response fact, not a licence for a psychiatrist to manage metabolic therapy independently.

A particularly useful negative finding is that folate plus vitamin B6 does not reduce stroke risk in adults with elevated homocysteine. The distinction matters because biochemical improvement and clinical vascular protection cannot simply be treated as interchangeable outcomes. It is the kind of nuance that strengthens an answer and, more importantly, prevents overconfident counselling.

10.6 Other named metabolic disorders: recognise the syndromic pattern

DISORDER	METABOLIC BASIS	CLUES THAT SHOULD REDIRECT THE FORMULATION	SPECIFIC THERAPEUTIC IMPLICATION
Galactosaemia	Autosomal recessive deficiency of one of three enzymes in galactose metabolism, including GALT	After milk introduction: failure to thrive, hepatosplenomegaly and cataracts	Galactose-free diet; if treated, intellectual disability is usually mild.
Lesch-Nyhan syndrome	X-linked mutation of the HGPRT gene causing a purine-metabolism defect with excessive uric acid production and excretion	Severe self-injurious behaviour with compulsive biting of fingers and lips, with spastic cerebral palsy and choreoathetoid movements	Behavioural risk management must be integrated with neurological and metabolic care.
Hurler syndrome (MPS I)	Autosomal recessive alpha-L-iduronidase deficiency	Storage disorder phenotype requires a multisystem formulation rather than an ID-only label.	Enzyme replacement for MPS and bone marrow transplantation for MPS I are listed interventions.

Galactosaemia teaches the importance of a feeding-linked systemic presentation. The triad of failure to thrive, hepatosplenomegaly and cataracts after milk introduction should redirect attention from an isolated developmental formulation toward galactose metabolism. The therapeutic link is equally direct: removing galactose is an intervention aimed at the pathway, not merely at a developmental consequence.

Lesch-Nyhan syndrome is especially memorable because the self-injury is severe and compulsive, rather than a generic behaviour problem. The child may be normal at birth before developing low-to-mild intellectual disability. The associated movement disorder and self-biting should prompt a syndrome-level formulation. In the consultation room, the practical priority is safety without losing explanatory precision: pain, communication difficulty, environmental response and the underlying neurobiological disorder can all shape the observed behaviour.

For the mucopolysaccharidoses, **Hunter syndrome (MPS II)** and Sanfilippo syndrome (MPS III) are both described as milder than Hurler syndrome (MPS I). The examination value is to recognise that MPS is a family of disorders rather than an eponym to be memorised in isolation. The clinical value is to retain the possibility that a developmental and behavioural presentation belongs to a multisystem storage disorder.

10.7 Treatability across metabolic disorders: use a mechanism-led therapeutic map

The key contrast across these conditions is not simply treated versus untreated. It is the kind of intervention that fits the disrupted biology. The Indian Psychiatric Society guidance identifies special or modified diets for many organic acidurias and aminoacidopathies, including PKU, glutaric aciduria type 1 and maple syrup urine disease. This should be read as an illustrative list, not as a claim that every metabolic disorder responds to dietary restriction.

- Replacement of deficient molecules includes thyroxine for hypothyroidism, enzyme replacement for mucopolysaccharidoses and copper histidine for Menkes disease.
- Small-molecule therapy includes creatine monohydrate for cerebral creatine deficiency syndromes and pyridoxine, vitamin B12 and folate for homocystinuria.
- Bone marrow transplantation is listed for alpha-mannosidosis and MPS I.
- Pharmacotherapy with vigabatrin and chelation of excess metals for Wilson disease and manganese transporter deficiency are further listed approaches.

These modalities are conceptually useful because they prevent therapeutic nihilism while also preventing indiscriminate treatment. A diet is appropriate when the disease mechanism supports dietary modification; replacement is appropriate when a missing molecule can be supplied; chelation addresses a harmful excess. The psychiatrist needs this map to understand why treatment is being offered and to recognise how psychiatric symptoms, adherence and family burden may affect it. Selecting the regimen, monitoring response and managing disease-specific risk remain specialist tasks.

When intellectual disability is linked to a metabolic pathway, the decisive question is whether recognition permits pathway-directed treatment before or alongside irreversible developmental consequences.

10.8 Psychiatric practice: formulate, do not overshadow

Metabolic diagnosis should sharpen psychiatric assessment, not end it. A person may require assessment for depression, anxiety, attentional difficulty, obsessive-compulsive symptoms, personality change, severe self-injury or behavioural distress. The diagnosis helps the clinician ask better questions about timing, cognitive baseline, neurological change, diet, treatment burden and family interpretation. It must not become a blanket explanation that erases a treatable psychiatric syndrome or a preventable safeguarding problem.

Conversely, a familiar psychiatric label should not absorb clues that point beyond psychiatry. In PKU, executive and attentional impairment can coexist with mood and anxiety disorders. In homocystinuria, a psychiatric presentation sits beside ocular, skeletal and vascular clues. In Lesch-Nyhan syndrome, self-injury needs immediate compassionate management, yet its form and accompanying neurological signs carry diagnostic information. The senior clinical move is to hold both truths at once: symptoms deserve direct care, and their biological context may alter risk, referral and prognosis.

For examination writing, organise the answer around the prototype, then demonstrate range. Begin with PKU - inheritance, enzyme, biochemical block, phenotype, newborn screening and dietary treatment. Use homocystinuria to show why systemic findings and psychiatric symptoms must be integrated. Add galactosaemia, Lesch-Nyhan syndrome and mucopolysaccharidoses as concise pattern-recognition examples, and finish with the principle of mechanism-matched intervention. That organisation conveys clinical understanding rather than memorised accumulation.

11 · Fetal alcohol spectrum disorder

Why this diagnosis earns its place in an intellectual-disability chapter. Fetal alcohol spectrum disorder (FASD) is the clinical umbrella for developmental consequences attributed to prenatal alcohol exposure. It matters because the developmental phenotype is often broader than the visible phenotype: a child may have substantial difficulty with learning, regulation and everyday functioning without displaying the full facial pattern of fetal alcohol syndrome (FAS). Conversely, the recognisable physical pattern should prompt a careful account of neurodevelopment, rather than being treated as a diagnosis made from appearance alone. FAS is **probably the most common nongenetic cause of intellectual disability**, but FASD must not be reduced to intellectual disability alone.




Fig 21.10 Fetal alcohol spectrum disorder is an umbrella for developmental consequences associated with prenatal alcohol exposure. Neurodevelopmental impairment may occur without the complete growth and facial pattern of fetal alcohol syndrome.

The useful clinical stance is therefore developmental and longitudinal. Ask whether prenatal exposure is documented, look deliberately for the phenotype, establish whether growth or brain-development indicators are present, and describe the functional profile in its own right. The label should bring order to a scattered presentation; it should not become a shortcut that explains every behaviour, nor should an absent facial phenotype be used to dismiss a credible exposure-related neurodevelopmental disorder.

11.1 Scope, language and diagnostic lineage

FASD is an umbrella, not a synonym for FAS. Its subdiagnoses include FAS, partial FAS (PFAS), alcohol-related neurodevelopmental disorder (ARND), alcohol-related birth defects (ARBD), and neurobehavioural disorder associated with prenatal alcohol exposure (ND-PAE). This terminology is clinically useful because it separates a full syndromic presentation from partial physical phenotypes and from neurodevelopmental presentations in which physical signs are not

demonstrable. ARBD is primarily a physical diagnosis and is not the focus of this neurodevelopmental discussion.

The original FAS criteria were set out by **Jones and Smith in 1973**. Contemporary clinical criteria are commonly described as the **Hoyme criteria**, updated from earlier Institute of Medicine criteria. For examination answers, the important point is not to recite a lineage of labels. It is to show that FASD diagnosis is organised around a coherent set of diagnostic domains: prenatal alcohol exposure, facial anomalies, growth deficiency, deficient brain growth or abnormal morphogenesis, and neurobehavioural impairment. These domains provide the framework for deciding which diagnosis best fits the available evidence.

■ **RULE Diagnostic principle.** In FASD, exposure history, physical findings and neurodevelopmental findings are interpreted together; none should be allowed to stand in for the others.

This framework also explains an apparently counterintuitive feature of the full syndrome. A diagnosis of **FAS** rests on the characteristic physical and neurobehavioural domain combination, so documentation of confirmed prenatal alcohol exposure is not required for that diagnosis. That is not permission to treat exposure history as unimportant. It reflects the specificity attributed to the combination when it is present, while recognising that reliable retrospective histories can be unavailable.

MNEMONIC

Use the spine, not a checklist blur

Think *exposure, face, growth, brain, behaviour*. It keeps the assessment ordered and makes it less likely that a clinician will mistake a single facial observation or a behavioural complaint for the whole diagnosis.

11.2 Biology and the syndromic phenotype

Alcohol is a developmental teratogen, so the clinical phenotype should be understood as the outcome of disruption during development rather than as a static cosmetic syndrome. A proposed mechanism is inhibition of **NMDA receptors** that mediate postsynaptic excitatory effects of glutamate, with effects on cell proliferation. This provides a biologically intelligible bridge between structural findings and later developmental difficulties. It does not, however, permit one to infer a particular cognitive profile from mechanism alone. Developmental outcomes are heterogeneous, and clinical formulation remains more informative than mechanistic shorthand.

Recognising the face requires discipline. The classic sentinel pattern is a **triad of short palpebral fissures, a smooth philtrum and a thin upper lip**. These are midfacial findings, not a general impression that a child “looks dysmorphic”. The examiner should observe and document each component carefully, because casual facial judgement is vulnerable to expectation, ancestry-related variation and poor-quality photographs. In the facial-anomaly domain, **at least two** characteristic features are required. A palpebral fissure length at or below the **10th centile**

qualifies as a facial feature; a thin vermilion border or smooth philtrum is assessed on a racially normed lip-philtrum guide at **rank 4 or 5**.

■ **TRAP Examination trap.** Do not diagnose FASD from a nonspecific statement such as “typical facies”. State the individual facial findings, then relate them to the wider diagnostic domains.

Growth and brain-development indicators add clinical weight when present. Growth deficiency is represented by height or weight at or below the 10th centile. Deficient brain growth or abnormal morphogenesis may be reflected by head circumference at or below the 10th centile, structural brain abnormalities, or recurrent nonfebrile seizures. These are diagnostic indicators, not interchangeable measures of the same process. A growth measurement answers a different question from head growth, and neither replaces a careful account of neurobehavioural functioning.

DIAGNOSTIC DOMAIN	WHAT THE CLINICIAN ESTABLISHES	WHY IT CHANGES REASONING
Prenatal alcohol exposure	A reliable history when it is available.	It is essential for some FASD diagnoses and strengthens causal attribution.
Facial anomalies	The sentinel midfacial pattern, measured rather than guessed.	It distinguishes a recognisable physical phenotype from a purely behavioural presentation.
Growth deficiency	Appropriate anthropometric evidence.	It contributes a separate developmental domain.
Brain growth or morphogenesis	Head-growth, structural or seizure indicators.	It links the physical assessment to neurodevelopmental risk.
Neurobehavioural impairment	Impairment on a global estimate or a relevant neurobehavioural or behavioural domain.	It prevents the diagnosis becoming a morphology exercise.

The threshold for neurobehavioural impairment in this clinical framework is performance at least 1.5 SD below the mean on a global cognitive estimate or on a single relevant neurobehavioural or behavioural domain. The key clinical inference is modest but important: a child can require a detailed functional evaluation even when the physical findings are incomplete. The physical phenotype and the developmental phenotype overlap, but they are not identical.

11.3 Distinguishing FAS, partial FAS and ARND

The diagnostic categories are best used as descriptions of the configuration before you, not as a hierarchy of “real” and “less real” disorders. FAS is the full syndromic pattern described above. PFAS retains facial anomalies and neurobehavioural impairment in the presence of confirmed exposure. When exposure is not confirmed, PFAS requires those findings together with impairment in either growth or brain growth. This additional requirement is logically necessary: without a documented exposure history, the clinician needs a more specific configuration before making an exposure-attributed diagnosis.

Alcohol-related neurodevelopmental disorder is made in the absence of the physical indicators of face and growth, but requires confirmed prenatal alcohol exposure and neurobehavioural impairment. This category is especially important because it captures children whose most disabling difficulties are cognitive, behavioural and functional rather than dysmorphic. Its existence protects against a common diagnostic error: assuming that the absence of the FAS face excludes alcohol-related neurodevelopmental harm.

- For FAS, look for the characteristic syndromic combination and do not make documented exposure a false prerequisite.
- For PFAS, decide first whether prenatal exposure is confirmed, because that changes the required domain configuration.
- For ARND, do not demand facial or growth indicators when a documented exposure history and neurobehavioural impairment are present.

These distinctions should improve the quality of documentation. Record what was observed, what was measured, which part of the exposure history is corroborated and which diagnostic domain is not established. That language is more clinically defensible than forcing a label from an incomplete record. It also keeps diagnostic uncertainty visible, which is preferable to either overcalling a syndrome or making a false reassurance from an absent physical feature.

11.4 ND-PAE: the proposed neurodevelopmental diagnosis

Neurobehavioural disorder associated with prenatal alcohol exposure is placed in the DSM appendix under Conditions for Further Study, with proposed rather than full diagnostic criteria. This status matters in a written answer: ND-PAE is a structured clinical construct, but it is not a full DSM diagnosis. It should therefore be described accurately rather than presented as though it had identical nosological standing to every routine DSM diagnosis.

Its exposure criterion specifies more-than-minimal gestational exposure as more than 13 drinks per month during pregnancy, or more than 2 drinks on any one occasion. This threshold is a criterion for the proposed diagnosis, not a declaration that exposure below it is safe. There is **no established safe threshold** or safe period of gestational alcohol exposure. The effects of exposure remain difficult to reduce to a single number because developmental stage and maternal and fetal factors modify risk.

The ND-PAE structure makes the clinical interview more precise. It requires impairment in at least one neurocognitive domain, at least one self-regulation domain, and at least two adaptive-functioning domains, one being communication or social communication and interaction. The neurocognitive domains include global intellectual performance, executive functioning, learning, memory and visual-spatial reasoning. The self-regulation domains include mood or behavioural regulation, attention and impulse control. Adaptive difficulties can involve communication, social communication and interaction, daily living skills or motor skills.

Global intellectual performance may be demonstrated by an IQ or comprehensive developmental standard score at or below 70. That is one route within the neurocognitive criterion, not the

definition of the whole condition. A sound assessment asks which abilities fail in ordinary demands and whether the pattern coheres with the exposure history. Neurocognitive deficits, regulation problems and adaptive limitations are clinically separable, though they often compound one another in a child's life.

Diagnosis should be deferred at or below 3 years, because reliable neurocognitive assessment is limited at that stage. Deferral is not abandonment. It means recording exposure and developmental concerns, following the child's profile over time, and avoiding a prematurely fixed label. ND-PAE can be diagnosed in alcohol-exposed children with or without FAS, including those without facial dysmorphism. That independence is precisely why the clinician must inspect the neurodevelopmental phenotype rather than use the face as a gatekeeper.

11.5 Cognitive and behavioural clinical correlation

FASD creates a frequent conceptual trap in intellectual-disability teaching. The presence of FASD does not mean that every affected child has intellectual disability, and a normal-looking child should not be assumed to have intact higher functions. The majority of children with FASD do not have IQ below 70; observed ability spans from severe intellectual impairment to average performance. In children with FAS, the average full-scale IQ is around 72, compared with around 80 in children with heavy prenatal alcohol exposure without FAS. Intellectual disability in FAS is usually mild to moderate, though it may be severe.

Interpretation of intelligence testing itself belongs in the intellectual-disability assessment material.

GOOD TO REMEMBER

Do not let a single global score decide whether FASD is clinically important. A child with broadly higher measured ability may still struggle with executive demands, new learning, memory, visual-spatial reasoning, attention, impulse control or social communication. Translate these findings into difficulties at school, with home routines and in judgement; do not misread them as wilful noncompliance.

Psychiatric comorbidity adds a second layer of complexity. **Attention-deficit/hyperactivity disorder** is the most common co-occurring psychiatric diagnosis, with reported prevalence among children with FASD between 44% and 96%. Mental health problems have been identified in more than 90% of people with histories of significant prenatal alcohol exposure. These figures should not invite indiscriminate diagnostic labelling. They should prompt the clinician to ask whether attention, impulsivity and emotional dysregulation are being described carefully, rather than folded uncritically into a single explanatory story.

The absence of the classic face does not exclude an alcohol-related neurodevelopmental disorder; assess exposure history and functional phenotype together.

11.6 Differential diagnosis and disciplined attribution

Attribution is a clinical conclusion, not a moral judgement. Prenatal alcohol exposure may be difficult to establish and the behavioural presentation is not pathognomonic. The diagnostic task is therefore one of pattern recognition followed by active exclusion of alternatives. A credible history is valuable, but history alone cannot establish that every developmental difficulty is caused by alcohol. Equally, uncertainty in history should not lead to a careless assumption that a distinctive physical and neurobehavioural pattern is irrelevant.

The named differential includes genetic conditions such as Williams syndrome, Down syndrome and Cornelia de Lange syndrome, and other teratogenic conditions such as fetal hydantoin syndrome and maternal phenylketonuria. These conditions can share physical and behavioural characteristics with ND-PAE. The practical lesson is to seek positive evidence for the proposed diagnosis and contradictory evidence for alternatives. “Alcohol exposure reported” is not a substitute for a differential diagnosis; “facial features absent” is not a substitute for one either.

■ **TRAP Examination trap.** A documented exposure history does not remove the need for differential diagnosis, and an undocumented history does not automatically rule out FAS when the required syndromic configuration is present.

- Separate observations from inferences: document anthropometry, facial examination, neurodevelopmental findings and exposure evidence distinctly.
- Ask whether each finding belongs to the FASD configuration or whether it suggests an alternative developmental or teratogenic condition.
- State the limits of the available evidence rather than converting an uncertain history into false certainty.

For a broader developmental formulation, use the relevant sections of the *Child psychiatry: assessment, treatment, services and protection notes*. Assessment of adaptive behaviour, intellectual testing and the multidisciplinary management of intellectual disability are deliberately treated elsewhere in these notes. Here, the point is narrower: recognise when alcohol-related developmental injury is a plausible organising diagnosis and describe why.

11.7 Epidemiology, prevention boundary and clinical significance

Reported prevalence varies greatly with population, case definition and completeness of assessment, so a single headline figure can mislead. The mean global general-population prevalence of FASD reported for 2012 was 7.7 per 1,000. The corresponding mean for the Americas was 8.8 per 1,000. In the United States, FASD has been estimated at between 1.1% and 5.0%, whereas the reported estimate for FAS itself is between 2 and 7 per 1,000 children. These are not competing answers. They measure differently framed populations and, crucially, the spectrum is wider than the full syndrome.

7.7 per 1,000 FASD: GLOBAL MEAN	8.8 per 1,000 FASD: AMERICAS MEAN	1.1% to 5.0% FASD: UNITED STATES ESTIMATE	251.5 per 1,000 ND-PAE: CHILDREN IN CARE SETTINGS
---	---	--	--

Ascertainment is not a technical footnote. ND-PAE estimates in fully evaluated children have ranged between 31.1 and 98.5 per 1,000, and a meta-analytic estimate among children in care settings was 251.5 per 1,000. The latter should never be generalised to all children. It does, however, underline why a high-risk care context should lower the threshold for a thoughtful developmental history and a careful search for unmet needs.

The clinical preventive message is uncomplicated: there is no established safe threshold or safe period for alcohol exposure in pregnancy. The public-health framework for prevention is addressed in the *Prevention of intellectual disability* material. In this section, the practical diagnostic corollary is that clinicians should discuss exposure without blame. Shame drives concealment, while accurate histories and respectful care improve recognition of children who need a fuller developmental evaluation.

12 · Tuberous sclerosis, neurofibromatosis and neurocutaneous syndromes in ID

Why these disorders matter in intellectual disability. Neurocutaneous syndromes make the aetiological assessment of intellectual disability clinically visible. The examination of skin, eyes and nervous system can connect a developmental presentation to a multisystem disorder, while the psychiatric formulation must remain alert to seizures, cognitive change, behaviour, family burden and the practical implications of a long-term condition. They reward disciplined observation: a lesion on the skin is not a diagnosis by itself, but it may be the clue that makes the whole developmental history intelligible.

These syndromes are especially useful for teaching this approach. **Tuberous sclerosis complex** brings developmental, epileptic and systemic manifestations into a formulation.

Neurofibromatosis type 1 likewise has cutaneous signs that may point to neurodevelopmental and neurological needs. The task for the psychiatrist is neither to reduce the patient to a syndrome nor to overlook the syndrome while treating only behaviour.

1 in 7000 TSC POPULATION PREVALENCE	about 70% TSC DE NOVO MUTATIONS	above 90% TSC SEIZURES AT PRESENTATION	50% NF1 NEW CASES IN UNAFFECTED FAMILIES
--	---	---	---

12.1 Neurocutaneous syndromes as a clinical frame

A neurocutaneous syndrome should be approached as a pattern across organ systems rather than as a dermatological label. In a person with intellectual disability, the clinician asks whether physical stigmata, epilepsy, cognition and change in functioning can be explained by a shared condition. This is useful because the consequences of an established syndrome are not merely diagnostic. It changes the threshold for collateral history, medical liaison and surveillance for new symptoms.

For tuberous sclerosis, older texts may use **epiloia** or tuberoses sclerosis. The classical teaching spine comprises **three** linked manifestations: epilepsy, intellectual disability and facial angiofibromas on the malar surface. It remains a memorable clinical entry point, but it is not a substitute for examining the person or for recognising that features evolve across development.

In particular, facial angiofibromas may first become apparent in adolescence; their absence earlier in life should not falsely reassure the clinician.

MNEMONIC

E-I-F for the classical TSC spine

epilepsy, intellectual disability and facial angiofibromas. Use it to prompt a whole-person review, not as a rule-out checklist.

- **RULE** When intellectual disability coexists with characteristic skin or neurological signs, formulate the syndrome as a multisystem condition and reassess for changing neurological function.

Physical signs are most useful when they sharpen the examination rather than when they create diagnostic tunnel vision. In tuberous sclerosis, look deliberately for facial angiofibromas, hypomelanotic macules including ash-leaf spots, shagreen patches over the lower trunk or buttocks, ungual fibromas and retinal phakoma. These findings support a coherent clinical hypothesis, yet the psychiatrist should also document what they mean functionally: distress about appearance, stigma, pain, caregiving burden and barriers to education or work can all influence the presenting complaint.

12.2 Tuberous sclerosis complex: genetics and mechanism

Tuberous sclerosis is inherited in an **autosomal dominant** pattern, although many presentations are sporadic. What varies between affected people, and even within one family, is the **expressivity** of the disorder rather than its penetrance: current genetic sources put penetrance at essentially complete, so a relative who carries the pathogenic variant is affected, but the severity and organ pattern of that involvement differ widely. Older texts, including some in common examination use, describe this as variable penetrance. Say expressivity, and know why the distinction matters: it means a mildly affected parent is affected, not a non-penetrant carrier, and their apparently trivial skin sign is a reason to examine them properly. This matters when taking a family history: the history may be apparently negative without making a genetic disorder implausible. In conventional testing, an alteration in TSC1 or TSC2 is identified in 80% to 90% of people with the condition. The choice and sequence of tests belong to the ID aetiological work-up; the syndrome-specific result is interpreted alongside phenotype and family context rather than in isolation.

TSC1, located at 9q34, encodes hamartin. **TSC2**, located at 16p13, encodes tuberin. Their products act as tumour suppressors that restrain Rheb-mediated mTOR kinase activity. This molecular account is clinically valuable because it explains why mTOR-directed treatment has a rational place in selected manifestations; it does not imply that every cognitive or behavioural difficulty is directly reversible by treating a lesion.

The broad developmental lesson is that a molecular diagnosis describes risk and mechanism, not destiny. Cognitive ability in tuberous sclerosis has been reported as bimodally distributed, with a

group having normal IQ and another in the profoundly affected range. Such heterogeneity makes an individual developmental and adaptive assessment indispensable. It also prevents the opposite errors of assuming that every person with the syndrome has intellectual disability, or overlooking substantial support needs in someone whose physical signs dominate the consultation.

GENETIC OR BIOLOGICAL LEVEL	CLINICAL MEANING	PSYCHIATRIC IMPLICATION
TSC1 and TSC2 encode hamartin and tuberin	The disorder has a defined tumour-suppressor pathway	Family history and physical findings should be integrated with the syndrome phenotype
The complex restrains mTOR activity	Selected structural manifestations have a pathway-based treatment rationale	Do not equate lesion-directed treatment with a complete explanation or cure for developmental needs
Variable penetrance and sporadic presentation	A negative family history does not settle the question	Give counselling within the wider genetic-work-up framework and avoid premature certainty

12.3 TSC phenotype: seeing the syndrome without missing the person

The skin examination is often the most accessible part of recognition. **Facial angiofibromas** are classically malar, while hypomelanotic macules, shagreen patches and ungual fibromas provide other cutaneous clues. Describe what is actually seen rather than using a label as shorthand. A careful record helps later clinicians distinguish longstanding stigmata from new change and allows the family to understand why a psychiatric consultation involves a full physical review.

Central nervous system lesions account for much of the developmental and neurological burden. Subependymal tubers may line the ventricles. Their reported size is **1 to 3 cm**, but the clinical question is not their size alone: one must ask whether a growing lesion could be producing seizures, headache, altered alertness, loss of skills or progressive cognitive impairment. **Subependymal giant cell astrocytomas** may compress and irritate adjacent cortex, and obstruction at the foramen of Monro can cause obstructive hydrocephalus. New deterioration should therefore trigger medical rather than purely behavioural attribution.

■ **TRAP** Do not call a new behavioural change “part of the intellectual disability” before considering seizure activity, lesion-related change or hydrocephalus in a person with TSC.

Systemic review is equally important. Tuberous sclerosis can involve renal cysts and angiomyolipomas, and cardiac rhabdomyomas that may cause dysrhythmias. These are not incidental facts for the psychiatrist. Pain, sleep disruption, repeated admissions, frightening cardiac symptoms and treatment demands can present indirectly as irritability, avoidance, low mood or family conflict. A liaison formulation distinguishes a psychiatric syndrome from a psychological response to a medical burden, while allowing both to be present.

Liaison also requires intellectual humility. Each specialty may see a valid but partial account of the person: the lesion, the seizure, the renal issue, the visible skin finding, the behaviour, or the

family's exhaustion. The psychiatric assessment can integrate these accounts around daily functioning and distress. It should make uncertainty explicit, identify the next clinically answerable question and avoid implying that a behavioural description can settle a neurological concern.

For candidates, the practical inference is simple: findings in brain, skin, heart, eye and kidney make the syndrome multisystem. The specialist medical team directs organ surveillance. Psychiatry contributes longitudinal developmental description, a baseline of behaviour and cognition, assessment of emotional symptoms at the person's communication level, and support for carers faced with uncertainty.

12.4 Epilepsy, intellectual disability and behaviour in TSC

Epilepsy is central to the neuropsychiatric phenotype of tuberous sclerosis. Partial motor and complex partial seizures are particularly characteristic. Other seizure forms may occur, with classic absence seizures a probable exception; severe presentations may begin with infantile spasms. This seizure history should be taken with attention to onset, semiology, treatment response, injuries, sleep, postictal behaviour and apparent developmental plateau or loss. A family's account of "spells" may be much more informative than a single label in an old record.

Vigabatrin is particularly effective for infantile spasms in tuberous sclerosis. A randomised, single-masked trial in young children found the subgroup with TSC-related spasms particularly responsive. This is a responsiveness observation, not a licence to treat without specialist epilepsy oversight.

IN INDIA

The Indian psychiatric guideline also lists tuberous sclerosis among its relevant uses.

The limitation is clinically serious. Vigabatrin can cause progressive, permanent bilateral vision loss, including concentric visual field constriction. It has also been associated with behavioural disturbance or ADHD symptoms, depression and psychosis, with no beneficial psychotropic effect reported.

GOOD TO REMEMBER

When behaviour worsens after an antiseizure treatment change, the formulation must consider medication effect as well as seizure control, environment and the person's baseline vulnerabilities.

- Obtain a careful seizure and medication chronology before attributing agitation or decline to a primary psychiatric disorder.
- Ask carers for concrete examples of change in sleep, attention, mood, behaviour and daily functioning.
- Escalate new focal neurological symptoms or progressive cognitive change to the relevant medical team.

There are useful risk associations, but they must not become deterministic predictions. TSC2 mutations are associated with more severe epilepsy and cognitive impairment; moderate to severe intellectual disability is more common in people with infantile spasms. These associations support vigilance and early support planning; they do not replace an assessment of the individual's communication, strengths, behaviour or family resources.

TSC also has a limited reverse relationship with autism at population level: **less than 4%** of all autism is attributable to TSC. The point is diagnostic humility. TSC may be relevant to autism in an individual with the syndrome, but it does not explain autism generally, and autism assessment should still be conducted on its own clinical evidence.

12.5 TSC management: targeted treatment and longitudinal formulation

The psychiatric role in a multisystem syndrome is coordinative and longitudinal. First, identify the current problem in concrete terms: seizure-related risk, a change in cognition, behavioural disturbance, emotional symptoms, school or vocational difficulty, caregiver strain, or difficulty navigating multiple specialties. Next, establish the baseline against which change can be judged. Finally, communicate concerns in a form useful to neurology, paediatrics, dermatology, ophthalmology, cardiology and renal services as appropriate.

For subependymal giant cell astrocytomas, the mTOR inhibitors everolimus and sirolimus can shrink the lesion. Topically applied mTOR inhibitors can reduce facial angiofibromas. These examples clarify an important distinction: a medicine can be directed at a structural or skin manifestation, while the psychiatrist still addresses developmental support, distress, adherence, family understanding and the lived effects of a chronic disorder.

In TSC, a change in behaviour or cognition is a clinical event to explain, not a baseline feature to dismiss.

Generic genetic counselling principles and the sequence of laboratory investigation are addressed in the ID aetiological work-up. In this syndrome-specific setting, the clinician should explain that inheritance may be variable and that sporadic presentation is common, then ensure that the family receives appropriately coordinated counselling. Avoid giving a falsely precise recurrence discussion from a psychiatric interview alone, especially when the molecular result and the family phenotype have not yet been clarified.

12.6 Neurofibromatosis type 1: recognition and diagnostic reasoning

NF1 is associated with the tumour-suppressor gene product **neurofibromin**, at 17q11.2. Neurofibromin is involved in regulation of cell division and tumour suppression. As in TSC, the psychiatric significance lies in recognising a medical context for developmental and behavioural needs while retaining ordinary clinical discipline: a person can have NF1 and an independent mood, anxiety, psychotic or behavioural disorder.

Diagnosis usually requires **at least two** diagnostic features from the established list. The recognised features span skin lesions, peripheral tumours, regional freckling, ocular findings, optic-nerve involvement, bony lesions and an affected first-degree relative. The list is best used as

a structured prompt for referral and documentation, not as an invitation for the psychiatrist to make a final diagnosis without appropriate examination.

NF1 FINDING DOMAIN	FEATURES TO LOOK FOR OR ELICIT	WHY IT MATTERS IN AN ID ASSESSMENT
Skin	Café-au-lait macules , axillary or inguinal freckling	May supply the first clue to a syndromic formulation
Peripheral tumours	Neurofibromas, including plexiform neurofibroma	Symptoms and visible difference may affect distress, self-image and help-seeking
Eye, bone and family history	Lisch nodules, optic nerve glioma, bony lesions, affected first-degree relative	Signals the need for appropriately coordinated specialist assessment

The café-au-lait threshold is a common recall point but should be applied with precision. The cited criterion specifies **six or more** lesions, larger than 5 mm before puberty or 15 mm after puberty. An alternative source frames the sign as more than six lesions, each larger than 5 mm in children and 1.5 cm in adults, and notes that individual café-au-lait spots occur in normal individuals. Therefore, a single lesion should not be over-interpreted; the pattern, associated signs and clinical context matter.

■ TRAP

Do not diagnose NF1 from an isolated café-au-lait spot, and do not mistake a referral criterion for a psychiatric explanation.

12.7 A psychiatrist’s practical synthesis

When a neurocutaneous syndrome is known or suspected in a person with intellectual disability, begin by separating the immediate clinical question from the aetiological label. Is the consultation about new seizures, progressive cognitive difficulty, aggression, emotional distress, medication adverse effects, family conflict or a support failure? That distinction prevents a syndrome name from becoming a catch-all explanation and makes the response clinically useful.

A good formulation has parallel layers. The biological layer records the established or suspected syndrome, current neurological concerns and treatment exposure. The developmental layer records the person’s baseline abilities, communication and pattern of support. The psychological layer asks how the person understands symptoms, procedures, visible differences and restrictions. The social layer identifies who observes change, who carries care and what makes access to coordinated treatment difficult. Writing these layers separately helps the team avoid both diagnostic overshadowing and fragmented, organ-by-organ care.

Communication with families deserves the same precision as communication with specialists. Explain why a new symptom is being taken seriously, what the immediate safety concern is, and which clinician will address it. Invite carers to retain their observations rather than dismissing them as vague. In people with limited spoken language, an account of altered routine, sleep, facial expression, mobility, appetite, engagement or self-care may be the earliest available evidence of a


medical or emotional change. The psychiatrist's value is often to translate this observation into an actionable question for the wider team.

- Examine and document relevant skin signs, while asking what they mean to the patient and family.
- Seek a longitudinal account of development, seizures, cognition, behaviour and treatment exposure.
- Compare the present state with the person's own baseline rather than with an abstract stereotype of the syndrome.

These disorders are a reminder that psychiatry in intellectual disability is whole-person medicine. The syndrome gives a framework for anticipation; it does not erase the individual's personality, preferences, communication style, social context or capacity for psychological distress. The best answer in the examination, and the best practice in the clinic, links a recognisable phenotype to a thoughtful developmental and psychiatric formulation without claiming more certainty than the evidence supports.

13 · Behavioural phenotypes of genetic syndromes

The useful question is not “what behaviour does this syndrome cause?” A behavioural phenotype is a clinically recognisable pattern that can sharpen observation, guide anticipatory enquiry and prevent diagnostic overshadowing. It is not a substitute for seeing the person in front of us. In intellectual disability, knowing the aetiology may provide **clues to psychiatric comorbidity**, because some syndromes are repeatedly associated with particular behavioural patterns. The practical gain is disciplined curiosity: ask about the pattern that is plausible, then establish whether it is actually present, distressing and functionally important.



- Probabilistic association
- Vulnerabilities
- Not destiny
- Strengths
- Anticipatory enquiry

Fig 21.11 A behavioural phenotype is a raised probability of a pattern, not a behaviour caused in every person by a syndrome. It guides anticipatory enquiry while preserving individual strengths, context and diagnostic assessment.

The term **behavioural phenotype** was introduced by Nyhan. Earlier clinical observers such as Penrose had noted a distinctive social presentation in Down syndrome, but the modern concept is broader and more cautious. Its working definition is associated with Dykens: a syndrome is associated with a raised likelihood of a behavioural or developmental outcome when compared with people without that syndrome. That is a **probabilistic association**, not a diagnostic label attached to every individual.

13.1 What the concept does - and does not - mean

A behavioural phenotype is best understood as a pattern of vulnerabilities, strengths and recurrent clinical problems that occurs more often in a defined condition than outside it. This distinguishes a useful phenotype from an attractive anecdote. The clinician is looking for the **combination of features and its association with a condition**, rather than claiming that an isolated common symptom identifies a syndrome. Attentional difficulty, social anxiety, sleep

disruption, stereotypy and self-injury all occur across diagnostic groups. Their meaning comes from their form, timing, associated features and clinical setting.

-
- **RULE** A behavioural phenotype changes prior probability; it does not determine the individual's behaviour.
-

The concept explicitly rejects genetic determinism. Behavioural expression is shaped by developmental level, communication, physical illness, sensory experience, family responses, opportunity and the person's own temperament. A syndrome may create a particular vulnerability while the environment influences how often it is triggered, how it is understood and how much disability follows. This is why a behavioural phenotype should lead to more careful assessment, not to premature explanation.

Equally, no behavioural phenotype is invariant. The absence of a familiar feature should not be used to dismiss a molecular or clinical diagnosis; its presence should not be used to make one. This is especially important in adults, in people with co-occurring sensory or neurological problems, and where developmental history is incomplete. The phenotype is a clinical map, not a gatekeeping criterion.

Specificity lies on a spectrum. A broad feature may be shared across several syndromes, whereas a particular form of that behaviour may be unusually informative. For example, a particular form of stereotyped hand wringing is almost invariably seen in Rett syndrome, even though hand stereotypies occur in many clinical contexts. The distinction protects against the common error of treating the category "stereotypy" as though it carried the specificity of its most characteristic form.

-
- **TRAP** Do not turn a phenotype into a personality stereotype. A recurring association may help formulate a question, but it never excuses failure to assess pain, communication, mood, safeguarding, environmental demands or co-occurring psychiatric illness.
-

13.2 How to use a phenotype in assessment

Use the syndrome to improve the interview, not to abbreviate it. Start with a baseline account of communication, adaptive functioning and the person's usual temperament. Then ask about the behaviours linked to the syndrome in a neutral, concrete way: their form, onset, antecedents, consequences and effect on family life, education, work and health. A behavioural signature has most value when it prompts a question that might otherwise not have been asked.

When a behaviour is established, the next task is formulation. Food-seeking, self-injury, withdrawal, agitation or sleep disruption can be part of a phenotype, but can also signal an untreated medical problem, a mismatch between demands and ability, a communication failure or a psychiatric disorder. The syndrome changes the differential weighting; it does not make competing explanations disappear. The assessment of psychiatric comorbidity in intellectual disability is addressed in the chapter's discussion of dual diagnosis, and the general assessment framework belongs in the Child psychiatry: assessment, treatment, services and protection notes.

- Describe the behaviour precisely before naming it as syndrome-related.
- Establish what has changed from the person’s own baseline and what has changed around them.
- Look for the linked behavioural cluster, not an isolated headline symptom.
- Use the formulation to anticipate risk and support needs without attributing blame.

This approach matters to families. A **strong behavioural phenotype** describes a vulnerability that may be evident even in a supportive environment. In such situations, parents and teachers who have worked hard should not be invited to conclude that the behaviour proves a failure of care. The right message is more exact: environment still matters greatly for safety, distress and functioning, but it may not fully remove a syndrome-linked vulnerability.

The historical literature illustrates why this caution is necessary. Reviews in the 1970s discussed only **five** chromosomal syndromes in this context. The concept has since become clinically richer, but a larger catalogue does not make its inferences categorical. The contemporary task is to be specific enough to notice a pattern and humble enough not to overread it.

13.3 Cross-syndrome signatures: a working comparison

The table is a pattern-recognition aid, not a diagnostic algorithm. It deliberately pairs the most discriminating feature with the clinical caution that prevents overclaiming. Several entries contain symptoms that are common in many people with intellectual disability; their value comes from the surrounding phenotype.

SYNDROME	USEFUL BEHAVIOURAL OR COGNITIVE SIGNATURE	CLINICAL READING
Down syndrome	Relative strength in social communication, with a wide variety of behaviour.	Do not replace assessment with the old “very sociable” personality stereotype.
Fragile X syndrome	Gaze aversion, shyness, social and performance anxiety, social withdrawal and hyperarousal, often with attention and activity difficulties.	Social avoidance can coexist with interest in others and must not be equated automatically with autism.
Prader-Willi syndrome	Hyperphagia with food seeking, foraging or hoarding, and life-long food preoccupation.	Food access and supervision are central environmental issues, while the wider formulation must also consider distress and routine.
Angelman syndrome	Episodic paroxysmal or inappropriate laughter with absence of speech.	The apparently happy expression should not conceal major communication needs.
Williams syndrome	High, sometimes indiscriminate sociability with reduced social flexibility.	Warm approach behaviour can coexist with poor peer relationships, anxiety and vulnerability in unfamiliar social settings.

SYNDROME	USEFUL BEHAVIOURAL OR COGNITIVE SIGNATURE	CLINICAL READING
Smith-Magenis syndrome	Severe self-injury, disruptive or aggressive outbursts, tantrums and hyperactivity.	Ask specifically about sleep, because disturbed sleep can amplify daytime behavioural difficulty.
Cri-du-chat syndrome	High-pitched cat-like cry in infancy.	This is an early developmental clue; later assessment should address language, attention and activity rather than treating the cry as a complete phenotype.
Lesch-Nyhan syndrome	Severe self-injury, typically compulsive biting of fingers and lips.	Its form and apparent compulsion make this a high-priority safety phenotype.
22q deletion syndrome	Childhood ADHD, autistic features and mood swings, with an important later psychosis vulnerability.	Developmental and psychiatric surveillance must extend across developmental transitions.
Rett syndrome	Loss of purposeful hand skills, replaced by stereotyped hand wringing and clapping.	The course and the exact hand phenomenology matter more than the label “stereotypy”.
Cornelia de Lange syndrome	Self-injury, autistic features and pleasurable responses to vestibular stimulation.	Communication may be disproportionately impaired relative to adaptive function, which should shape the formulation.

Attentional problems demonstrate the point about shared symptoms: they are frequent in fragile X, Williams and cri-du-chat syndromes. They are therefore clinically important but not syndrome-specific. By contrast, relatively characteristic forms include the unusual hand behaviour in Rett syndrome and extreme hyperphagia in Prader-Willi syndrome. An examination answer is stronger when it says whether a feature is shared, relatively specific or simply one component of a wider pattern.

13.4 The phenotype is often richer than its headline feature

The classic “Down personality” stereotype is difficult to verify. A safer clinical account is that Down syndrome has a relative social-communication strength and a broad behavioural range. It is also associated with less frequent and less severe psychopathology than in other groups with intellectual disability, while still requiring ordinary clinical vigilance. In later life, the phenotype includes high rates of dementia or Alzheimer disease in middle-aged to late adulthood. The teaching point is not that people with Down syndrome are protected from psychiatric problems; it is that group-level tendencies should never flatten individual presentation.

Fragile X syndrome is often recognised through its anxious, avoidant social style. In full-mutation males, the tabled phenotype also includes anxiety, perseverative language, an ADHD-like presentation, aggression, stereotypies and ASD-like behaviours. Yet men with fragile X syndrome are usually affectionate and lack the aloof quality typical of autism, despite social impairment.

This is a useful discriminator, not a rule that excludes autism. It directs the clinician to observe the quality of social engagement rather than inferring it from gaze aversion alone.

The food phenotype of Prader-Willi syndrome extends beyond hunger. Skin-picking, hoarding, concerns with symmetry, exactness, ordering, cleanliness and sameness, tantrums, aggression, stubbornness, underactivity and emotional lability may form part of the behavioural picture. Sleep also deserves active enquiry: excessive daytime sleepiness with sleep-architecture abnormalities can contribute to apparent inattention, irritability or reduced participation. The assessment should separate an environmental response to restricted food access from a mood, anxiety or behavioural disorder, while recognising that these may coexist.

In Angelman syndrome, laughter is only the opening clue. Other observed features include overactivity with short attention span, enjoyment of social and physical contact, mouthing objects and fascination with water. The older label “happy puppet” arose from a particular combination of nonverbal presentation, ataxia, frequent laughter and flexed arms. It may help recall the history, but it is not respectful shorthand for the person’s needs. The clinically relevant point is severe to profound intellectual disability with absence of speech, requiring an active search for nonverbal communication and distress signals.

Williams syndrome is another corrective to superficial impressions. It may show loquaciousness with “cocktail party manners” and a relative strength in language, auditory processing and music alongside poor visuospatial performance. However, friendliness may coexist with anxiety, including specific phobia and generalized anxiety. Treating verbal fluency as evidence of social comprehension is the trap. The person may initiate readily yet struggle with reciprocal flexibility, boundaries or sustained friendships.

13.5 Self-injury, stereotypy and sleep: read the mechanism before the behaviour

Self-injury is a description, not an explanation. Its diagnostic and safety significance depends on form, function and context. Lesch-Nyhan syndrome illustrates a highly characteristic pattern: stereotypic dystonic movements and self-mutilation of fingers, lip biting and other self-injury occur unless the individual is restrained. This is not interchangeable with all self-injury in neurodevelopmental conditions. It should prompt urgent attention to tissue protection, pain, oral injury, family burden and specialist coordination.

In Rett and Cornelia de Lange syndromes, self-injury may arise secondarily from hand-to-mouth stereotypies. The mechanism matters because it changes the clinical questions: is the injury a direct repetitive motor act, a response to distress, a communication of pain, or an outcome of a hand-to-mouth movement? The same visible lesion may therefore require very different prevention plans.

Contrasting source lists make a useful methodological warning. The DSM-5-TR names Lesch-Nyhan, Rett, fragile X, Cornelia de Lange and Smith-Magenis syndromes as examples of neurogenetic disorders with stereotypies. Another source instead names Cornelia de Lange, fragile X, Smith-Magenis, cri-du-chat and Rett syndromes in relation to stereotypic movements and self-injurious behaviour. Neither list should be treated as an exclusion list. They illustrate

overlapping clinical territory, while the detailed phenotype supplies the discriminating information.

Smith-Magenis syndrome is especially instructive because its phenotype links daytime behaviour and sleep. Alongside self-injury and outbursts, a **self-hug** can be a useful signature gesture, and the sleep problem may include absence of REM sleep. The reported disruption of melatonergic control of the sleep cycle provides a plausible bridge between sleep management and behavioural or cognitive functioning. This is not an invitation to prescribe from a syndrome name; it is a reminder that sleep assessment may be central rather than incidental.

MNEMONIC

PATTERN before label

Precise form, Antecedents, Trajectory, Triggering pain or sleep problem, Effect on function, Related syndrome features, Needs and safety.

13.6 Developmental trajectories and later psychiatric risk

Some phenotypes are defined as much by change over time as by a single behaviour. In Rett syndrome, loss of hand skills occurs with language regression, communication deficits, social withdrawal, hyperventilation or breath-holding, bruxism, sleep irregularities and some autistic symptoms. The early withdrawal should not be read as a fixed autism phenotype: socialization may later improve, with intense eye gaze in many patients. Rett syndrome itself is considered in the regression discussion elsewhere in this chapter; here, its course demonstrates why a behavioural phenotype must be longitudinal.

For 22q deletion syndrome, the crucial temporal pattern is a childhood developmental and behavioural presentation followed by risk of psychosis in adolescence or adulthood. About **25%** develop schizophreniform psychosis. At the same time, no other single risk factor is so strongly associated with psychosis. These findings belong together: striking association does not imply universal penetrance.

nearly 100% OBESITY RISK IN PRADER-WILLI SYNDROME WITHOUT AGGRESSIVE DIETARY INTERVENTION	~25% SCHIZOPHRENIFORM PSYCHOSIS IN 22Q DELETION SYNDROME	up to 30% MAJOR PSYCHIATRIC ILLNESS IN VELOCARDIOFACIAL SYNDROME
---	--	--

The final statistic measures a broader outcome. In velocardiofacial syndrome, up to **30%** develop major psychiatric illness, a category that includes schizophrenia and bipolar disorder. It must not be substituted for the figure for schizophreniform psychosis. This is a classic exam error: the number may be familiar, yet the construct and denominator are wrong. Anticipatory follow-up should therefore ask about emerging psychotic and affective symptoms without assuming that developmental or autistic features are their explanation.

13.7 Communicating the formulation without blame or fatalism

A syndromic formulation can be relieving when it gives a family language for a pattern that had previously been interpreted as wilfulness, poor parenting or inevitable deterioration. It should be

communicated as a vulnerability profile: “this condition makes this cluster more likely; we will look carefully at it and address what is modifiable.” That preserves realism and agency. It also makes room for strengths, relationships and the person’s own preferences, which a list of syndrome behaviours can easily erase.

GOOD TO REMEMBER

Use the phenotype to organise monitoring, environmental support and safety planning. For a food-related phenotype, explore access, supervision and conflict around food. For self-injury, define the topography, injuries, pain and immediate safeguarding needs. For a sleep-related phenotype, obtain a sleep history before interpreting daytime behaviour as purely psychiatric. For social disinhibition or withdrawal, assess vulnerability, consent, peer context and anxiety. These are clinical applications of the phenotype, not a generic management programme.

Be precise when discussing uncertainty. “Characteristic” does not mean inevitable; “shared” does not mean trivial; and “not specific” does not mean unimportant. The most defensible clinical note describes observed behaviour, names the syndrome-linked hypothesis, records alternative explanations and states what will be monitored. This reasoning protects the person from diagnostic overshadowing as well as from ignoring a syndrome-specific safety signal.

Must remember: A behavioural phenotype is a raised probability of a characteristic developmental or behavioural pattern, not genetic predestination. Read the whole cluster, its course and its context - then use it to anticipate needs without stereotyping the individual.

Living with intellectual disability

14 · Comorbid psychiatric disorders in ID (dual diagnosis)

Dual diagnosis is ordinary psychiatry, adapted rather than abandoned. In intellectual disability (ID), a psychiatric disorder may coexist with the developmental condition and must be sought as an additional clinical problem. In this setting, **dual diagnosis** means the coexistence of ID with an identifiable mental illness or personality disorder. The phrase originated in learning-disability psychiatry, although it has subsequently acquired a different substance-use meaning; state the context whenever using it.

The examination task is not to label every difficult behaviour as illness, nor to excuse illness as inevitable in ID. It is to establish whether there has been a meaningful change from the person's usual functioning, determine the function and context of that change, exclude physical and environmental explanations, and then decide whether a psychiatric syndrome is present. This is clinically consequential because communication differences can alter both the account available to the clinician and the outward form of distress.

3-5-fold RELATIVE RISK OF ANY PSYCHIATRIC DISORDER	40% AROUND CUMULATIVE PREVALENCE	9-18% ESTIMATED ADHD IN ID
--	--	--------------------------------------




Fig 21.12 Diagnostic overshadowing, drawn as the reasoning it actually is. One trigger, a behaviour that is new or an increase on a long-standing one in a person with intellectual disability, is put to two clinicians, and the plate follows each of them through three steps to a different place. The amber strip is why the error is expensive: on the IPS CPG's figures psychiatric disorder is 3 to 5 times more common in intellectual disability and around 40% meet criteria at some point, a prior probability the left column discards. No rate for overshadowing itself is drawn, because no source in this corpus gives one.

14.1 Burden and the meaning of excess risk

Psychiatric comorbidity is not an occasional complication of ID. Across the lifespan, the relative risk of any psychiatric disorder is **3-5-fold** that in the general population. In children and adolescents, studies similarly report a 3-5-fold excess compared with typically developing peers. These figures should change the clinician's threshold for asking an additional psychiatric question, but not the threshold for making a diagnosis without evidence.

Prevalence estimates differ substantially between studies, with reported values ranging from **10-39%**. The variation is informative: samples differ in age, severity of disability, setting, informants, case definitions, and whether the measure captures a current disorder, symptoms, or cumulative morbidity. A rate is therefore a prompt to assess, not a shortcut to diagnosis. Historical comparisons also remind us that ascertainment matters. In an Isle of Wight comparison, psychiatric disorder was identified in **50%** of children with intellectual disability and in 7% of children with normal IQ.

Risk is unevenly distributed. The severity gradient reported across epidemiological studies means that psychiatric disorder is more often identified with greater intellectual impairment. By contrast, the evidence is less clear for excess prevalence in adults with mild ID. Child and family risk markers include more substantial intellectual impairment, adaptive-behaviour and language difficulties, poor socialisation, and adverse family social circumstances. These are not diagnostic criteria and not explanations that make a person's symptoms less worthy of assessment. A co-occurring autistic spectrum disorder is itself an important risk context for anxiety and depression, bipolar-spectrum presentations, severe obsessional behaviour, anger disorders, and psychosis-like episodes that do not meet schizophrenia criteria.

■ **RULE** **Rule:** treat a changed mental state or behaviour in ID as a clinical question with several possible answers, not as proof of either psychiatric illness or developmental disability.

14.2 Diagnostic overshadowing, masking and the baseline

The central diagnostic hazard is **diagnostic overshadowing**: emotional or behavioural problems are attributed to ID itself instead of being considered as a coexisting disorder. At its extreme, the developmental diagnosis becomes so dominant that a second condition is never detected. Overshadowing is a clinician error of attribution. It can lead to a missed depressive episode, psychosis, anxiety disorder, delirium, pain state, or an environmental crisis because the referral is prematurely translated into “part of the disability”.

Diagnostic masking is different. Here, a developmental disorder modifies or obscures the manifestations of a primary psychiatric illness through cognitive, language, or speech limitations, particularly when disability is more severe. The illness has not been dismissed as ID; rather, it is harder to recognise because its usual verbal and phenomenological markers are inaccessible or altered. A mood disturbance, for example, may appear as excessive laughter or an increase in stereotyped behaviours rather than as a verbally reported low mood.

A third concept is **baseline exaggeration**. An increase in a long-standing behaviour may be the first, and occasionally the only, evidence of psychiatric disorder. Thus an established repetitive

act, irritability, aggression, or self-injury should not be dismissed merely because it is familiar. The question is whether its frequency, intensity, setting, consequences, or controllability has changed. The baseline is not background detail. It is the reference point that converts a behaviour into a possible symptom.

■ **TRAP** **Examination trap:** “Self-injury in a person with ID” is a presenting problem, not a diagnosis. Do not infer a psychiatric syndrome from the behaviour alone, and do not use the developmental diagnosis to end the enquiry.

PROBLEM IN REASONING	WHAT GOES WRONG	CORRECTIVE CLINICAL MOVE
Diagnostic overshadowing	A new psychiatric or physical problem is assigned to ID.	Ask what is new, then actively consider coexisting illness.
Diagnostic masking	The primary disorder has an altered, less verbal presentation.	Use behaviour, chronology, and informants alongside the interview.
Baseline exaggeration	A familiar behaviour is treated as clinically meaningless.	Compare current frequency, intensity, context, and impact with baseline.
Intellectual distortion	Answers to mental-state questions are overinterpreted.	Adjust language and seek corroboration rather than forcing adult phenomenology.
Psychosocial masking	Developmental symptom expression is mistaken for mental illness.	Interpret the observation in developmental context before diagnosing.
Cognitive disintegration	Acute disorganisation is equated immediately with chronic psychosis.	Assess for an evolving illness, stressor, physical cause, and change in supports.

14.3 Why the mental-state examination needs translation

Increasing severity of ID and associated communication impairment make diagnosis more complex. This does not imply that psychiatric diagnosis is impossible. It means that the clinician must translate the clinical question from a reliance on subjective report to a synthesis of observable behaviour, longitudinal change, collateral history, environmental context, and the person's own communication at their best level.

One useful framework describes **four** sources of diagnostic uncertainty: baseline exaggeration, **intellectual distortion**, **psychosocial masking**, and **cognitive disintegration**. Intellectual distortion concerns how the person understands a question about mental state and whether its implication can be appreciated. Psychosocial masking, in this formulation, describes developmental symptomatology that is misconstrued as mental illness. Cognitive disintegration describes disproportionate disorganisation during an emerging mental illness when cognitive reserve is limited, which can make the person look apparently psychotic. These are prompts for better assessment, not alternative diagnoses.

Behavioural equivalents are particularly important when verbal description is limited. In severe to profound ID, mood disturbance may be signalled by irritability, crying, dysregulation of sleep or appetite, agitation, mood lability, withdrawal, or isolation. Aggression or self-injury may express dysphoria at any developmental level, but neither is specific to mood disorder. Anxiety may present with aggression, agitation, compulsive or repetitive behaviour, self-injury, or insomnia. Panic may appear as agitation, screaming, crying, or clinging and may even look paranoid or delusional to an unwary observer.

Conversely, unfamiliar does not mean pathological. Self-talk can be a normal behaviour that is mistaken for psychosis. A complaint of being picked on may be the accessible form of persecutory ideation, but it also needs ordinary contextual testing. The task is phenomenology by approximation: ascertain what the person experiences where possible, establish what others observe, and decide whether the whole pattern forms a syndrome over time. In mild to moderate ID, mood disorders do not show striking differences in expression from those of typically developing peers; do not exoticise a presentation simply because ID is present.

14.4 A disciplined assessment sequence

Begin with change, not with labels. Assessment begins with the individual's baseline and interprets change with collateral information, behavioural observation, and context.

GOOD TO REMEMBER

Before treating current behaviour as psychiatric evidence, establish the person's previous baseline state. Family, friends, and carers often provide observations more useful than the brief assessment encounter. Use the interview alongside a period of behavioural observation. From first contact, consider approach, activity level, attachment to the informant, response to transitions, spontaneous communication, affective range, and reaction to demands against that baseline.

Build the history around the following dimensions rather than around a checklist of adult symptoms:

- onset and chronological evolution of the change;
- intensity and frequency of the behaviour or symptom;
- the context in which it appears and what follows it;
- precipitating factors, including changes in relationships, routine, supports, or expectations;
- relieving factors and the response to ordinary environmental adjustment.

This sequence gives the history a causal structure. Newly developing challenging behaviour can indicate a comorbid physical or mental disorder, an environmental or support change, or physical illness. Long-standing self-injury or aggression is more often understood developmentally or behaviourally. When change is sudden, give particular consideration to environmental changes, life events, painful physical conditions, and medication adverse effects. In the most disabled person with sensory deficits, exclude physical illness and stressful events before considering whether communicative frustration and understimulation are contributing.

Antecedent-Behavior-Consequences analysis is a functional method for clarifying what precedes a behaviour, what the behaviour achieves, and what consequences may maintain it. It is not a substitute for diagnosis. It prevents a false dichotomy: a psychiatric disorder can coexist with behavioural reinforcement, and a behaviour with an understandable function can still be an expression of distress. Prospective simple ABC or sleep charts are particularly useful when the care environment produces discrepant staff reports, since variation may reflect differing tolerance thresholds or attributions.

The most reliable discriminator is a documented departure from the individual's own baseline, interpreted with collateral observation and context.

14.5 Disorder patterns: recognise without overcalling

Detection of psychiatric syndromes in ID is shaped by communication, developmental level, and context. Schizophrenia should be diagnosed from a coherent longitudinal syndrome rather than from odd behaviour, self-talk, or a single report of suspiciousness. In a community sample of people with learning disability, point prevalence of schizophrenia was **over 4%**. Among people with learning disability and schizophrenia, auditory hallucinations are reported in around 90%; this is helpful when a reliable account can be obtained, but its absence or inaccessibility cannot settle the diagnosis.

Attention-deficit/hyperactivity disorder is also relevant: estimated prevalence among people with ID is 9-18%. The core diagnostic safeguard is developmental proportionality. Symptoms must be excessive for the person's mental age, and the threshold should be higher in severe to profound ID. Inattention, overactivity, impulsivity, poor frustration tolerance, and a chaotic environment can mimic each other. The clinician should seek a pervasive pattern rather than equating high activity in one setting with ADHD.

Anxiety estimates vary widely, from 1-25%, in part because diagnosis is difficult. This is a reminder to search for anxiety when distress is expressed behaviourally, not a basis for diagnosing it indiscriminately. Mood symptoms require the same restraint. Appetite and sleep change, social withdrawal, loss of usual enjoyment, tearfulness, agitation, or altered stereotyped behaviour gain significance when they are new, clustered, sustained, and not better explained by pain, medication, or altered circumstances.

■ **TRAP Examination trap:** do not diagnose psychosis from self-talk, depression from self-injury, or ADHD from activity level without establishing change, phenomenology, developmental proportionality, and context.

14.6 Instruments and adapted diagnostic systems

Questionnaires and structured interviews designed for children of average ability have a serious norming limitation in ID: almost none are normed for ID populations, and their scores or cut-offs may therefore be invalid. They may inflate symptoms or underestimate them depending on the setting. Use them as aids to gathering information, never as a rule-in or rule-out mechanism. This caution does not mean that all measures are unsuitable. The **Developmental Behavior**

Checklist is comprehensive and provides norms and subscales for the ID population, making it particularly useful where general-population measures fit poorly.

The clinical interview, observations, informant account, and functional data must retain primacy. ID-specific instruments may identify general psychopathology, yet studies do not show clear agreement between such instruments in detecting specific psychiatric disorders. A positive screen should trigger a more focused assessment; a negative screen should not override a persuasive longitudinal history. This is why an apparent numerical score is less important than the quality of the behavioural description from which it arose.

Modified diagnostic systems can support clinical reasoning where conventional criteria rely heavily on subjective experience or developmentally inappropriate thresholds. **DC-LD** was developed to address diagnostic issues in moderate to severe learning disability. **DM-ID** is another adapted diagnostic resource. Such systems do not create a separate psychiatry for ID; they make explicit the adaptations required to apply psychiatric nosology fairly. For eating disorders whose criteria depend substantially on subjective experience, conventional diagnoses such as anorexia nervosa and bulimia are effectively precluded in severe or profound ID. Pica is perhaps the most common eating disorder in this population.

14.7 Formulation and diagnostic accountability

A defensible formulation has parallel tracks. One track describes the person's developmental profile and baseline repertoire. A second describes the current change in behaviour, affect, cognition, biological rhythm, and functioning. A third examines physical illness, pain, sensory impairment, epilepsy, medication effects, interpersonal stress, care environment, and reinforcement contingencies. A fourth asks whether the data support a psychiatric syndrome, and if so what evidence distinguishes it from the alternatives. Writing these tracks separately prevents the developmental diagnosis from swallowing the entire formulation.

Behaviour that challenges has person-specific associations, including male sex, lower adaptive functioning, poorer communication, autism, and painful physical conditions. Unlike oppositional defiant or conduct disorder, it is not associated with family adversity. This distinction guards against importing a formulation from typically developing populations without testing whether it fits. Similarly, a psychiatric presentation may coexist with environmental contingencies; neither cancels the other.

Carefully document the source of every observation, the degree of confidence in the person's self-report, baseline evidence, collateral agreement or disagreement, and the physical and environmental exclusions considered. This makes diagnostic reasoning reviewable and protects against therapeutic drift toward medication for an unclearly defined target. Management of ID itself, including multidisciplinary support and educational planning, is addressed in the management section of this chapter. The full child psychiatric assessment battery, services and protection are covered in the Child psychiatry: assessment, treatment, services and protection notes.

Formulation should also preserve the person's perspective. A carer's account may be essential, but it should not replace direct engagement with the person. Adapt the pace, language, and mode of communication; allow time for response; and test important inferences gently rather than relying on a single answer. The purpose is not to force adult verbal psychopathology into a developmentally mismatched interview. It is to make the available evidence as fair, transparent, and clinically useful as possible. Where uncertainty remains, state it explicitly, define what further observation would discriminate between competing explanations, and avoid converting uncertainty into a confident label.

The practical standard is neither diagnostic minimalism nor diagnostic enthusiasm. It is intellectual humility with enough structure to notice a second disorder. When the pattern, chronology, collateral history, and context converge, people with ID deserve the same diagnostic seriousness afforded to anyone else.

15 · Management of ID (multidisciplinary, early intervention, special education)

Management is the organisation of opportunity. Intellectual disability is not managed by a single clinic, a prescription, or a school placement in isolation. The clinical task is to turn an assessment of abilities, barriers, health and family circumstances into a coherent plan that improves participation and preserves dignity. This makes management a longitudinal exercise: the clinician must help the person acquire useful skills, reduce avoidable barriers, respond to mental or physical illness when present, and keep the family able to continue caring.

Marks are usually earned by showing this architecture rather than by naming a succession of therapies. A good answer begins with coordinated goals, places early intervention and education within the person's everyday settings, explains how behaviour is understood before it is treated, and recognises that medical treatment has a limited, target-led role. Severity should be named only as a descriptor of support need; its classification belongs with the assessment and severity sections of this chapter.

15.1 The multidisciplinary plan

Multidisciplinary care is the appropriate treatment model because intellectual disability has biological, psychological, social and developmental consequences at the same time. It is therefore multimodal: psychoeducation, psychological work and medication, where indicated, are coordinated rather than offered as competing alternatives. The plan also has to restore wellbeing within the family system. In practice, the psychiatrist's distinctive contribution is diagnostic clarification, assessment of mental and physical contributors to change, risk formulation, judicious prescribing, and helping the team keep a shared hierarchy of goals.

The team should not be imagined as a referral list. Its work is to translate a common formulation into actions that can actually happen at home, in education, and in the community. For optimal management, coordination includes at least six disciplines, including psychiatry, psychology, occupational therapy, speech and language therapy, genetic counselling and nursing. This is not an exhaustive job description. It illustrates why a locally available team should be assembled around the person's needs rather than around an idealised service chart.

- **RULE** **Rule:** Start with shared functional goals, then assign one accountable person for coordination; referrals without a joint plan merely move fragmentation from one service to another.

SERVICE FUNCTION	CLINICAL PURPOSE	WHAT COORDINATION PREVENTS
Prevention and prompt recognition	Identify needs before they become fixed barriers to development or participation.	Late referral after family strain or school failure has already escalated.
Strengthening family capacity	Make everyday care, communication and goal-setting possible.	Advice that is technically sound but impossible to sustain.
Learning and productive roles	Build participation and future roles around meaningful skills.	Care that remains custodial when the person can learn more.
Health and mental-health care	Address health needs and emotional or behavioural presentations without diagnostic overshadowing.	Attributing a new problem automatically to the disability.
Living support and self-care	Match support to real-world functioning and safety.	Discharge plans that fail because ordinary living demands were ignored.

The level of specialist involvement should follow complexity. Mainstream mental-health services may prescribe for mild or borderline intellectual impairment, while assessment and treatment of behavioural or emotional disorders in more marked or atypical cognitive impairment should be undertaken with specialist clinical input. This is a service-triage principle, not a claim that a mainstream clinician should work alone when communication barriers or atypical presentation make assessment uncertain.

Family work is clinical work. Families need an explanation that can be revisited, not a single overwhelming disclosure. Parents may require repeated discussion before the implications can be absorbed; the clinician should return to prognosis, available help and the family’s active role as understanding changes. Family support must be individualised to the caregiving burden and its meaning for that particular household. Parents and primary caregivers should also be routinely screened for stress-related disorders. This is not a courtesy add-on: depleted carers have less capacity to implement a plan, and their distress deserves treatment in its own right.

15.2 Early intervention as a developmental service

Early intervention begins before a child has entered an educational system. The Indian guideline permits a starting point from the prenatal period, with recognition of high-risk pregnancies, appropriate health care and attention to psychosocial adversity. After birth, the plan is not simply stimulation exercises. It begins with accurate identification of developmental and associated needs, then creates opportunities for sensory-motor, communication and socioemotional development in the child’s ordinary relationships and routines.

The central relational mechanism is healthy bonding and attachment between the child and mother. The practical implication is important: early stimulation can also be delivered by any

stable caregiver. Programmes should stabilise current developmental achievements while opening the next achievable task. Play-based work and culturally rooted early-child-care practices are often more usable than imported materials; when materials are inaccessible or culturally mismatched, parents can become overwhelmed and the programme loses its place in daily life.

Evidence supports a measured, not triumphalist, account. **Family-centred intervention** can be effective, but uncertainty remains about the active components and how help is best delivered. This is why the clinician should set observable functional goals and revise them with the family, rather than promise a uniform developmental result. One intensive compensatory-education experience illustrates both the possibility and the limits of inference: teaching began at 3 months; at a later preschool assessment the intervention group had a mean advantage of 27 IQ points over controls. The finding does not prove a durable individual IQ gain, and its design limitations matter. Its defensible teaching point is that sustained, skilled effort with socially disadvantaged families can make a meaningful difference.

3 years INITIAL LEARNING PHASE	3-6 SCHOOL-READINESS PHASE	6-18 CONSOLIDATION PHASE
--	--------------------------------------	------------------------------------

A life-span skill plan has three **developmental stages**. In the initial years, work addresses sensory-motor ability, socio-communication, basic self-help and concepts. The subsequent school-readiness period emphasises readiness skills and culturally appropriate adaptive behaviour. Later childhood and adolescence consolidate academic and independent personal skills in preparation for vocational training, employment and adult living. These are organising priorities, not rigid age-based ceilings. The immediate task is always to find the skill that will increase autonomy in the child’s own context.

IN INDIA

For formal early intervention in India, referral can be made to the District Early Intervention Centers of the Rashtriya Bal Swasthya Karyakram and the Anganwadi Centers of the Integrated Child Development Services. The referral should carry a concise developmental formulation and the family’s priorities, so that the service encounter becomes part of a continuing plan rather than a one-time certification exercise.

The clinical details of normal milestones and attachment are covered in the Normal child development and milestones notes.

15.3 Education, participation and placement

Normalization is the guiding principle of arranging a pattern of life as close to ordinary life as possible, supported by specialist help that allows the person to reach potential. It directs attention away from segregation as a default and toward family life, community participation, education, work and self-advocacy. It does not mean pretending that support needs do not exist. Normalisation becomes clinically useful only when the necessary teaching, communication support, supervision and environmental adjustments are actually provided.

Placement is a clinical and ethical judgement. Ordinary-school education may offer more normal social surroundings, social integration, expectation of progress and learning for peers that participation is ordinary. It may also lack the teaching skills, equipment and methods required by some learners, particularly where specific language and communication teaching is needed. The evidence cited here does not settle which children benefit from ordinary schooling. Therefore, do not turn inclusion into a slogan or special education into a failure. Consider the child's learning needs, social experience, available supports and family preference, then review the placement when the fit changes.

■ **TRAP** **Exam trap:** "Mainstream" and "beneficial" are not synonyms. Educational inclusion without the support needed to learn or remain safe is not meaningful participation.

The ethical tension is real. In secondary schools, children with special needs were reported to be bullied **three times more often** than other children. This should prompt active safeguarding and a discussion of the individual child's present interests, not a simplistic conclusion that community education is always unsafe. At a service level, a systematic review found better outcomes with community-based services than with institutional care. The practical inference is to design community support well, monitor the child's experience, and address exclusion rather than treating it as inevitable.

Education should be functional. Literacy, numeracy, communication, self-care and social learning have value insofar as they enlarge the person's choices in daily life. At the end of school, reassessment and vocational guidance are necessary before a young person is expected to make an adult transition. Opportunities should be ambitious but realistic, and should not be narrowed merely because others hold limited expectations. The fuller approach to disability certification, benefits and statutory entitlements belongs in the disability certification and benefits section of this chapter.

15.4 Behavioural formulation before behavioural control

Behaviour that challenges is a question, not a diagnosis. Its onset or worsening should lead first to a search for communication difficulty, environmental demand, reinforcement, health problems and mental disorder. A **comprehensive personalised functional assessment** considers behaviour in its setting and improves the chance of a plan that can be managed psychologically and environmentally. It can consequently reduce, and sometimes remove, the need for psychotropic medication.

Functional behaviour analysis asks what consequence may be maintaining the behaviour. Challenging behaviour may serve three functional classes: obtaining a social positive reinforcer such as attention, escaping a demand through social negative reinforcement, or obtaining sensory reinforcement. These possibilities are hypotheses to test against observation, not labels to attach from a brief history. Parents, teachers and other caregivers are indispensable informants because they see the antecedents and consequences that a clinic visit cannot reproduce.

The behavioural plan should be least intrusive and culturally appropriate. The Indian guideline describes **three** linked levels: restructure the environment to alter antecedents, reinforce adaptive behaviour differentially, and prevent inappropriate reinforcement of problematic behaviour. Its basic premise is functional substitution. If a problematic behaviour communicates a need, the plan must create and reinforce a socially acceptable behaviour that serves the same function. Suppression alone leaves the original need unanswered and is likely to fail once the therapist has left.

- Define a small number of observable target behaviours in the settings where they occur, and prioritise those causing exclusion, stigma or interference with learning.
- Record antecedents, behaviour and consequences with carers, then decide which maintaining contingencies are plausible.
- Teach and reinforce an adaptive alternative that achieves the relevant function more safely.
- Change environmental triggers and caregiver responses together, then review whether the person's participation has improved.

Positive behavioural support is a useful framework because it combines reinforcement with changes to triggers, minimal use of aversive strategies and planned reactive responses such as distraction. Its components should not be applied as a menu of techniques. Selection follows the formulation. Techniques using negative reinforcement should not be used. A discipline-heavy behaviour programme is best developed and implemented by a behavioural psychologist or board-certified behavioral analyst, although psychosocial interventions delivered by non-specialist carers can produce better outcomes than no treatment or treatment as usual. In settings with few specialists, this supports task-sharing with supervision and training, not abandonment of formulation.

15.5 Medication stewardship and specialist liaison

GOOD TO REMEMBER

Medication treats a defined target symptom, coexisting mental disorder or physical contributor; it does not replace a formulation of behaviour. **Psychological intervention**, including functional behavioural analysis, should be considered first-line when no identifiable mental illness with clear treatment implications is present, except in the most serious or intractable behavioural disturbance. The same discipline applies when diagnostic information is inadequate: try psychosocial intervention before converting uncertainty into a prescription.

Antipsychotics remain the most widely prescribed psychotropic class in intellectual disability, yet their current positioning is as a later option for older children or adults when intensive behavioural intervention or other medication options have failed. This apparent contradiction is a warning about service reality, not a reason to disregard behavioural care. A functional formulation may reveal a modifiable trigger that makes medication unnecessary, whereas a clear psychiatric syndrome or severe risk may require simultaneous medical treatment and behavioural support.

When a dopamine antagonist is considered for behaviour that challenges, the NICE guideline NG11 identifies **three** alternative qualifying circumstances: psychological or other intervention alone has not changed the behaviour within an agreed time, treatment of a coexisting mental or physical problem has not reduced it, or risk is very severe. These are alternatives, not a checklist in which every condition must be met. Once medication is begun, prescribe the lowest possible dose and reassess whether continuation remains justified after target symptoms improve, when there is non-response, or when adverse effects affect functioning or quality of life.

The management sequence is: clarify function and illness, strengthen environmental and behavioural supports, prescribe only for a defined target, and keep asking whether the drug is still needed.

Deprescribing needs the same care as initiation. The NICE guideline NG54 suggests slow withdrawal of antipsychotics for people without psychotic symptoms. This does not mean that withdrawal is consequence-free: discontinuation studies commonly, though not invariably, show re-emergence of problem behaviours. The appropriate response is preparation, monitoring of the original target behaviour, and readiness to revise environmental support, rather than assuming either that every drug must continue or that every drug can be stopped without difficulty. The UK STOMP programme promotes antipsychotic deprescribing but also shows a crucial pitfall: an antipsychotic may be replaced by another psychotropic rather than reducing overall medication burden.

Drug choice for a diagnosed condition may in general resemble practice in the wider population, but good prescribing practice is to start at lower doses and increase more slowly. This distinction resolves an apparent contradiction between using standard treatments and being more cautious in a person who may have communication barriers, neurological vulnerability or difficulty reporting adverse effects. A review should document the target symptom, benefit, adverse effects, family observation and whether non-drug supports remain active. Annual health checks are generally recommended for people with intellectual disability; they are health surveillance, not a substitute for medication review whenever treatment changes.

15.6 Continuity into adult life

Management succeeds when it survives transitions. A community-care model is judged less by its label than by the detail of planning and the commitment with which it is carried out. Planning needs both an understanding of the local population and an individual account of what this person can do, wants to do, and needs help to do. A named coordinator can prevent families from receiving inconsistent advice across health, education, social and voluntary-sector services.

Families often need additional help at puberty and school leaving; moving from child to adult services can be especially stressful. These periods should trigger anticipatory planning rather than a crisis referral. Reassessment and vocational guidance before school leaving protect against the common drift from education into inactivity. For school leavers with mild intellectual disability, vocational planning may lead to ordinary or sheltered work; adults with severe

disability are more likely to need the support of a day centre. The point is not to predict a person's future from a label, but to plan enough support for a real adult role.

Referral and linkage in India should map the person's needs to education and vocational routes, rehabilitation centres, national institutes and local nongovernment agencies. Whenever possible, link the family to agencies in their locality so that rehabilitation costs and travel do not quietly defeat the plan. The clinician should retain a longitudinal role: explain the plan repeatedly, detect emerging health or psychiatric problems, coordinate with the relevant team, and protect the person's right to participate in family and community life.

16 · Prevention of intellectual disability

Prevention is a life-course clinical task. Prevention of intellectual disability is not a promise that every case can be avoided. It is the organised reduction of preventable risk before conception, during pregnancy and birth, through the newborn period, and after disability is recognised. The examination value of the topic lies in distinguishing prevention of a condition from prevention of avoidable functional disadvantage once a person already has intellectual disability. A good answer therefore shows a framework, gives a few high-yield examples at the correct level, and retains ethical humility.

The field uses several overlapping taxonomies. The most useful approach in an intellectual-disability answer is to state the purpose of the intervention rather than force every programme into a label. Public health measures reduce exposure to risk; screening creates an opportunity for timely action; and habilitative care limits the secondary disabilities that arise when a person's needs are neglected. These are connected steps in one pathway, not competing lists.

16.1 What prevention means in intellectual disability

The conventional framework has **three** categories. **Primary prevention** acts before a disorder develops, so its clinical purpose is to reduce new cases. For intellectual disability, this means combining medical care with public-health, educational and social measures before an avoidable insult has produced disability. The point is causal timing: an intervention is primary when it removes or reduces the exposure before the outcome arises.

Secondary prevention aims to reduce the burden of established disease in the general terminology, but the intellectual-disability application is more clinically practical: recognise a condition that can lead to disability as early as possible and treat it before disability follows. This makes screening meaningful only when a result can lead to confirmation, treatment, counselling or follow-up. A test with no route to action may collect information, but it does not by itself achieve prevention.

Tertiary prevention concerns disability associated with an existing illness. In intellectual disability it is best understood as **habilitation**: helping the person attain the best attainable functioning despite the disability. This includes biomedical and sociocultural work, but it is not a synonym for cure. The target is participation, competence and protection from avoidable loss of function.

- **RULE** **Rule:** Classify a preventive intervention by its intended point of action: before disability arises, while a remediable pathway is still open, or after disability is present to prevent additional functional loss.

PREVENTIVE LEVEL	CLINICAL QUESTION	INTELLECTUAL-DISABILITY APPLICATION	COMMON ERROR
Primary prevention	Can the causal exposure or pathway be avoided?	Genetic counselling, population health, safer pregnancy, nutrition, immunisation and prevention of toxic or traumatic injury.	Calling every antenatal test prevention without asking what action follows, or filing genetic counselling under care that comes after a diagnosis.
Secondary prevention	Can a disorder that may cause disability be found and treated early?	Early recognition and treatment of metabolic conditions or congenital hypothyroidism.	Equating screening with treatment.
Tertiary prevention	How can further functional disadvantage be reduced after disability is present?	Habilitation, education, family support and treatment of comorbid conditions.	Presenting support as an afterthought rather than prevention of secondary disability.

The alternative language of universal, selective and indicated prevention is also useful in public health. Universal work reaches whole populations, selective work addresses groups with increased risk, and indicated work is directed at people with individual evidence of heightened risk. Do not casually equate every use of indicated across classifications. The later Institute of Medicine model, published in 1994, uses indicated intervention for people with early or prodromal symptoms, whereas the earlier approach used it for asymptomatic people identified as higher risk. The boundary between selective and indicated work can be clinically blurred; the safer answer is to state the target population and the reason for intervention.

- **TRAP** **Exam trap:** Primary, secondary and tertiary prevention are not interchangeable labels for antenatal, postnatal and rehabilitation services. Timing matters, but the decisive issue is what the intervention is meant to prevent.

16.2 Primary prevention before and during pregnancy

Primary prevention begins before pregnancy is recognised. Good preconception and antenatal care allows the clinician to identify modifiable risks, improve maternal health and support informed reproductive decisions without converting counselling into coercion. The Indian guideline specifically identifies malnutrition, diabetes, teratogenic drugs and substance use as maternal factors that may be treatable or preventable in subsequent pregnancies. A prior child with intellectual disability should prompt a respectful etiological review and a plan for the next pregnancy, not an assumption of parental fault. That review is where **genetic counselling** belongs, and it belongs here rather than later: the standard account of primary prevention opens with genetic counselling and continues with education of pregnant women about behaviours that may put the fetus at risk, early detection of fetal abnormalities during pregnancy, and good

obstetric and perinatal care. Counselling a couple before a further conception acts on the next pregnancy, so it removes the exposure before the outcome arises, which is the test this section has just set. Reading it as an after-the-fact service, because the conversation happens after a diagnosis in an older child, is the classic misplacement, and naming the primary preventive measures is a canonical examination question whose first answer is this one.

Risk reduction works through ordinary care. Primary measures include prevention of lead poisoning, avoidance of substance exposure during pregnancy, prevention of teenage pregnancy, improved neonatal care and injury prevention such as infant car seats. Prenatal vitamin supplementation is included because it prevents neural tube defects. The phrasing matters: prenatal supplementation should not be inflated into a guarantee against all developmental disorders. Likewise, childhood lead exposure is a risk factor for low IQ and intellectual disability, which supports environmental prevention without claiming that every exposed child will develop disability.

Maternal infections are another preventable pathway. Toxoplasmosis, cytomegalovirus, rubella and herpes are recognised maternal infections associated with infant morbidities that include intellectual disability. Rubella immunisation is therefore an important prevention measure. Prenatal care may begin before conception, including rubella immunisation for women without immunity. Prevention of rhesus sensitisation uses **anti-D antibody**. These examples should be used to show an integrated maternal-health strategy, not as a memorised infection list detached from antenatal care.

Alcohol deserves direct enquiry in all settings where it is accessible. In such societies, alcohol is the leading toxin cause of intellectual disability. Fetal alcohol syndrome has an incidence of **0.5-3 per 1000 births** and may account for as much as 3 percent of all intellectual disability. These are different denominators, and they must not be swapped. Apparent rates also depend on diagnostic practice: use of both physical signs and a history of heavy prenatal alcohol exposure produces lower rates than use of exposure history alone, with underdiagnosis and overdiagnosis possible at the two extremes.

Nutritional prevention has its own logic. Iodine deficiency is described as the most common preventable cause of learning disability worldwide. Where endemic iodine deficiency threatens pregnancy, periconceptional injection of **iodized oil** has attenuated endemic cretinism and subsequent intellectual disability in at-risk populations. A pharmacology source also states that daily maternal iodine intake of **150-200 microgram** can prevent endemic goitre and cretinism due to iodine deficiency, with iodised edible salt as the preferred delivery method. This is a prevention claim, not an invitation to prescribe an unsupervised supplement outside local obstetric and public-health practice.

one-third ID/GDD CASES WITH NONGENETIC CAUSES	over eighty POTENTIALLY TREATABLE DISORDERS IDENTIFIED	over 50 CONDITIONS DETECTED BY CONTEMPORARY NEWBORN SCREENING
---	--	---

The limits are as important as the opportunities. Nongenetic factors account for approximately one-third of ID/GDD cases, so environmental prevention is necessary but cannot remove all intellectual disability. At the same time, advances in therapeutics have identified over eighty potentially treatable disorders. This justifies careful diagnosis and prompt action without selling prevention as total control. Prevention priorities differ by setting: maternal and perinatal health are central public-health opportunities in low- and middle-income countries, while reduction of genetic causes of severe intellectual disability remains important in high-income countries. It is unlikely, however, that prevention can substantially alter the incidence of mild intellectual disability.

16.3 Antenatal screening, diagnosis and ethical care

Non-invasive prenatal screening is based on cell-free fetal DNA in maternal plasma. Its value is risk estimation and selection for counselling or diagnostic evaluation, not a diagnosis by itself. First-trimester screening combines an ultrasound marker with maternal blood testing. An ultrasound assessment for major malformations is often performed at approximately 20 weeks of gestation. These routes should be explained as complementary rather than as competing tests.

CLINICAL ROUTE	WHAT IT CAN CONTRIBUTE	TEACHING POINT
First-trimester screening	Risk assessment through ultrasound and maternal blood analytes.	A positive screen requires counselling and a diagnostic pathway.
Mid-trimester AFP testing	Screening for neural tube defects in the specified screening context.	An analyte result is not a diagnosis and is not specific to one disorder.
Amniocentesis	Diagnosis of chromosomal and metabolic disorders.	Use the test for the question it can answer.
Chorionic villus sampling	Chromosomal and molecular genetic studies.	Do not transfer amniocentesis metabolic role to CVS.

In one psychiatric-text account, amniocentesis or chorionic villus sampling is recommended for all women aged **35 years or more**. Another account frames amniocentesis as an indication-based decision for pregnancies with relevant prior or familial history, or a clinically significant current risk. The defensible clinical response is to know that recommendations are guideline- and context-dependent, establish the indication, explain benefits and limitations, and refer for current specialist obstetric or genetics advice rather than use a textbook threshold as a stand-alone rule.

Screening needs careful language. Maternal serum alpha-fetoprotein is a marker with more than one possible clinical explanation; it is not a disease label. For in-utero diagnosis of Down syndrome, the source describes a blood-based route, ultrasound assessment and invasive testing. These options differ in what they establish and what follow-up they entail. Carrier screening should be offered to all persons in high-risk populations, but named examples should follow locally relevant, verified services rather than stereotype or assumption.

Prenatal diagnosis followed by elective termination raises a boundary that clinicians should not obscure. It is a matter of deeply personal values because termination prevents childbirth rather than treating or preventing the disorder in the fetus. The psychiatrist's role is to support informed, non-directive decision-making, recognise family distress, avoid coercive framing, and ensure that a decision is not driven by misinformation about disability or by a lack of support for families who continue a pregnancy. The syndrome-specific genetics and recurrence discussion belongs with the relevant aetiology sections of this chapter.

16.4 Newborn screening and time-sensitive treatment

Newborn screening refers to tests undertaken within the first hours or days of life when early action can prevent severe health problems. Contemporary systems detect over 50 different conditions. A blood spot collected by heel stick is the established specimen method in the settings described in the source.

GOOD TO REMEMBER

An abnormal screen is a signal for confirmation, not a diagnosis or a moral judgement about the family. Communicate promptly and compassionately, then ensure confirmatory testing, treatment and longitudinal follow-up.

Congenital hypothyroidism illustrates the time-sensitive logic. With newborn screening, levothyroxine may be initiated by **4-6 weeks of life**. Screening of neonates is described as the best preventive strategy for cretinism. This should not be reduced to a prescription detail: early identification links a population programme to a child developmental outcome. The quoted treatment timing is not the time at which the blood sample must be collected.

The overall verdict on newborn metabolic screening is appropriately strong but bounded: it has been extremely successful in reducing the incidence and severity of some intellectual-disability syndromes. The qualifier *some* is crucial. A screening programme may also generate false-positive results, which can produce acute parental distress and unnecessary follow-up unless communication is prompt and compassionate.

Phenylketonuria is a classic screenable metabolic disorder, but its genetic mechanism and diet belong in the inborn errors of metabolism section. Here, the prevention lesson is narrower: early diagnosis and treatment of PKU, other metabolic conditions and congenital hypothyroidism are examples of secondary prevention because they intervene before disability develops.

16.5 Preventing secondary disability after diagnosis

Once intellectual disability is present, the preventable outcome is often loss of function rather than intellectual disability itself. Tertiary prevention therefore uses early support, education, family support and timely attention to mental or physical conditions that suppress functioning. This is habilitation in action. It safeguards developmental opportunity and reduces the additional disability created by untreated epilepsy, pain, sensory difficulty, psychiatric illness, behaviour problems, exclusion or caregiver exhaustion.

Do not wait for a crisis to begin support. Early intervention may start in the prenatal period by identifying high-risk pregnancies, arranging appropriate health care and addressing psychosocial adversities.

IN INDIA

For formal early intervention, the guideline names the District Early Intervention Centres of the Rashtriya Bal Swasthya Karyakram and the Anganwadi Centres of the Integrated Child Development Services. The preventive point is timely connection before a modifiable disadvantage becomes entrenched.

The service model, developmental methods and general evidence for early intervention are covered in the management section of this chapter.

Comorbidity management is central, not peripheral. Epilepsy affects 15-30 percent of persons with intellectual disability. Problematic behaviours may occur in 20-80 percent of the ID population. These figures do not authorise diagnostic overshadowing. A new behavioural change may be communication of pain, distress, environmental difficulty or mental illness. Prevention in tertiary care means looking actively for such contributors and treating them before they further reduce learning, relationships or family capacity. The psychiatric-comorbidity assessment itself is addressed in the dual-diagnosis section of this chapter.

Families may need genetic counselling alongside causal investigation and reproductive planning. It should be offered without blame and without making a risk discussion substitute for the family's own values or choices. The generic genetic-counselling process and diagnostic work-up are developed in the aetiology section.

16.6 Public-health priorities and realistic expectations

Prevention is most effective when it reaches the conditions in which children develop. ID risk factors are considerably increased in low-income countries, contributing to their increased rate of intellectual disability. This is why a prevention answer should move beyond rare disorders and include maternal health, nutrition, safety, infection control, toxic exposure and access to perinatal and neonatal care. It is also why prevention must be paired with equity: a family cannot implement advice that depends on services, food security or safe housing it cannot access.

Evidence supports interventions at more than one level. Programmes aimed at psychosocial stimulation were more effective than nutritional supplementation for improving both cognition and behaviour in growth-retarded children in the cited comparison. This does not diminish nutrition. It teaches that development depends on both biological provision and responsive social experience, and that a single nutrient-focused programme may not repair the effects of deprivation. Similarly, prevention programmes should be evaluated for whether they can be delivered consistently and whether they improve real developmental participation.

A comprehensive service begins with prevention and early detection and then connects families to the educational, health, social and practical supports that sustain participation. Prevention is therefore the first element of a continuum of care, not a separate campaign that ends after a

screening test. Appropriate referrals should follow the person across early, educational and adult life. The statutory eligibility and benefits themselves are dealt with in the disability certification and benefits section of this chapter.

MNEMONIC

PREVENT

Prepare before conception; **R**educe maternal, toxic and nutritional risks; **E**valuate antenatal and newborn screens; **V**erify a positive result; **E**ngage family support; **N**otice comorbidity; **T**arget function through habilitation.

The mature answer is neither fatalistic nor triumphalist. It recognises preventable pathways, acts early when an actionable condition is found, and continues to prevent additional disability through support and treatment. It also respects the person with intellectual disability as more than the outcome of a risk factor. Prevention should enlarge future choice, not become a language for blaming parents, narrowing reproductive autonomy or measuring a life only by what might have been avoided.


- Ask which preventable exposure or remediable condition is relevant at the present point in the person's life course.
- Make sure a positive screen has a confirmation, treatment and follow-up pathway.
- After diagnosis, look for treatable illness and avoidable barriers to participation rather than assuming that decline is inevitable.

For intellectual disability, prevention is a continuum: reduce avoidable risks before disability arises, detect and treat remediable conditions early, and use habilitation to prevent avoidable loss of function after diagnosis.

17 · Disability certification and benefits for intellectual disability in India (IDEAS, National Trust Act)

Certification is a clinical document with legal consequences. For a person with intellectual disability, it can convert an assessment into access to education, work, social protection and supported decision-making. The psychiatrist's task is therefore not merely to write a diagnosis. It is to distinguish clinical formulation from statutory quantification, obtain defensible evidence of functioning, explain what the certificate does and does not establish, and help the family navigate a system that can otherwise feel opaque.

The contemporary legal frame is the **Rights of Persons with Disabilities Act, 2016**, notified on **28 December 2016**. It replaced the **Persons with Disabilities (Equal Opportunities, Protection of Rights and Full Participation) Act, 1995**. Intellectual disability is a specified disability under this framework, alongside other conditions relevant to psychiatric practice. This section concerns the certification route for intellectual disability, its practical consequences, the limited role of IDEAS, and the National Trust framework for adulthood and guardianship.



17.1 The legal frame: from diagnosis to enforceable participation

The Act is not a diagnostic manual. Its purpose is to translate disability into rights, duties and institutional remedies. It has 17 chapters and 102 sections; its Schedule names 21 specified disabilities, compared with 7 disability types in the earlier law. Within the Schedule, intellectual disability appears at serial 7 and autism spectrum disorder at serial 9. The nomenclature matters: the Act replaces **“mental retardation” with “intellectual disability”**. In clinical records and communication, use the current term consistently.

The statutory definition recognises limitation in intellectual functioning together with adaptive behaviour in everyday social and practical skills. That overlap with clinical assessment is important, but the questions asked are different. A clinical assessment asks what the person can do, what has changed, which supports work and whether there are treatable coexisting problems. A certificate establishes disability in the form required for rights and entitlements. It should never flatten the person into a percentage or be allowed to substitute for an individualised care plan.

■ **RULE** **Clinical severity and certified disability are related, but they are not interchangeable labels.** Clinical formulation guides care; statutory quantification establishes eligibility under a legal scheme.

For intellectual disability, the statutory schedule describes significant limitation in intellectual functioning together with limitation in adaptive behaviour. It does not legislate an IQ number, and a candidate should resist the temptation to supply one: the percentage on the certificate is derived from adaptive functioning, not from the IQ score. This is an easily missed distinction. A numerical cognitive test can corroborate intellectual impairment, but it does not itself produce the statutory percentage. Candidates who equate the two have confused an assessment construct with a legal outcome, and quoting a legal IQ cut-off that the statute never states is the sharpest form of that error.

- Use the diagnosis to organise clinical understanding and intervention.
- Use the certificate to document statutory disability and enable access to rights.
- Explain to caregivers that the certificate records function for a stated purpose, not the person's potential or worth.

17.2 The intellectual-disability certification assessment

Start with the correct question. The assessment must establish intellectual disability and then quantify disability in the prescribed manner. Careful administration and a credible developmental history are especially important; a rushed score, divorced from the person's actual opportunities and supports, is a poor foundation for a legal document.

IN INDIA

The Indian guideline identifies the VSMS as the currently available standardised adaptive-behaviour measure. The prescribed adaptive-functioning instrument is the **Vineland Social Maturity Scale (VSMS)**; it yields a **social quotient (SQ)** and an adaptive profile. Intellectual functioning is assessed using the **Binet-Kamat Test** or **Malin's Intelligence Scale for Indian Children**.

Cognitive testing in this pathway provides diagnostic confirmation, not percentage derivation. Where IQ and SQ suggest different grades, the SQ should guide the decision, because social maturity more closely reflects the ability to meet culture-appropriate demands of daily life.

GOOD TO REMEMBER

An implausible or discordant result should prompt review of the history, test conditions, language, sensory or motor barriers, educational opportunity, behaviour during testing and the informant's understanding of the items.

A statutory assessment should be clinically thorough even when its output is concise. Establish the developmental course and current adaptive abilities from an informant who knows the person across settings. Consider whether communication difficulty, behavioural disturbance, sensory impairment, physical illness or an untreated psychiatric disorder has distorted apparent function. Record observed abilities as well as deficits. The certificate process should protect against both overestimation, which deprives a person of support, and underestimation, which may unnecessarily restrict autonomy.

■ **TRAP** **Do not use an IQ score to calculate the intellectual-disability percentage.** In this certification pathway, the percentage follows the VSMS-based adaptive-functioning assessment; IQ testing confirms the intellectual impairment.

The categories below are statutory bands, not a substitute for the clinical severity discussion elsewhere in this chapter. They answer the specific legal question of what disability percentage is certified. The row should be read as a mapping: identify the applicable VSMS band, then document the corresponding percentage without improvising a conversion.

CERTIFICATION CATEGORY	VSMS SCORE BAND	CERTIFIED DISABILITY	PRACTICAL INTERPRETATION
Profound	0-20	100%	Highest statutory disability band
Severe	21-35	90%	Benchmark-disability threshold is met
Moderate	36-54	75%	Benchmark-disability threshold is met
Mild	55-69	50%	Benchmark-disability threshold is met
Borderline disability	70-84	25%	Does not meet benchmark-disability status

The borderline-disability row is clinically useful because it makes the eligibility boundary explicit. It is not a reason to dismiss need. A person may still need educational, vocational, family or mental-health support even when the statutory benchmark threshold is not met. Conversely, a benchmark certificate creates a legal entitlement pathway but does not tell the clinician which supports will be effective. Keep these two conversations separate and honest.

17.3 Application, medical authority and validity

The route should be explained before the family is sent from counter to counter. An application for a disability certificate is made in **Form IV**. When a person with intellectual disability cannot apply, a legal guardian may apply; an organisation registered under the Act may also apply where it has a minor in its care. This provision is practical rather than merely administrative: it avoids making access to certification contingent on an ability that is itself affected by the disability.

The medical authority is headed by the Medical Superintendent, Chief Medical Officer, Civil Surgeon or an equivalent state-notified authority. Its clinical membership includes a paediatrician or paediatric neurologist where available, with age-appropriate alternatives specified for adults, a clinical or rehabilitation psychologist, and a psychiatrist. The psychiatrist’s contribution is more than signing a completed form. It includes diagnostic coherence, review of developmental and behavioural information, recognition of coexisting psychiatric illness, communication with the family and protection of the person’s dignity during examination.

Certification can be issued after at least **1 completed year of age**. The label is age-sensitive: the guidance uses Global Developmental Delay from ages 1 to 5 years and Intellectual Disability after age 5 years. This distinction avoids premature certainty while preserving access to assessment and support. Do not delay a needed certificate simply because the child’s development is still being clarified; use the appropriate certification label and plan review.

WHEN THE CERTIFICATE IS ISSUED	VALIDITY APPROACH	CLINICAL IMPLICATION
Before age 5 years	Temporary; it ends at the earlier of age 5 or 3 years.	Plan reassessment rather than treating the early certificate as final.
Between ages 5 and 17 years	Review is scheduled at ages 5, 10 and 18 years.	Make renewal part of transition planning, not a crisis near a deadline.
At age 18 years	Lifelong validity	Use the adult review to address future support and decision-making.

The medical authority must issue the certificate or communicate the reason for rejection within at most **1 month from receipt of the application**. In practice, clinicians can reduce avoidable delay by ensuring that reports identify the tool used, the functional basis of the percentage, the diagnosis appropriate to age and any need for follow-up. Keep a copy of the assessment record: it is clinically useful at renewal and can clarify an apparent discrepancy without making the family repeat an entire evaluation.

17.4 Benchmark disability and the rights that certification can activate

A **person with benchmark disability** has certified disability of at least **40%**. This is an eligibility threshold, not a clinical threshold. It determines access to several statutory provisions, whereas care should respond to need at every level of disability. Explain this plainly. Families often hear “eligible” as “severely disabled” or “ineligible” as “no help is possible”; both inferences are wrong.

40% BENCHMARK THRESHOLD	5% HIGHER-EDUCATION SEATS	4% GOVERNMENT VACANCIES	1% SHARED JOB-RESERVATION CLUSTER
-----------------------------------	-------------------------------------	-----------------------------------	---

A child with benchmark disability has a right to free education between 6 and 18 years. Government and government-aided higher-education institutions reserve at least 5% of seats for persons with benchmark disabilities, with an upper-age relaxation of 5 years for admission. In every Government establishment, at least 4% of vacancies are reserved. These provisions should not be presented as a promise of a particular educational placement or job. The clinician should document function accurately and direct the family toward the relevant institutional process rather than make an outcome guarantee.

The employment reservation has a subtle but examinable structure. Autism, intellectual disability, specific learning disability and mental illness form one listed cluster. That cluster does not receive a separate one-percent allocation by itself: it shares exactly 1% of vacancies with the multiple-disabilities cluster. This clarification prevents the common answer that incorrectly assigns the whole sub-quota to intellectual disability or to the listed cluster alone.

- Discuss educational entitlement early enough for the family to use it meaningfully.

- Frame reservation as an access mechanism, alongside realistic skill-building and accommodation.
- Check the current administrative pathway locally before advising on an application or placement.

17.5 IDEAS: important scale, wrong shortcut for ID certification

The acronym is a frequent source of error. **IDEAS** means Indian Disability Evaluation and Assessment Scale. It was developed by the Rehabilitation Committee of the Indian Psychiatric Society in December 2000. IDEAS is designed for disability due to mental disorders and is recommended for disability associated with mental illness. It is publicly available and remains important for psychiatric disability work.

IDEAS rates four functional domains: self-care, interpersonal activities, communication and understanding, and work. Each domain is scored from 0 to 4, from no disability to profound disability, and duration of illness is incorporated into the global score. These features explain why IDEAS may look superficially attractive in an adult with longstanding functional impairment. They do not make it the intellectual-disability certification instrument.

■ **TRAP** **IDEAS must not be substituted for the ID certification route.** Its place is the mental-illness pathway, whereas intellectual-disability certification follows its own prescribed assessment route.

This distinction is more than a technicality. Intellectual disability is a neurodevelopmental condition, and its certification requires an account of adaptive functioning. IDEAS includes illness duration, which answers a different question. Importing it into an ID certificate can create a legally vulnerable assessment and obscure the actual statutory basis of the percentage. Where intellectual disability and mental illness coexist, formulate and treat both conditions, but do not let the presence of psychiatric symptoms alter the prescribed ID percentage route.

17.6 Supported decision-making and the National Trust framework

Certification often brings families to the question they had postponed: who will help the person make decisions in adult life? Begin with the person's own preferences, abilities and participation. The **limited guardianship** model under the Act is offered by the District Court. It is a joint decision-making arrangement based on mutual understanding and trust, confined to a specified period and decision or situation, and intended to operate according to the person's will. This is conceptually different from taking over a person's life because support is convenient for others.

Clinicians should avoid treating guardianship as an automatic consequence of diagnosis or a certificate. Assess the decision at hand, the person's capacity to participate with support, the risks of coercion and the least restrictive way to enable a meaningful choice. Documentation should describe function and support needs rather than make sweeping declarations of incapacity. This approach also keeps the consultation aligned with rights, not merely family anxiety.

The **National Trust for the Welfare of Persons with Autism, Cerebral Palsy, Mental Retardation and Multiple Disabilities Act, 1999** covers four conditions: autism, cerebral

palsy, mental retardation and multiple disabilities. It creates a national body intended to support independent, community-close living and meaningful participation, rather than segregation or passive care. With an adult who has intellectual disability, the clinician must provide information about guardianship and National Trust provisions.

MNEMONIC

SUPPORT before substitution

Start with the person's wishes, identify the decision and the support required, protect against undue influence, and use the least restrictive lawful arrangement.

- Raise future decision-making before an emergency forces the issue.
- Include the adult with intellectual disability in the discussion in a communication-accessible manner.
- Separate financial or property concerns from a global assumption that the person cannot decide anything.

17.7 Documentation, advocacy and protection from procedural harm

Good certification is disciplined advocacy. The final document should make its reasoning auditable: diagnostic conclusion, adaptive assessment, IQ assessment where used, the statutory basis of the percentage, validity and the need for review. Avoid pejorative wording, casual estimates and unexplained contradictions between narrative and score. Discuss the result with the person and family in plain language. A certificate can open doors, but careless language can also follow a person into school, employment and adult services.

Rights under the Act have enforcement mechanisms. A first contravention of the Act or its rules attracts a **fine which may extend to Rs 10,000**, and a subsequent contravention a **fine of not less than Rs 50,000 and up to Rs 5 lakh**. Neither tier carries imprisonment, and the first-tier figure is a ceiling rather than a flat penalty. Fraudulently availing a benefit meant for persons with benchmark disabilities is a separate and graver offence, punishable with imprisonment which may extend to 2 years, or fine which may extend to Rs 1 lakh, or both. A separate group of offences of atrocities carries imprisonment of not less than 6 months and up to 5 years, and with fine, and their elements are examinable precisely because dropping them broadens the offence. Insult or intimidation qualifies when it is intentional, done with intent to humiliate, and in a place within public view. The sexual-exploitation limb requires the offender to be in a position to dominate the will of a woman or child with disability and to use that position to exploit her. The section runs to several further limbs, each with its own elements, and they are worth reading in the Act itself rather than in summary: a paraphrase that drops an element is precisely how these offences get overstated. Special Courts are designated in each district to handle violations.

The psychiatrist is not expected to become the family's lawyer. The clinical role is to recognise when rights may be at stake, document the relevant functional facts clearly, avoid false reassurance and direct the family toward appropriate legal or administrative assistance. For

broader assessment, service planning and protection issues, consult the Child psychiatry: assessment, treatment, services and protection notes. The practical aim is simple: certification should increase access, participation and safety without becoming a mechanism for unnecessary exclusion.

Must remember: In intellectual-disability certification, diagnose clinically, confirm intellectual functioning with the Binet-Kamat Test or Malin’s Intelligence Scale for Indian Children, and calculate the statutory percentage from VSMS-based adaptive functioning - not from IDEAS, and not from the IQ score, which does not enter the percentage at all.


Autism: concept, causes and criteria

18 · Historical evolution (Kanner, Asperger, DSM-IV subtypes to spectrum)

Why this history matters. Autism did not move from obscurity to its current spectrum formulation simply by acquiring a new name. Each classificatory change reflected a different answer to a clinical question: is the presentation a distinct childhood syndrome, a developmental disorder with several recognisable patterns, or a dimensional condition whose boundaries cannot be reliably divided into neat subtypes? For the clinician, this history explains why older notes use unfamiliar terms, why families may still identify with a former label, and why current assessment must not be constrained by obsolete diagnostic partitions.

The examinable spine is therefore not a recital of dates. It is the movement from carefully described clinical prototypes, through a broad pervasive-developmental-disorder framework, to a single spectrum diagnosis. That movement also records an important correction in psychiatric thinking: explanatory models have to be accountable to observation and evidence, particularly when they risk locating blame in parents.

1943 FIRST PAPER	11 CHILDREN DESCRIBED	1944 PARALLEL ACCOUNT	2013 SPECTRUM SHIFT
---------------------	--------------------------	--------------------------	------------------------



18.1 From descriptive observation to a clinical syndrome

The starting point is the clinical description associated with **Leo Kanner, an Austrian-born child psychiatrist working at Johns Hopkins University**. He is credited with the first clinical portrayal of autism in the scientific literature. The importance of that contribution lies in the act of delineation: a set of children who might otherwise have been subsumed under broad and unhelpful psychiatric categories were described as sharing a distinctive developmental pattern. Good historical reading is not hero worship. It is an appreciation that a diagnostic entity begins when clinicians can show, with enough precision, what belongs together and what does not.

The paper, *Autistic disturbances of affective contact*, concerned a series of **11 children** accrued over 5 years. It appeared in *The Nervous Child*, volume 2, pages 217-250. The title itself is instructive. Its language belongs to its period, but it directs attention to a disturbance in the child's way of relating rather than merely to disruptive conduct or low measured ability. That emphasis helped establish autism as a problem of developmental organisation and social relatedness, rather than a label for any child who was withdrawn, unusual, or difficult to manage.

A later follow-up is a useful corrective to retrospective romanticising of early accounts. Kanner's 1971 follow-up traced 9 of the original 11 children into adulthood; only two were eventually employed, and they remained single and lived with their parents. These observations should not be converted into a modern prognostic rule. They arose from a small historical cohort and a very different service context. Their value is conceptual: a diagnostic history should remain connected to the life course, participation, support, and family circumstances of the people being described.

■ **RULE** Read the early descriptions as clinical prototypes, not as a checklist that limits whom autism can include. A prototype establishes a recognisable pattern; later classification must test whether its boundaries remain valid in wider clinical populations.

18.2 Asperger's parallel account and its later influence

Independently of Kanner's work, **Hans Asperger** published an account based on **4 cases**. In his 1944 paper, *Die "Autistischen Psychopathen" im Kindesalter*, he used the term "autistic personality disorder". The parallel descriptions matter because they prevent an overly simple origin story. Autism was not manufactured by a later committee out of a single author's observations. Different clinicians, working separately, identified related developmental phenomena, while giving different weight to their most striking features.

The feature that especially distinguished that account was attention to **extraordinary skills in particular cognitive domains**. This is often simplified into a stereotype of universal talent, which is clinically unhelpful. The historical point is narrower: ability can be strikingly uneven, and the presence of a particular strength does not erase social-developmental difficulty or establish a separate disease entity. Conversely, the absence of a striking skill never argues against autism. A mature formulation should describe the person's actual profile rather than infer ability from a historical label.

The work was published in German and was not widely disseminated in English-speaking settings until nearly 40 years later. **Lorna Wing**, in 1981, introduced these findings to the

English-language literature and described Asperger's syndrome. This delay helps explain why the label became prominent after many clinicians had already learned the original account of autism. It also explains why the label acquired cultural and personal significance for some patients and families. Respecting that significance does not require preserving a classification that later evidence found difficult to delimit consistently.

- Historical labels may illuminate how a person and family first understood a developmental profile.
- They should not be treated as a shortcut to current assessment or assumptions about cognition, language, education, or independence.
- When reviewing an old record, translate the label into a current description of developmental history, functioning, and support needs.

18.3 Early labels, aetiological error, and scientific correction

An influential early label emerged when **Kanner and Leon Eisenberg, in 1956, used “early infantile autism” and emphasised obsessive insistence on sameness and extreme self-isolation**. The label concentrated attention on early developmental presentation and on behavioural patterns that clinicians continue to recognise. Yet it also shows why diagnostic terminology should not be mistaken for a complete explanation. A name can draw attention to a pattern while still carrying assumptions about timing, causation, or typical presentation that later evidence will revise.

One of the most damaging examples was the **“refrigerator mother” theory, furthered in the 1960s by Bruno Bettelheim**. It interpreted autism through a psychogenic account of parenting and emotional deprivation. Its clinical legacy is not merely historical embarrassment. It can lead to guilt, adversarial relationships with caregivers, and neglect of the child's developmental and practical needs. The lesson for a psychiatrist is to separate careful inquiry about family relationships, which remains clinically relevant for every child, from unsupported causal attribution. Families require collaborative assessment, not retrospective blame.

Empirical work displaced this theory by providing **evidence for biological and neurological foundations, while comparative studies distinguished autism from childhood schizophrenia through associated symptoms, phenomenology, age of onset, and family history**. This was a methodological advance as much as an aetiological one. It required investigators to compare syndromes rather than rely on surface impressions, and to ask whether apparently similar social withdrawal had the same developmental meaning. The distinction remains clinically valuable: resemblance in a single symptom is not enough to establish diagnostic equivalence.

■ **TRAP Examination trap:** do not present the refrigerator-mother account as an historical “cause” that later became less important. It was a discredited psychogenic theory, not a competing aetiology to be balanced against contemporary evidence.

For present practice, history supports a disciplined stance. A developmental diagnosis does not make family context irrelevant, nor does it justify reducing the child to biology. It means that causal claims should match evidence, while formulation still attends to caregiver stress, communication, schooling, access to services, stigma, and the child's strengths. The causes of autism and the clinical features used for diagnosis are addressed elsewhere in this chapter; history here teaches why those domains must not be confused.

18.4 DSM-III and the pervasive developmental disorder framework

Autism entered a formal diagnostic manual in **1980**, when DSM-III used the label “infantile autism”. Formal recognition mattered because it placed the presentation within a reproducible diagnostic framework rather than leaving it dependent on individual clinical tradition. It also created the possibility that criteria could be studied, criticised, revised, and compared across settings.

The broader home created for it was **pervasive developmental disorder (PDD)**, a term developed specifically for DSM-III. It denoted pervasive and severe impairment across basic functions, including socialisation and communication. The word “pervasive” captured an attempt to distinguish a broad developmental organisation from an isolated symptom. Its limitation became clearer over time: a useful umbrella can inadvertently suggest that the conditions beneath it are naturally separate, even when the boundaries between them prove unstable.

The PDD framework nevertheless performed an important transitional task. It organised clinical observation around developmental patterns and made room for presentations that did not resemble a narrow prototype. It also created a common language for research and services. Candidates should understand this as a stage in nosology, not as a rival to the current diagnosis. The historical PDD term is still encountered in case records, educational reports, and older research, so it must be recognised and translated rather than dismissed as meaningless.

There is a broader lesson in this transitional category. Classification is most useful when it makes a clinician ask better questions, not when it supplies a prematurely final answer. The PDD framework encouraged clinicians to look across domains of development, but its named divisions could tempt them to treat a label as a complete account of the person. The later spectrum formulation retained the need for careful description while moving the centre of gravity from deciding which box fits best to explaining the individual's pattern. That is a more demanding task: it requires longitudinal history, direct observation, collateral information, and attention to functioning in ordinary settings.

The key historical transition is from separately named pervasive developmental disorders to one autism spectrum diagnosis: the clinical task moved from choosing a former subtype to applying a common framework precisely.

18.5 DSM-IV: expansion into named subtypes

The DSM-IV and DSM-IV-TR PDD category contained exactly 5 disorders: autistic disorder, Asperger's syndrome, Rett's syndrome, childhood disintegrative disorder, and PDD-NOS. This

scheme is worth knowing because it shaped generations of clinical records and examination questions. It was also an attempt to represent heterogeneity, especially differences in developmental course and the degree to which an individual resembled the original prototypes. The weakness was not that heterogeneity was unreal; it was that named categories did not reliably carve that heterogeneity at its natural joints.

DSM-IV / DSM-IV-TR SUBTYPE	DIAGNOSTIC CODE
Autistic disorder	299.00
Asperger's disorder	299.80
PDD-NOS, including atypical autism	299.80
Rett's disorder	299.80
Childhood disintegrative disorder	299.10

Do not let the table mislead you into seeing the former codes as a clinical hierarchy. Codes record a classification decision; they do not tell you how a person communicates, what repetitive patterns are present, how development unfolded, or which supports are needed. The full diagnostic criteria, severity framework, and specifiers of current ASD are addressed elsewhere in this chapter. Here, the important point is that a subtype name in an older file is historical data to be interpreted, not a final formulation.

- Do not infer that former subtypes represented neatly separated biological entities.
- Do not convert a former subtype automatically into a contemporary severity statement.
- Do preserve the original developmental account, because it may contain useful information that a code alone cannot carry.

18.6 DSM-5: from subtype boundaries to a spectrum

The decisive revision came with **DSM-5 in 2013. Autistic disorder, childhood disintegrative disorder, Asperger's syndrome, and PDD-NOS were reconceptualised as a single autism spectrum disorder.** This was not a claim that all autistic people are alike. It was a claim that the prior labels did not consistently identify separate disorders, whereas variation could be described more meaningfully within a shared diagnostic framework.

The change also altered the architecture of symptoms: the 3 DSM-IV-TR symptom categories became 2 DSM-5 categories, and the item architecture beneath them was rebuilt. DSM-IV-TR required a total of 6 or more items across its 3 categories, with at least 2 drawn from the social-interaction category, which itself listed 4; DSM-5 carries 3 items in the single social-communication and social-interaction category. The threshold rule is worth reading twice, because it is widely misquoted as though the six were six social items: the six was the total across all three categories, and the social category alone never contained six. These are historical contrasts, not a substitute for learning current criteria. The critical conceptual shift was to bring social communication and social interaction together, while retaining restricted and repetitive

patterns as the other major domain. The diagnostic-criteria discussion should be used for the present rule set; history explains why it was reorganised.

The onset wording changed as well. DSM-IV-TR required onset before age 3, whereas DSM-5 specifies early development. This prevents an arbitrary historical cut-off from overriding a careful developmental history. It does not mean that onset history has become unimportant. Rather, the clinician has to establish whether the overall developmental pattern is compatible with ASD and whether later recognition reflects changing social demands, masking, missing early history, or another explanation.

- **RULE** **Spectrum does not mean vague.** It replaces unstable subtype labels with a common diagnostic framework while requiring a more precise individual description of developmental history and current functioning.

18.7 Rett syndrome, coding legacies, and reading old records safely

The spectrum consolidation had an explicit exception. **Rett syndrome was not retained as a distinct autism-spectrum category; it is cited under the associated-condition specifier as an example of a known genetic condition.** This is a useful guard against a common historical error: assuming that every condition previously housed within PDD was simply renamed ASD. The detailed clinical account of Rett syndrome and childhood disintegrative disorder belongs with their dedicated discussion; the classification point here is that historical placement and present diagnostic status are not the same thing.

The manual pairs that reclassification with a differential instruction, set out where each of those syndromes is discussed in full. The classification principle is what belongs here, and it generalises: a symptom that appears during a particular phase of another condition must not be treated as sufficient evidence for a lifelong co-occurring diagnosis. Diagnostic overlap calls for full criteria, developmental context, and longitudinal reasoning.

Older ICD-10 documentation may use the following labels. The list is useful for record translation, but it must not be read as a DSM-IV row-by-row conversion.

ICD-10 CODE	LABEL PRINTED IN THE HISTORICAL PDD LISTING
F84.0	childhood autism
F84.1	atypical autism
F84.2	Rett's syndrome
F84.3	childhood disintegrative disorder
F84.4	overactive disorder associated with mental retardation and stereotyped movements
F84.5	Asperger syndrome
F84.8	other pervasive developmental disorders
F84.9	pervasive developmental disorders, unspecified

For the ICD-11 ASD stem, the back-mapping permits F84.0 childhood autism, F84.1 atypical autism, or F84.5 Asperger syndrome; use F84.1 when onset before age 3 is unclear, and use F84.5 regardless of onset when there is no general impairment in intellectual functioning or functional language. This is a coding bridge, not a licence to reintroduce obsolete diagnostic distinctions into the clinical formulation.

Finally, modern ASD encompasses historical terms that will appear in legacy material:

- early infantile autism, childhood autism, and Kanner's autism
- high-functioning autism, atypical autism, and PDD-NOS
- childhood disintegrative disorder and Asperger's disorder

GOOD TO REMEMBER

When encountering a historical label, first establish which classification system was used, then recover the descriptive evidence behind it, and finally apply the current framework without erasing the person's history. Taxonomy should improve clinical understanding, not replace it.

19 · Diagnostic criteria of ASD (DSM-5 dyad, ICD-11)

The diagnostic task is to recognise a pattern, not to count isolated oddities. Autism spectrum disorder is formulated from a persistent developmental pattern that is organised around **two** core domains. The value of the formal criteria is that they force the clinician to establish both the social-communication disturbance and the restricted or repetitive pattern, then to ask whether the pattern began developmentally and matters in the person's present life. This protects against diagnosing autism from one striking mannerism, a narrow interest, or social discomfort alone.

The criteria are best understood as a clinical argument. Symptoms must be described in their developmental context, linked to real-world functioning, and weighed against the person's general developmental level. The classification language is concise; the clinical work is not. A good answer therefore states the diagnostic structure first and then explains why each element is necessary. This section addresses the criteria in DSM-5-TR and ICD-11, before showing how their different formats lead to the same disciplined diagnostic reasoning.

2 DSM CORE DOMAINS	3 CRITERION A ELEMENTS REQUIRED	at least 2 CRITERION B ELEMENTS REQUIRED	6A02 ICD-11 PARENT STEM
------------------------------	--	---	-----------------------------------




Fig 21.15 The DSM-5-TR autism dyad, and the two counting rules inside it. Two symptom domains, two different rules. Criterion A needs all 3 of its 3 sub-criteria, so one intact sub-criterion defeats it; Criterion B needs any 2 of its 4, and which 2 does not matter. Criteria C, D and E add onset, impairment and the not-better-explained rule against intellectual developmental disorder. Specifiers hang off the diagnosis, not the counts, where language and severity sit, outside the criteria set. ICD-11 6A02 has the same two domains and no counting rule, so these counts are DSM-5-TR's alone. No onset age is drawn: DSM-5-TR attaches none to the early developmental period, and the specifier axes are listed, never numbered, because it does not state how many there are.

19.1 The diagnostic architecture: a dyad with developmental and functional anchors

The DSM-5-TR dyad has one domain concerning social communication and social interaction, and another concerning restricted, repetitive patterns of behaviour, interests, or activities. The dyad is not merely a shorter checklist. It requires the examiner to decide whether social difficulty is part of a broader reciprocal-interaction disturbance and whether the behavioural pattern has the characteristic quality of repetition, inflexibility, restriction, or unusual sensory engagement.

A useful way to read the criteria is vertically as well as horizontally. Vertically, the clinician first establishes the core phenotype, then asks about developmental onset, present impairment, and competing explanations. Horizontally, each core domain must be examined across ordinary contexts rather than inferred from a single consultation. A person may use rehearsed social behaviours in a structured setting and still have difficulty when reciprocity, flexibility, or contextual understanding are demanded. Conversely, a single unusual habit without the wider pattern does not establish the diagnosis.

This structure also explains why a symptom inventory is insufficient. Checklists are useful prompts for enquiry, but their items are not interchangeable. An affirmative response about a repetitive act does not repair missing evidence from the social domain, and a history of social discomfort does not establish the repetitive domain. The clinician must decide whether each observation belongs to the required construct, whether it is persistent, and whether the observations form one developmental syndrome. That is the safeguard against both overinclusive diagnosis and the dismissal of a coherent but subtle presentation.

■ **RULE** **Diagnostic rule:** establish the complete pattern across both core domains before using onset, impairment, and exclusion criteria to confirm that it is autism spectrum disorder.

The printed DSM diagnosis carries the code F84.0. In an examination answer, however, the code is less important than showing the logic of the dyad. The examiner should be able to see that the candidate understands why a social-communication presentation without the required restricted or repetitive pattern is not simply labelled autism by default.

19.2 Criterion A: social communication and social interaction

Criterion A concerns deficits that are persistent and span the social-communication field. DSM-5-TR requires **all three** listed components. This matters because conversational difficulty, poor eye contact, or social isolation in isolation can arise for many reasons. The criterion asks for a coherent disturbance in reciprocity, in the use of nonverbal communication for social purposes, and in relationships.

- **Deficits in social-emotional reciprocity** concern the give-and-take of social exchange. The clinical question is whether the person can initiate, respond to, and sustain a reciprocal interaction in a manner appropriate to the context.
- **Deficits in nonverbal communicative behaviours used for social interaction** concern how nonverbal signals are used to regulate and share interaction, rather than a superficial judgement about one gesture or one gaze behaviour.

- **Deficits in developing, maintaining, and understanding relationships** require attention to the person's grasp of social expectations, adjustment of behaviour to context, and capacity to participate in relationships.

The phrase “social communication” should not be reduced to spoken language. A fluent speaker can have marked difficulty with reciprocity, social interpretation, or relationships, while a person with limited speech may show a clear wish to engage and share. The diagnostic question is how social communication functions between people. **Language impairment is not itself a core DSM-5-TR criterion**; its specifier framework is addressed in the severity-level and specifier discussion in this chapter. Detailed language phenomenology belongs with the clinical features of ASD.

Clinical interpretation. Criterion A is often easiest to misread when an observer relies on a socially compelling moment during an interview. The purpose is not to decide whether a person can make contact at all; it is to understand the consistency, flexibility, reciprocity, and context-sensitivity of social communication over development. A single competent performance, particularly in a familiar or highly structured interaction, is not by itself a substitute for that broader account.

For the same reason, Criterion A should not be phrased as a personality judgement. Reserve descriptive language for observable interaction and its functional meaning. Ask how the person shares attention, responds when another person changes the topic, reads the interpersonal situation, and adapts communication to the relationship. Such questions convert a loose impression of being “socially awkward” into evidence that can be placed, or not placed, within the criterion. They also make it easier to explain the reasoning to the person and family without reducing the assessment to a label.

19.3 Criterion B: restricted and repetitive patterns

Criterion B captures a style of behaviour and interest that is repetitive, circumscribed, inflexible, or unusually related to sensory experience. DSM-5-TR requires **at least two** of the listed manifestations. The threshold deliberately prevents a single preference, hobby, or repetitive action from carrying the diagnosis. What counts is the pattern's persistence, context, intensity, and relationship to the rest of the clinical picture.

The manifestations include stereotyped or repetitive motor movements, use of objects, or speech. The point is not simply that behaviour is repeated. Repetition must be interpreted as part of the characteristic pattern, rather than as an isolated self-soothing act, play preference, or response to a transient stressor.

A further manifestation is insistence on sameness, inflexible adherence to routines, or ritualized patterns of verbal or nonverbal behaviour. Here, the clinical task is to ask what happens when an expected sequence changes. The diagnostic relevance lies in the rigidity and the way the need for predictability shapes functioning, not in whether a household happens to have routines.

The domain also includes highly restricted, fixated interests that are abnormal in intensity or focus. A strong interest should be evaluated for its quality and consequences, not judged by whether its topic seems unusual to the clinician. An ordinary subject can become diagnostically relevant when engagement is unusually narrow or dominant; an unusual subject can be entirely non-pathological when it is flexible and proportionate.

Finally, DSM-5-TR includes **hyper- or hyporeactivity to sensory input, or unusual interest in sensory aspects of the environment**. This places sensory experience within the restricted and repetitive domain. It does not mean that sensory sensitivity alone is diagnostic. Its value is greatest when it helps explain a broader pattern of behaviour, distress, avoidance, fascination, or difficulty adapting to ordinary environments.

MNEMONIC

Memory spine: A is all, B is a selected set

Establish every social-communication element in A, then establish the required threshold within B. Use this as a recall aid only after understanding what the observations mean clinically.

- **TRAP Examination trap:** do not diagnose ASD from social-communication deficits alone. The restricted and repetitive domain is required, and it must be demonstrated rather than assumed.

19.4 Criteria C to E: timing, impact, and developmental explanation

The core phenotype is necessary but not sufficient. Criterion C requires that symptoms are present in the **early developmental period**. Recognition may occur much later because demands become more complex or because a person has learned strategies that conceal difficulty. The criterion is therefore about developmental presence, not the age at which a family, school, or clinician first names the concern. A late recognition narrative should prompt careful developmental reconstruction, not an automatic rejection of ASD.

Criterion D requires **clinically significant impairment** in social, occupational, or other important areas of current functioning. Impairment is the bridge from descriptive traits to a disorder. It should be considered in the person's actual environment, including the effort needed to maintain functioning and the consequences when familiar supports are absent. The clinician should avoid both extremes: treating every difference as disorder, or dismissing substantial hidden burden because outward performance appears adequate.

Criterion E says that the presentation must not be better explained by intellectual developmental disorder or global developmental delay. It also permits co-occurring diagnoses when **social communication is below that expected for general developmental level**. This is a comparative judgement, not a shortcut based on an intelligence score. The issue is whether the social-communication profile is disproportionately affected relative to the person's overall developmental functioning. Intellectual disability and its assessment are covered in the relevant sections of this chapter.

- **Timing:** establish a developmental history that can account for the current pattern, including why it became visible when it did.
- **Impact:** describe what the pattern changes in daily participation, relationships, education, work, or other important roles.
- **Explanation:** decide whether the social-communication disturbance exceeds what would be expected from the person's general developmental level.

The transition note is clinically important: an individual with a well-established DSM-IV diagnosis of autistic disorder, Asperger's disorder, or PDD-NOS should receive the diagnosis of autism spectrum disorder. This is a diagnostic transition rule, not an invitation to recreate older subgroup boundaries in current formulation. For a person with marked social-communication deficits who does not meet the full ASD criteria, the DSM directs that they should be evaluated for social (pragmatic) communication disorder. The full differential is considered elsewhere in this chapter.

19.5 ICD-11: essential features and a clinically judged threshold

ICD-11 places ASD under the parent stem 6A02. Its essential features have two required symptom domains, structurally paralleling the DSM dyad. The practical message is the same: social-communication disturbance and restricted or repetitive patterns are both indispensable to the syndrome.

The important structural contrast is that ICD-11 does not set a numerical sub-criterion threshold within either domain. Instead, manifestations are illustrative rather than a required count. This calls for disciplined clinical judgement, not diagnostic laxity. The clinician must still establish that the required domains are convincingly represented and persistent. A list of examples is not a licence to infer a domain from a vague or isolated concern.

When working from ICD-11, document the evidence that makes each domain essential rather than simply reproducing several examples. This keeps the record auditable: another clinician should be able to see why the pattern was judged to meet the domain, how it has persisted developmentally, and how it affects functioning. The same discipline is useful when DSM criteria are used. Counting is a decision aid, not a replacement for a coherent account of the person's development and current life.

The ICD onset requirement is that onset occurs during the developmental period, usually becoming apparent early, although characteristic symptoms may become fully manifest only when social demands exceed the person's capacities. Its functional requirement is significant impairment in important areas of functioning; diagnosis remains appropriate when adequate functioning is achieved only through exceptional effort. This wording is especially useful for presentations in which compensation can obscure the burden of the condition.

ICD-11 also notes that a true regression pattern may typically begin between **1 to 2 years of age**, while loss of previously acquired skills is rarely observed above **3 years of age**. This is a narrow

classification observation, not a substitute for the clinical assessment of regression or its differential diagnosis.

19.6 DSM-5-TR and ICD-11: compare the logic, not merely the wording

DIAGNOSTIC AXIS	DSM-5-TR	ICD-11	CLINICAL CONSEQUENCE
Core structure	Two core domains	Two essential symptom domains	Both require a dyadic phenotype rather than a single-domain formulation.
Within-domain threshold	Every Criterion A component is required; Criterion B requires its stated minimum.	No required sub-criterion count is specified.	DSM is more operationally counted; ICD relies more explicitly on judgement about the essential pattern.
Developmental timing	Symptoms are present in the early developmental period.	Onset occurs during the developmental period.	Recognition can be delayed without converting a developmental condition into an adult-onset one.
Functional significance	Clinically significant current impairment is required.	Significant impairment remains relevant despite exceptional compensatory effort.	Assess current consequences and effort, not symptoms in abstraction.

The classifications should therefore not be treated as rival definitions. They are two formats for the same central clinical problem: determining whether a developmental social-communication pattern is accompanied by the required restricted or repetitive pattern and causes meaningful difficulty. DSM makes the internal threshold more visible. ICD makes the need for judgement in a variable phenotype more visible. In either system, the best formulation is evidence-based, developmental, and functionally anchored.

19.7 A disciplined way to write and defend the diagnosis

GOOD TO REMEMBER

In practice, begin by naming evidence under each core domain. Then state the developmental course, the present functional impact, and the developmental comparison required when intellectual disability or global developmental delay is relevant. This sequencing prevents the common error of beginning with a favourite label and searching selectively for confirming behaviours.

A concise formulation can therefore be written as a chain of reasoning: the person has evidence for the social-communication domain; evidence for the restricted or repetitive domain; a developmental history compatible with the condition; current functional consequences; and no more complete developmental explanation for the social-communication profile. If any link is weak, the right response is to clarify that link rather than to stretch another criterion. This

approach is especially valuable when the presentation is compensated, when history is incomplete, or when more than one developmental difficulty is present.

Do not turn the criteria into a proxy severity scale. Severity levels and accompanying intellectual or language impairment are specifiers, not replacements for the core diagnostic argument. Likewise, screening and diagnostic instruments may organise information, but they do not remove the need to establish the criteria. Their selection and use are addressed in the assessment section of this chapter.

ASD requires the dyad, developmental presence, clinically meaningful impact, and a developmental explanation that does not account for the social-communication profile more fully.

The final formulation should make its reasoning transparent. It should show why the observed pattern belongs together, why it is developmental, why it matters now, and why the chosen diagnosis is more accurate than a partial description. That is the difference between reproducing a checklist and using a diagnostic classification responsibly.

20 · Epidemiology and changing prevalence of ASD

Epidemiology is more than a prevalence figure. In autism spectrum disorder, the number reported by a survey depends on whom it reaches, how cases are found, which diagnostic concept is used, and whether a record, a service register, or direct assessment establishes the diagnosis. The examination task is therefore not to recite one figure as though it were universal. It is to state the population and method to which a figure applies, distinguish occurrence from detection, and explain why apparently conflicting estimates may all be credible within their own settings.

Autism is diagnosed across the life course, but ascertainment is shaped by developmental surveillance, referral pathways, educational provision, clinical expertise, and families' ability to reach assessment. These influences make epidemiology clinically relevant. They determine which children come to attention, which adults remain unidentified, and how a service should interpret a rising caseload. Prevalence describes the burden recorded in a defined population; it should not casually be converted into a claim about causation or a proof that the underlying incidence has changed.

20.1 Reading an ASD prevalence estimate

Prevalence is the proportion of a defined population meeting a case definition at a stated time or over a stated period. For ASD, it is an output of both the condition in the population and the process used to recognise it. A direct, structured assessment can identify people who have never entered services, whereas an administrative dataset captures only those who have acquired a diagnostic label. Neither approach is automatically superior in every setting: each answers a somewhat different question.

A useful distinction is between a population survey, a clinical series, and a service count. A clinic denominator is strongly influenced by referral, affordability, and local diagnostic practice. A

school or disability register may be especially sensitive to eligibility rules. A population study can reduce some of these effects, but still depends on participation and on the screening and diagnostic pathway. Thus, when comparing papers, the candidate should ask: who was sampled, what constituted a case, and how was that case established?

Case definition is equally important. Research that asks whether a person meets a current diagnostic formulation is not directly equivalent to a historical study that used a narrower formulation. Nor is a parent-reported label equivalent to an assessment made after direct observation and developmental history. The point is not to rank methods in the abstract. It is to understand the direction in which each method may shape a result and to avoid comparing unlike estimates as though they measured precisely the same construct.

For clinical readers, this also explains why a service's experience can diverge from published population data. A tertiary service may receive people with complex presentations, delayed recognition, or accompanying needs that make referral more likely. A community survey may capture people whose needs have not brought them to a specialist. Both can be valuable sources of information, but neither should be used to invalidate the other without considering its sampling frame. Epidemiology becomes clinically useful when it trains that habit of interpretation.

■ **RULE** **Never quote an ASD frequency without naming its population, period, and ascertainment pathway.** The same label can yield different observed frequencies when the pathway into the numerator changes.

Ascertainment means the process by which potential cases are found and verified. It is not a minor methodological footnote. A wider net may identify people whose presentations were previously overlooked; an altered threshold may change who qualifies; and better access to assessment may turn an unmet need into a recorded diagnosis. These processes can change measured prevalence even when the underlying population risk has not changed.

- State whether the estimate is local, national, or multinational.
- Identify whether it is a survey estimate, a diagnosed-rate estimate, or a service-derived estimate.
- Describe a temporal increase as a change in **measured prevalence** unless the evidence separately establishes an increase in incidence.

20.2 Contemporary frequency: use the denominator carefully

Published figures support a broad conclusion: ASD is not rare in contemporary clinical and public-health practice. They do not support treating one jurisdiction's figure as the world's figure. The DSM-5-TR describes a reported frequency in the United States of **between 1% and 2%**. In the same source, the median global prevalence is **0.62%**, while prevalence across non-United States countries has approached 1%. These are estimates from differently assembled evidence, not interchangeable constants.

1-2% UNITED STATES REPORTED FREQUENCY	0.62% MEDIAN GLOBAL PREVALENCE	approaching 1% NON-UNITED STATES REPORTS	3:1 WELL-ASCERTAINED EPIDEMIOLOGICAL SEX RATIO
--	---	---	--

The Kaplan & Sadock's Comprehensive Textbook of Psychiatry gives a current estimate of about 69 per 10,000, or about 1 in 145 children. Lewis's Child and Adolescent Psychiatry describes 6.6 per 1,000 as a reasonable overall estimate. The immediate lesson is not to calculate a false reconciliation between these reports. Rather, each estimate needs its source and denominator attached when it is quoted.

IN INDIA

An international or United States surveillance number may be useful as context, but it must not be relabelled as an Indian prevalence estimate. The Indian Psychiatric Society guideline cites a United States surveillance estimate of 1 in 68; its provenance must travel with the number. A responsible answer acknowledges the need for locally representative ascertainment rather than importing a denominator from another health and education system.

20.3 Sex and gender: distinguish epidemiology from diagnosis

Sex ratio is often quoted as a compact epidemiological fact, but it carries an important methodological warning. In well-ascertained epidemiological samples, the global male:female ratio is 3:1. That figure should not be conflated with the ratio among people who receive a diagnosis. The DSM-5-TR states that ASD is diagnosed between three and four times more often in males, while the ICD-11 Clinical Descriptions and Diagnostic Requirements describes males as diagnosed at a 4:1 male:female ratio.

The difference between a diagnostic ratio and an epidemiological ratio matters because diagnosis is a social and clinical event. Referral expectations, recognition of less stereotyped presentations, language and cognitive profile, and co-occurring difficulties can all shape who crosses the threshold into a dataset. Therefore, a male predominance in a clinical register must not be interpreted automatically as a direct measure of the sex distribution of autistic traits in the wider population.

Other textbook summaries similarly report a male excess, but their scope should be retained. Kaplan & Sadock's Comprehensive Textbook of Psychiatry reports on the order of three to five boys for every girl in clinical and epidemiological samples, and notes that the predominance is greater among those with higher cognitive ability. Lewis's Child and Adolescent Psychiatry reports that boys are affected between three and four times more often. These formulations should be presented as source-specific descriptions, not as permission to under-recognise ASD in girls or women.

■ **TRAP Examination trap:** a diagnosed male:female ratio is not automatically a population prevalence ratio. Name the numerator before interpreting the ratio.

20.4 Why recorded prevalence has risen

The historical contrast is striking, but the correct inference is nuanced. Early epidemiological studies described autism as roughly 1 per 2,000, whereas current estimates are much higher. This temporal pattern establishes that the recorded frequency has changed. It does not, by itself, identify why it changed. Trends are especially difficult to interpret when the diagnostic category, awareness, service incentives, and methods of finding cases evolve together.

REPORT	FIGURE AND INTENDED USE	INTERPRETIVE BOUNDARY
Early studies	Roughly 1 per 2,000	Historical baseline, not a current estimate.
Kaplan & Sadock	About 69 per 10,000, or about 1 in 145 children	Contemporary textbook estimate.
Lewis	6.6 per 1,000	Reasonable overall estimate in that text.
Indian Psychiatric Society guideline	1 in 500 historically, compared with 1 in 68	Both figures are historical comparison and United States surveillance context, not Indian population prevalence.

Secular increase refers to an apparent change across time. It may reflect a real change in occurrence, an altered measurement process, or both. Lewis interprets the **apparent rise** through evolving diagnostic concepts, broader recognition of the spectrum, and access to relevant services, rather than as proof that population occurrence has increased. This is a disciplined reading of time trends: the conclusion should remain proportionate to what the underlying studies can establish.

The evidence base has expanded substantially. Kaplan and Sadock refers to over 60 epidemiological studies. More studies improve the visibility of variation and methodological limitations; they do not erase those limitations. An increased research base should make the clinician more careful about denominators, not more casual about generalisation.

20.5 Mechanisms that inflate or alter measured frequency

Diagnostic broadening or change in diagnostic practice can alter measured frequency. When a category is conceptualised differently, people previously placed outside it, or given another developmental label, may now meet the contemporary definition. Awareness also matters: families, teachers, and clinicians who recognise a broader range of presentations are more likely to refer and document concerns. Service availability can have a similar effect, because visible assessment and educational support make diagnosis more feasible and more relevant to a family.

Diagnostic substitution is a related, but more specific, phenomenon. In the United States in particular, a single diagnostic label is commonly used to identify students for service eligibility. As awareness rose, schools and parents became more likely to use the autism label; estimates can consequently be inflated when service records, rather than direct assessment of children, establish caseness. This is not a claim that every increase is artificial, nor should this United States-specific mechanism be exported uncritically to all countries.

Kaplan and Sadock likewise regards **the measured rise** as partly shaped by evolving diagnosis, professional practice, educational policy, and public recognition; higher rates often occur where awareness and services are more available. It leaves the possibility of a true incidence increase unresolved. That restraint is the correct clinical posture. A service observing more diagnosed children may be seeing a mixture of changing need, changing recognition, and changing access.

MNEMONIC

FRAME the rising count

Find out how cases were found; **R**everview the diagnostic rule; **A**sk about awareness and access; **M**ap service and educational incentives; **E**xpress uncertainty about true incidence.

- Do not call a rise an epidemic merely because a surveillance figure rose.
- Do not dismiss all increase as artefact merely because ascertainment has improved.
- State which mechanisms are supported by the source and which remain unresolved.

20.6 Ethnoracial, social, and cultural interpretation

Within United States child data reported in the DSM-5-TR, prevalence is lower in African American children at 1.1% and Latinx children at 0.8% than in White children at 1.3%, even after socioeconomic resources are taken into account. This is an observed comparison in that source. It should not be rewritten as established proof that the difference is wholly caused by underdiagnosis or differential ascertainment. The same source says reported prevalence may be affected by misdiagnosis, delayed diagnosis, or underdiagnosis, which is a cautious possibility rather than a completed causal explanation.

Ascertainment bias is the distortion introduced when some people are more likely than others to be identified, referred, or recorded. It is a useful hypothesis when interpreting group differences, but it is not a substitute for evidence. A good answer separates the observed distribution from proposed explanations and avoids converting a plausible clinical concern into a settled epidemiological conclusion.

Social-class arguments require the same historical discipline. **Kanner's original observation** involved highly achieving, high-social-status parents, but Kaplan and Sadock regards the sample as likely biased by case ascertainment and notes that epidemiological studies have generally not found such an association. The lesson extends beyond ASD: impressions formed from a selective clinic population should never be mistaken for a population association.

20.7 Applying epidemiology in clinical and service work

What the clinician should do with a rate. Epidemiology should improve case-finding and service planning, not replace individual assessment. A family does not become more or less likely to need a careful developmental evaluation because the clinician has memorised a national estimate.

Conversely, a service cannot plan workforce, assessment capacity, or educational liaison responsibly if it assumes that a historical clinic count captures all people who may need support.

GOOD TO REMEMBER

Treat an apparent rise in caseload as an audit question, not a reflex conclusion. Check referral, assessment pathways, education or disability processes, age groups, and the catchment population. This helps distinguish increased recognition from a changed denominator, so planning is based on demand visible to the service.

At the level of a research appraisal, ask whether the study has separated screening from confirmation and whether non-participants might differ from participants. Ask whether a diagnostic record had to pre-exist for inclusion, and whether the study would identify people with a subtle or previously unrecognised presentation. Such questions are transferable across settings. They also keep the clinician from treating an attractive headline figure as more precise than the study design permits.

When counselling families, use population data with humility. Explain that frequency figures vary across settings and methods, and that an epidemiological association cannot determine the cause of one person's ASD. Avoid implying blame from parental characteristics, social position, or the fact that a diagnosis became visible after a change in school or service access. The aetiology of ASD, including genetic and environmental architecture, belongs in the etiological discussion of this chapter; epidemiology here explains how cases are distributed and counted.

For a written answer, a compact but high-quality structure is: define prevalence and ascertainment; quote a source-qualified contemporary estimate; state the male predominance while distinguishing diagnosed from epidemiological ratios; then explain the rise through diagnostic concepts, awareness, practice, educational policy, and service access. End with the unresolved point: changing measured prevalence alone cannot settle whether underlying incidence has changed.

The rise in recorded ASD prevalence is real as a measurement trend; its causes are multiple, and it must not be equated automatically with a proven rise in true incidence.

20.8 High-yield synthesis

The epidemiology of ASD is best understood as a conversation between population occurrence and the systems that recognise it. Contemporary estimates demonstrate substantial need, while their variation demonstrates why source, denominator, geography, and ascertainment must be stated. Sex ratios have the same requirement: a well-ascertained epidemiological ratio and a diagnosed-rate ratio answer different questions. Historical increases should prompt examination of diagnostic change, awareness, education, services, and record-based case definition before claims are made about causation.

This approach protects against opposite errors. Minimisation assumes that ASD is uncommon because an older estimate was lower. Overstatement treats every rise in a recorded rate as proof of a new environmental cause. The clinician's role is to recognise the public-health burden, assess

the individual in front of them, and communicate uncertainty precisely. That combination is more useful than a detached list of figures and more defensible than a single dramatic number.

21 · Etiology of ASD (genetic, neurobiological, environmental)

Aetiology is plural, not a single cause. Autism spectrum disorder arises through developmental pathways in which inherited liability, rare genetic events, brain development and environmental exposures may intersect. The clinical task is therefore not to offer families a simplistic cause, but to explain that association is not destiny, that aetiological formulation can identify relevant medical conditions, and that a cause is not required before support can begin. The examination task is similar: distinguish evidence for familial aggregation from proof of an individual mechanism, and distinguish a risk factor from a cause.

The evidence is strongest for a substantial genetic contribution, but ASD is genetically heterogeneous. Its neurobiology is likewise a set of converging developmental observations rather than a diagnostic test. Environmental findings need particularly careful reading because many exposures are correlated with parental characteristics, obstetric events and ascertainment. A good answer therefore moves from genetic architecture to neurodevelopmental correlates, then appraises environmental association and vaccine misinformation without conflating any of these levels of explanation.




Fig 21.16 The aetiological architecture of autism. Six tiers, each printed with the strength its own source claims for it and against its own denominator. The plate teaches one shape: no row beneath twin-based heritability is both common and large, and the one row the ledger ties to a clinical action is syndromic autism. The common-variation lane is drawn empty because no such figure was sourced, and a hole in the map is not a hole in the world. The two traps and the refuted vaccine hypothesis are marked on the plate itself.

21.1 What an aetiological formulation means in ASD

Aetiological heterogeneity means that similar clinical phenotypes can arise through different causal routes. In practice, this prevents the error of treating all ASD as one disease with one lesion. Some presentations occur alongside a recognisable medical or genetic disorder; others reflect the combined effect of many inherited and developmental influences.

GOOD TO REMEMBER

Use the formulation to connect a person’s developmental history, physical examination and family pattern to an appropriate medical work-up; choice of ASD genetic tests belongs to the ASD medical workup section.

RULE Read causation at the right level: a population estimate of genetic influence does not tell us what caused ASD in one child.

Heritability is a population statistic: it describes the share of variation in a population that is associated with genetic differences under the conditions studied. In some studies, twin-based estimates for autism lie **between 80-90** percent of phenotypic variance. This is compelling evidence that genetic factors matter, but it does not mean that this proportion of any individual’s ASD was “caused by genes,” nor does it make environmental influences irrelevant. Heritability can change when a population or its exposures change.

Historical twin work made the importance of genetic factors visible after **Folstein and Rutter, 1977**. Twin and family designs are complementary. Twin concordance asks whether co-twins share the phenotype; family studies ask whether traits cluster among relatives. Neither design, on its own, identifies a gene or predicts the outcome for an individual pregnancy. A careful answer also states whether it is discussing narrowly defined autism or the broader phenotype.

approximately 60% MZ CONCORDANCE: NARROWLY DEFINED AUTISM	80-90% MZ CONCORDANCE: BROADER AUTISM PHENOTYPE	approximately 19% PROSPECTIVELY FOLLOWED INFANT SIBLINGS	5-15% SINGLE-GENE DISORDER OR CHROMOSOMAL ABNORMALITY
---	---	--	---

21.2 Familial aggregation and the broader phenotype

For strictly defined autism, monozygotic twin concordance is approximately 60%. When the outcome includes the **broader autism phenotype**, concordance is between 80-90%. The broader phenotype refers to subthreshold autistic-like traits in relatives, not to an assertion that every relative has ASD. This contrast is clinically instructive: genetic liability may be expressed as traits rather than as the full diagnostic phenotype, and the phenotype being measured changes the estimate.

Older retrospective family studies reported sibling recurrence between 2-3%. In prospectively followed infant siblings, the estimate is approximately 19%. These figures should not be presented as a contradiction. The later design observes siblings from birth and works with broader contemporary diagnostic ascertainment; both features can increase detection. For

counselling, it is more honest to explain that familial recurrence is elevated and that an individual family needs a personalised assessment rather than a memorised population figure.

Subthreshold autistic-like traits occur in above 20% of parents and siblings of ASD probands, compared with between 5-10% in control families. Such aggregation supports a dimensional account of inherited liability. It must not be turned into blame: a parent’s communication style, temperament or isolated trait is not evidence that parenting caused ASD.

■ **TRAP** **Examination trap:** do not quote the broader-phenotype twin figure as the concordance for narrowly defined autism, and do not use an early retrospective sibling estimate as though it came from a prospective infant-sibling design.

21.3 Genetic architecture: common liability, rare variants and syndromic ASD

Genetic architecture describes the kinds of genetic contribution found across a condition, rather than seeking one “autism gene.” ASD includes polygenic liability, rare inherited or de novo variation, copy-number variation and recognised syndromic disorders. These categories can overlap in a person, and the presence of a genetic finding should be interpreted alongside the phenotype. A result may clarify associated medical surveillance and family discussion, but it does not reduce the child to a laboratory label.

ARCHITECTURE	WHAT IT CONVEYS CLINICALLY	IMPORTANT DISTINCTION
Familial liability	Traits and diagnoses may aggregate in relatives.	Aggregation is not a single-gene explanation.
Rare de novo copy-number variation	A rare genomic event may contribute substantially in an affected individual.	Each individual rare event is below 1% of the ASD population, yet such variants cumulatively account for between 15-20%.
Chromosomal abnormality	May place ASD in a broader medical and developmental formulation.	Reported abnormalities involve many loci; 7q and 15q are described as common sites.
Syndromic association	Prompts attention to syndrome-specific physical and neurological features.	Association does not imply that every person with ASD has a syndrome.

Recognisable syndromic associations include fragile X, tuberous sclerosis, and neurofibromatosis. ICD-11 additionally lists cerebral palsy and early-onset epileptic encephalopathies among medical disorders associated with ASD, alongside chromosomal abnormalities and neurocutaneous disorders. These are associations that enrich clinical assessment; syndrome-specific genetics, phenotype and recurrence belong to their respective disorder sections.

Copy-number variation refers to a genomic deletion or duplication. The useful clinical contrast is frequency against collective contribution: any one rare de novo CNV occurs in **below 1%** of the ASD population, but the group taken together may account for between 15-20%. This is why a negative search for one famous variant never excludes a genetic contribution, and why genetic architecture cannot be understood by memorising a short gene list.

- Candidate genes named in the Indian guideline include neuroligins, SHANKs, CNTNAP2 and FMR1.
- FMR1 is the fragile X gene, and it is the single commonest monogenic contributor a candidate is asked to name here; its locus, repeat mechanism and phenotype are taught with the syndrome itself.
- CHD8 mutations illustrate a genotype-first association: above 80% of carriers meet diagnostic criteria for autism spectrum conditions.

Direction of a proportion is a frequent exam trap. Between 20-30% of boys with fragile X syndrome meet criteria for ASD, whereas approximately 1% of boys with ASD carry the fragile X mutation. These are different denominators and neither can be inverted. This distinction is also why a syndrome association supports targeted clinical consideration, not indiscriminate causal certainty.

21.4 Neurobiological findings: correlates, pathways and limits

Neurobiology informs models; it does not diagnose ASD. Studies of neurotransmitters, brain growth, imaging and postmortem tissue have identified group-level differences. They are heterogeneous, often developmentally timed, and may be affected by sampling, clinical variation and methodology. The appropriate conclusion is that ASD involves altered neurodevelopmental processes, not that any one scan, blood measure or neuropathological feature establishes the diagnosis.

Peripheral hyperserotonaemia is the most consistently replicated biochemical observation: elevated serotonin in blood and platelets is reported in between 25-50% of children and adolescents with autism. It is peripheral, not a direct measure of cerebrospinal fluid or brain serotonin. A candidate should therefore resist the tempting but invalid inference that it specifies a universal central neurotransmitter abnormality or tells the clinician how to treat an individual child.

Structural findings have focused on development and timing. In the first MRI study, total brain tissue volume in ASD was approximately 6% larger than in controls. **Megalocephaly** may be observable as young as 2 years of age, but was not present at birth in the cited observation. The teaching point is not that all autistic children have enlarged heads. It is that a developmental trajectory may differ after birth, and a cross-sectional image cannot be read as a universal sequence.

Postmortem studies describe increased neural packing, reduced cell size in the limbic system, reduced cerebellar Purkinje-cell number and altered cortical minicolumn organisation, as well as macroencephaly and altered brain-growth trajectories. Such work is constrained by small samples and tissue quality. It must be kept separate from in-vivo imaging. In a voxel-based meta-analysis, grey matter reduction was reported in the bilateral amygdala-hippocampus complex and bilateral precuneus, with a small increase in the middle-inferior frontal gyrus. These are group findings, not a radiological signature for clinical diagnosis.

21.5 Synaptic pruning as a mechanistic example

Synaptic pruning is a useful model for linking cellular findings with developmental circuitry. In the cited postmortem study, typically developing controls lost approximately 40% of dendritic spines across the age span, while individuals with ASD lost half as much as controls. The observation supports a hypothesis of less efficient pruning in at least some ASD tissue. It does not establish that deficient pruning is the sole pathway to ASD, nor that it is measurable in routine practice.

In the linked mechanistic work, **mTOR hyperactivity** was associated with impaired autophagy and disrupted synaptic pruning. In mice, rapamycin, an mTOR inhibitor, corrected pruning deficits and ASD-like behaviours. This is an important translational distinction: a correction in an animal model is not a recommendation for standard ASD treatment. The pathway is being examined in the tuberous-sclerosis subgroup, but the same source cautions that rapamycin is unlikely to become standard treatment for ASD. Mechanism-guided research is valuable precisely because it identifies testable subgroup hypotheses, not because it offers a universal cure.

MNEMONIC

Formulate, do not flatten

family pattern, genomic architecture, developmental brain evidence and exposure history belong in one formulation, but none should be promoted alone into “the cause.”

21.6 Environmental and perinatal associations: how to appraise them

Risk factor means an exposure or characteristic associated with a higher probability of outcome; it is not automatically a necessary or sufficient cause. Obstetric complications have some consistency across reviews, but associated odds ratios tend to be **below 3.0**. This magnitude and the possibility of confounding demand restraint. Complications may be on a causal pathway, may mark an earlier developmental difference, or may share determinants with ASD risk. The clinician should neither dismiss a careful obstetric history nor use it to assign parental blame.

Prenatal valproate exposure is a clinically important exception because it is recognised as a teratogenic exposure. ASD prevalence among those exposed prenatally is between 8-18 times that of the general population. ICD-11 also recognises prenatal valproate as an ASD risk factor. This is a risk statement about prenatal exposure, not permission to alter an individual's medication without specialist review. In clinical practice, risk communication must balance maternal illness, treatment alternatives and reproductive planning.

Parental age and migration findings illustrate why context matters. Maternal birth abroad is associated with a risk increase of exactly 28% in the general estimate, while the association in immigrant mothers in Nordic countries is a risk increase of exactly 58%. These findings may reflect a complex mixture of migration-related exposures, access to care, social context and ascertainment; they must never be used to stigmatise migrant families.

- The greatest paternal-age risk is reported for fathers over the age of 50, compared with fathers below 30 years.

- For mothers over the age of 30 years, the estimated increased risk is exactly 27%.
- Evidence supports advanced parental age as a risk factor, with a possibly greater paternal contribution. Proposed pathways differ: maternal ageing is linked with chromosomal imbalance, while paternal ageing is linked with point mutation, imprinting, sperm-cell changes, de novo CNVs and epigenetic influences.

The professional response to these associations is prevention-oriented but non-punitive: obtain a sound exposure history, coordinate medication decisions before conception where possible, and avoid retrospective certainty. Most importantly, do not imply that ordinary parental conduct produced the child's ASD.

21.7 Vaccines, parenting and the duty to correct misinformation

This is a safety conversation, not an optional controversy. Claims that vaccines cause autism can lead to delayed or absent vaccination and can damage trust at the point when families need clear guidance. The evidence should be explained calmly, directly and without amplifying the myth through excessive repetition. The same principle applies to blaming parenting: it is inaccurate, harmful and clinically obstructive.

Natural experiment evidence is particularly useful when an exposure changes across a whole population. The Japanese cessation of MMR showed that MMR was not responsible for an epidemic of autism. Separately, Scandinavian cessation of the mercury preservative thimerosal gave the same negative finding. These are distinct questions and should not be blurred: MMR and thimerosal are not interchangeable exposures.

■ **TRAP Examination trap:** do not write “vaccines are controversial causes.” The supported conclusion is that vaccines do not cause autism; withholding routine vaccination can harm the child.

IN INDIA

The Indian guideline, citing the CDC, rejects vaccine causation of autism and cautions that missing routine vaccination can harm the child. It also cites evidence that increasing exposure to vaccine antibody-stimulating proteins and polysaccharides was not associated with autism risk. These statements permit an unambiguous clinical recommendation while still making room for parents' fear to be heard respectfully.

When communicating a diagnosis, frame ASD as a lifelong neurodevelopmental disability with prenatal origins. Explain clearly that neither vaccination nor poor parenting produces it, and that children need not look alike clinically. This framing validates uncertainty without inventing a culprit. It also redirects attention toward assessment of strengths, support needs, associated medical conditions and intervention planning.

21.8 Clinical synthesis and examination answer

A complete aetiology answer begins with multifactorial neurodevelopmental causation and then separates evidence streams. Describe strong familial and twin evidence without treating heritability as individual causation. Explain the range from polygenic liability to rare CNVs,

chromosomal abnormalities and syndromic ASD, while referring genetic-test selection to the ASD medical workup. Use neurobiological observations as developmental correlates, state their limitations, and appraise environmental findings by study design, effect magnitude and confounding. End by explicitly correcting vaccine and parenting myths, because they are clinically consequential errors rather than harmless opinions.

For families, the formulation should be usable. It should say what is known, what remains uncertain, and what can be acted upon now. A syndrome-associated presentation may change surveillance and counselling; a prenatal exposure may affect future reproductive planning; neither changes the need for developmental support. Avoid giving a long list of alleged causes, which often leaves parents with blame rather than understanding.

ASD aetiology is best understood as heterogeneous neurodevelopmental causation: genes and developmental biology predominate, environmental associations require cautious causal inference, and vaccines and parenting do not cause autism.

The clinical picture of autism and how it is recognised

22 · Clinical features and core domains of ASD

The clinical task is to recognise a pattern, not to count isolated oddities. Autism spectrum disorder becomes visible when differences in social communication are understood alongside restricted, repetitive or inflexible patterns of behaviour and experience. A child may be quiet, highly verbal, bright, anxious, disruptive, apparently aloof, or warmly attached to familiar people. None of those impressions settles the question. The useful clinical formulation asks how the person shares attention, uses communication with another mind in view, manages reciprocity, and responds when routine, expectation or sensory input changes.

Recognition matters because the presentation is developmental and contextual. The same behaviour can have different meanings at different developmental levels, and a behaviour that looks unusual in a brief consultation may be an adaptive response to uncertainty. History, direct observation and reports from the settings in which the person lives should therefore be brought together. The diagnostic criteria are addressed separately; this section concerns the clinical phenotype that makes those criteria intelligible at the bedside.

approximately 20% REPORTED REGRESSION OR DEVELOPMENTAL STAGNATION	12-24 months USUAL RECOGNITION WINDOW	12-24 months USUAL WINDOW FOR SKILL LOSS, WHEN PRESENT	10-50% ESTIMATED RANGE FOR SAVANT OR SPLINTER SKILLS
---	---	--	--

22.1 Recognising a developmental social-communication difference

Autism is usually recognised through a qualitative difference in how social information is noticed, interpreted and used. The central issue is not whether the person wants company in an abstract sense. It is whether interaction is organised as a reciprocal exchange in which another person's

attention, knowledge, feeling and response alter one's own communication. A person may seek proximity, enjoy affection, or initiate contact for a need, yet still have difficulty sustaining the back-and-forth adjustments that make interaction feel shared.

Social reciprocity includes initiating interaction, responding contingently, sharing interests and affect, and allowing the partner's response to shape the next move. In young children this may be seen as limited showing, bringing, pointing out or checking back after an event. In older children and adults it may appear as a one-sided conversation, a tendency to return rapidly to a preferred topic, or difficulty recognising when the listener is confused, bored or trying to change direction. These are descriptions of the quality of exchange, not moral judgements about empathy, attachment or motivation.

A careful history separates capacity from performance. Some people communicate much less when overloaded, unfamiliar with the examiner, tired, frightened or expected to respond rapidly. Others have learned scripts that make initial interaction appear fluent but that do not readily flex when the conversation departs from the script. The clinician should look for developmental continuity and for the effort required to maintain apparently typical social behaviour. A socially successful encounter does not erase a pervasive developmental difficulty, just as an awkward encounter does not establish one.

-
- **RULE** Interpret the social behaviour in its communicative context: the question is how the person shares a mind with another person, not whether eye contact or sociability is simply present or absent.
-

GOOD TO REMEMBER

At the bedside, ask the caregiver for ordinary examples of sharing an achievement, reacting to another person's distress, joining group play, managing a misunderstanding or telling a story to a listener who does not already know it. Then observe the response to an unexpected but benign shift in topic or activity: does the person repair communication, seek clarification, withdraw, perseverate or become distressed? This helps distinguish a stable developmental pattern from a situational reaction.

CLINICAL DOMAIN	WHAT TO EXPLORE	HOW IT MAY PRESENT IN CONSULTATION	INTERPRETIVE CAUTION
Reciprocal exchange	Initiation, response, turn-taking and sharing	Conversation may be asymmetrical, unusually passive, or driven by an immediate need	Shyness and anxiety can reduce performance without creating the developmental pattern
Non-verbal integration	Gesture, facial expression, gaze and timing	Signals may be limited, atypical, poorly coordinated with speech, or difficult to read	Do not convert one gaze style into a diagnosis
Relationships	Peer interest, social imagination and adaptation to roles	Rules of friendship may be understood literally; group interaction may be hard to enter or sustain	Solitude may be a preference, a coping strategy, or both
Repetition and inflexibility	Patterns, routines, focused interests and sensory responses	Distress, persistence or repetitive action becomes prominent when expectations shift	Determine function and impairment before labelling a preference pathological

22.2 Non-verbal communication and shared attention

Non-verbal communication is not a decorative adjunct to speech. Gesture, posture, facial expression, gaze, distance, timing and prosody carry much of the meaning of social exchange. In autism, the difficulty may lie in producing these signals, coordinating them with words, noticing those used by others, or inferring the unstated message they convey. The pattern is variable. Some individuals use abundant gesture but in an idiosyncratic or poorly integrated way; others have a relatively neutral facial style yet understand familiar relationships well.

Joint attention is especially useful clinically because it captures the difference between requesting and sharing. A person can point to obtain an object while finding it harder to point simply to make another person notice something interesting. Similarly, a child may follow a hand because it predicts access to a desired item but not because the other person's focus is meaningful in itself. The history should ask about following another's attention and actively directing it. Direct observation can then assess whether the person shifts attention between an object and a partner, checks the partner's reaction, or incorporates the partner into the experience.

Do not reduce this to an eye-contact test. Gaze is influenced by culture, anxiety, visual ability, temperament and the demands of the interaction. Sustained eye contact may be uncomfortable, distracting or consciously performed. Conversely, apparently good eye contact may coexist with marked difficulty using its social meaning. A better formulation is that visual social signals can be used differently, sometimes with little coordination between gaze, gesture and spoken language.

- Explore whether gestures are used to share as well as to request.
- Observe whether facial expression and voice match the intended message.
- Ask how misunderstandings are repaired when the listener does not follow.

- Consider whether social signals are missed, over-read or interpreted literally.

■ **TRAP** Examination trap: neither reduced eye contact nor a preference for solitude is a sufficient clinical shortcut. The developmental pattern across reciprocal communication, non-verbal integration and flexibility is what carries meaning.

22.3 Language phenomenology: form, use and reciprocity

Language in autism must be described at the level of use, not merely the presence or absence of words. The language phenomenology of autism includes differences in developmental progress, conversational reciprocity, prosody, literal interpretation and the ability to adjust communication for another person's perspective. Language ability is heterogeneous. Some people have very limited spoken language; others have extensive vocabulary and grammatically complex speech while finding the social use of language persistently difficult. Language impairment is considered separately as a specifier and should not be confused with the social-communication phenotype itself.

Pragmatic language concerns what speech does between people. It includes selecting relevant information, recognising implied meaning, taking turns, maintaining an appropriate topic, judging what the listener knows, and repairing breakdowns. A child may answer factual questions accurately but not volunteer information that would make the account coherent for someone else. An adult may speak fluently at length about a preferred interest yet have trouble sensing the reciprocal rhythm of conversation. These patterns are often most evident in unstructured conversation rather than in direct question-and-answer exchanges.

Echolalia refers to repetition of heard language. It may be immediate or delayed, but the clinician should not assume that it is meaningless. Repeated phrases can request, protest, regulate emotion, preserve a familiar sequence, or communicate an association that is not obvious to the listener. The clinical task is to determine the communicative function and the degree to which repetition substitutes for flexible, generative language. Similarly, pronouns may be used in an unusual perspective, and speech may have an atypical rhythm, volume or melody. Such features are useful only when understood as part of the wider developmental pattern.

History taking should locate the change in ordinary life. Ask how the person tells a story, joins a conversation already in progress, copes with jokes and indirect requests, or reacts when someone asks an ambiguous question. A family may report that the person is articulate but seems not to "read the room". That everyday observation can be more clinically revealing than a formal vocabulary performance. Equally, a bilingual environment should not be mistaken for a cause of autism or used to excuse a pervasive social-communication concern. The relevant question is whether the pattern is present across the languages and relationships available to the person.

For the intervention plan, speech and language therapy belongs with the broader support approach described in the relevant section of this chapter. At assessment, the immediate responsibility is more basic: document what communication achieves, what breaks down, and what support makes the person's intention understandable.

22.4 Restricted patterns, insistence on sameness and sensory experience

Restricted and repetitive behaviour is clinically broader than obvious motor stereotypy. It includes repetitive movements or speech, a marked need for predictability, unusually intense or narrowly focused interests, and atypical responses to sensory input. Its common thread is a reduced ease with variation or a distinctive way of regulating attention, arousal and meaning through repetition. The behaviour may be quietly internalised and therefore missed in an articulate adolescent or adult whose outward presentation appears composed.

Insistence on sameness is often best recognised through the consequences of disruption. The person may rely on a fixed route, sequence, object arrangement, conversational script or rule. A change may produce distress, anger, shutdown or prolonged preoccupation, not because the preference is trivial but because predictability is serving an organising function. Obtain concrete examples from home, school, workplace and travel. Then ask whether the routine can be modified with preparation, whether the response is proportionate, and whether the restriction limits participation or causes conflict.

Sensory reactivity may involve unusually strong, weak or seeking responses to sound, touch, light, smell, taste, movement or visual detail. It should be elicited as lived experience rather than treated as a box to tick. A child who covers their ears may be in pain, anticipating noise, avoiding a social demand, or doing more than one of these. An adult who avoids crowded places may be managing sensory overload, social uncertainty, panic, or a combination. The formulation must link trigger, response, function and consequence.

- Ask what the person repeatedly does, says, collects, watches or arranges.
- Ask what must happen in a particular order and what follows if it does not.
- Explore focused interests for intensity, flexibility and impact, rather than judging the topic itself.
- Map sensory triggers, protective strategies and the effect on ordinary participation.

Focused interests can be a route to competence, pleasure, identity and connection. They become clinically relevant when their intensity or inflexibility constrains functioning, blocks reciprocal engagement, or makes transition difficult. This distinction protects against pathologising expertise or temperament. It also enables the clinician to use an interest respectfully as a bridge during assessment rather than treating it only as a symptom.

MNEMONIC

Clinical spine - SHARE

Explore **S**ocial reciprocity, **H**ow communication is used, **A**daptation to change, **R**epetitive or focused patterns, and **E**xperience of sensory input. It is an enquiry sequence, not a diagnostic shortcut.

22.5 Developmental regression and developmental stagnation

Developmental regression means loss of abilities that were previously acquired and used, whereas developmental stagnation refers to a plateau in expected gains. Both deserve a careful timeline. A report may be imprecise because a skill was inconsistent, context-bound or only recently emerging. The clinician should therefore ask for examples of the earlier ability, its ordinary use, the sequence of change, and what else changed around the same period. In autism, regression or stagnation is reported in approximately **20%** of cases.

Autism symptoms are typically recognised between **12 and 24 months**. Where previously acquired social or language skills are lost, the loss itself also typically occurs between **12 and 24 months**. These statements describe distinct clinical questions: recognition concerns when the pattern comes to notice, while regression concerns the timing of actual skill loss. Keeping that distinction prevents a history of late recognition from being incorrectly recorded as regression.

Loss of previously acquired skills is usually described during the second year of life and is rarely observed after **3 years of age**. When it occurs, the loss most often concerns language use and social responsiveness. If the reported course is later or broader, document it precisely rather than forcing it into an autism narrative. If a loss occurs after 3 years of age, it is more likely to include changes in cognitive and adaptive functioning, continence, sleep, language, social ability, emotion and behaviour. Such a presentation needs separate evaluation; the differential diagnosis of regression is addressed elsewhere in this chapter.

Regression is emotionally potent for families. They often remember a before-and-after story, and it is important neither to dismiss that account nor to turn it prematurely into an aetiological explanation. Clarify whether the child lost a functional word, stopped responding to name, ceased sharing interest, became less socially responsive, or simply failed to add new skills. Review contextual changes and establish whether the apparent loss was global or selective. The record should preserve the caregiver's observations in concrete behavioural language, because that is more useful than the label alone.

A plateau and a loss are not interchangeable: establish the prior skill, its functional use and the sequence of change before calling a course regressive.

22.6 Cognitive style, uneven ability and exceptional islands of skill

There is no single cognitive profile that defines autism. What is clinically common is an **uneven profile of abilities**, including in people whose general intellectual ability is average or high. A person may have striking factual knowledge, rapid pattern recognition or strong rote learning alongside difficulty applying ability flexibly in everyday social and practical contexts. The gap between demonstrated intellectual skill and adaptive functioning can be conspicuous. This is why a superficially impressive verbal or factual performance should not end the assessment of support needs.

Several cognitive accounts have helped clinicians describe aspects of the phenotype: **theory of mind**, weak central coherence, and executive dysfunction. Difficulty representing another

person's perspective can illuminate social misunderstandings. A detail-focused processing style can illuminate difficulty deriving the broader context. Problems with planning, shifting set, organising and inhibiting a habitual response can illuminate inflexibility. None should be used as a universal explanatory key. The cognitive theories do not establish a unitary cognitive origin for all manifestations of autism.

Savant or splinter skills describe competencies that stand out above the individual's general level of ability or above population expectation. Estimates vary with how such skills are defined, ranging between **10% and 50%** of people with autism. Their presence does not negate disability in another domain, and their absence has no diagnostic implication. Clinically, ask what the skill enables, whether it is enjoyable or burdensome, and whether it can be incorporated into education, work, self-advocacy or engagement.

Mental-state, contextual-integration, executive and motor features often discussed in autism are descriptive aids, not a substitute for developmental assessment. They can explain why an individual may know a rule yet fail to use it in a fast-moving social situation, or complete a complex task yet struggle with the unstructured demands around it. The practical lesson is to look for discrepancies: between knowledge and use, verbal expression and reciprocal conversation, performance in a preferred task and performance under change, or stated independence and actual day-to-day functioning.

22.7 A disciplined clinical formulation

A high-quality formulation is developmental, functional and contextual. It anchors current observations to the early pattern, describes what the person can do independently and what becomes difficult, and identifies environmental adjustments that alter performance. Home, educational, occupational and peer settings may reveal different aspects of the same difference. This framing matters when a person camouflages difficulties or when family routines have hidden impairment in consultation.

Use examples rather than labels wherever possible. "Talks at length about one topic and does not notice attempts to change it" is more clinically useful than "poor reciprocity" alone. "Becomes unable to continue when the usual route changes" is more useful than "rigid". Concrete descriptions allow later review, enable communication across disciplines and protect against diagnostic overshadowing. They also help the person and family identify accommodations without requiring the clinician to infer motives from a behaviour.

Changes from the person's usual baseline need particular care. Do not attribute new distress, marked withdrawal, loss of function, self-injury, agitation, sleep change or altered eating automatically to autism. Medical and psychiatric comorbidity, including catatonia, is considered in the medical comorbidities and workup section of this chapter. The clinician's immediate job is to recognise a baseline change and investigate it as a change, rather than accepting it as an inevitable expression of the developmental condition.

The assessment must also leave room for the person's strengths, preferences and account of their own experience. A formulation that records only deficits will miss the conditions in which

communication works, the interests through which rapport can be built, and the strategies that already reduce overload. The aim is not to make every distinctive preference disappear. It is to understand the pattern well enough to distinguish difference from impairment and to identify where support can make participation more possible.

23 · Severity levels and specifiers (with or without ID, language impairment)

Severity is a statement about current help, not a shorthand for ability. In autism spectrum disorder (ASD), a useful formulation has to keep several descriptions in view at once: the intensity of core autistic difficulties, intellectual functioning, language functioning, associated conditions and the person’s changing circumstances. Collapsing these into one label makes the record look tidy but makes clinical planning less accurate. The diagnostic formulation should therefore preserve distinctions that matter to the person, family and team.

The immediate examination value is equally practical. A severity level describes the support needed for the core autistic presentation. It does not quantify intelligence, does not replace a language description, and does not exhaust the associated clinical picture. Specifiers make those separations visible. They turn a broad diagnostic name into a clinically communicable account of the individual sitting in front of the clinician.

3 DSM-5-TR SUPPORT LEVELS	2 DOMAINS RATED SEPARATELY	3 ICD-11 FUNCTIONAL-LANGUAGE LEVELS
------------------------------	-------------------------------	--

23.1 What an ASD severity level is intended to convey

Start with the question behind the label. The DSM-5-TR severity framework is a description of **support need in the core autistic domains**. It asks how much assistance is currently required because of social-communication difficulty and because of restricted, repetitive patterns of behaviour. It is thus a functional clinical description, not a global ranking of the person.

There are **three** named support levels: **Level 1, requiring support**; **Level 2, requiring substantial support**; and **Level 3, requiring very substantial support**. The wording is deliberately about the assistance needed. A level should prompt the clinician to ask what supports permit participation, communication, flexibility and safety, rather than being treated as a fixed essence of the person.

■ **RULE** **Rule:** Record ASD severity as support need in the core domains, and record intellectual and language impairment separately.

That distinction matters because observed difficulty can be shaped by context. A familiar, predictable setting with effective communication supports may reveal capacities that are obscured in an unfamiliar assessment room. Conversely, a change in demands, sensory load, illness, distress or available support may expose need that was not apparent previously. The severity description must therefore be anchored in current functioning and meaningful participation, not in a superficial impression formed from a brief encounter.

The phrase “severity” can mislead candidates because the same word is used elsewhere in developmental psychiatry. Intellectual disability has its own severity classification, which is dealt with in the severity classification of intellectual disability section. The ASD framework answers a different question. It does not establish the presence of intellectual impairment and it does not substitute for an assessment of adaptive functioning. Keeping the questions separate prevents a common category error.

23.2 Rate the domains separately, not the person globally

DSM-5-TR asks that the level of support be recorded separately for **two** domains. One rating concerns social communication; the other concerns restricted, repetitive behaviour. This is not clerical fussiness. These domains can create quite different practical barriers, and their support requirements may not move together.

A person may need intensive help to manage inflexibility, transitions and repetitive behaviour while communicating more effectively in a structured interaction. Another may have relatively less conspicuous repetitive behaviour in a short consultation yet need extensive support to sustain reciprocal communication across daily settings. A single global level would conceal that pattern and can lead the receiving team to organise support around the wrong problem.

The social-communication rating asks how much support is needed for reciprocal communication. The restricted and repetitive behaviour rating asks how much support is needed for rigidity, repetitive behaviour and related interference. The intellectual-impairment specifier asks whether intellectual impairment accompanies ASD, while the language-impairment specifier asks whether language impairment accompanies it and, if so, what current verbal functioning is like. None of these questions can be answered by simply borrowing the answer to another.

GOOD TO REMEMBER

In a clinical note, record the two domain ratings with the observed barriers and the supports that alter them. A later change in level may reflect a change in support need, setting or the effect of effective intervention; it is not a prognosis.

■ **TRAP** **Classic trap:** Do not derive the ASD level from an IQ score. Intellectual impairment is carried by a separate specifier, not by the ASD severity level.

Equally, do not infer the support level merely from speech quantity. Language impairment and social-pragmatic difficulty overlap in everyday presentation but are not interchangeable descriptions. The language phenomenology of ASD belongs with the clinical features of ASD section; here the task is to show how language is captured in the diagnostic formulation without converting it into a core-severity score.

23.3 DSM-5-TR specifiers: add the clinically relevant dimensions

Specifiers prevent the broad ASD diagnosis from becoming clinically empty. They should be stated positively where present and negatively where absent, after assessment rather than by assumption. The first pair addresses accompanying intellectual and language impairment: **with**

or without accompanying intellectual impairment, and **with or without accompanying language impairment**.

The intellectual specifier is not an invitation to re-grade intellectual disability within the ASD note. It flags a co-occurring dimension that changes communication, assessment interpretation, educational planning and the kind of support likely to be required. The underlying assessment and severity classification of intellectual disability should be documented using their own standards. In particular, an autistic person's social difficulty should not be mechanically treated as evidence of global adaptive limitation.

The language specifier is similarly an addition, not a restatement of ASD itself. When language impairment is present, the clinician describes the current level of verbal functioning. This avoids the unhelpful shorthand of treating all language difficulty as equivalent. It also guards against the opposite error: assuming that fluent speech rules out significant social-communication disability. The diagnostic criteria section explains why language impairment is not itself a DSM-5 core criterion; the present specifier tells the reader whether it accompanies the syndrome.

Other specifiers place ASD in its wider clinical setting. The formulation may state that it is associated with a known genetic or other medical condition or environmental factor. This wording records an association or context; it must not be casually converted into a causal conclusion. It may also state that ASD is associated with a neurodevelopmental, mental, or behavioural problem. The full map of co-occurring conditions and how to detect them belongs in the medical comorbidities and workup in ASD section.

- Use the intellectual-impairment specifier to make accompanying intellectual impairment explicit, not to estimate it from presentation.
- Use the language-impairment specifier to record an accompanying language problem and current verbal functioning.
- Use the known-condition wording to state context accurately without claiming causation that the evidence has not established.
- Use the associated-problem wording to ensure that important co-occurring neurodevelopmental, mental or behavioural difficulties do not disappear behind the ASD label.

For the candidate, this is a disciplined form of diagnostic writing. “ASD” alone describes the syndrome but not the clinically relevant modifiers. A complete formulation makes visible which parts of the presentation are core, which are accompanying impairments, and which require parallel assessment or treatment. That distinction also protects against diagnostic overshadowing, where all distress or disability is wrongly attributed to autism.

23.4 Catatonia: a specifier that requires active clinical attention

Do not let a specifier become a footnote. DSM-5-TR permits the specification **with catatonia**. This should prompt careful assessment of a change from the individual's usual state, including

motor behaviour, responsiveness, activity and functional capacity. A diagnostic label should never replace a longitudinal account of what has altered and what the family or carers have observed.

When catatonia accompanies ASD, DSM-5-TR directs use of the additional code **F06.1**. It is a separate diagnostic code, not an ASD severity modifier and not a reason to raise a support level by default. The distinction is clinically important: support-level language describes core autistic support needs, whereas catatonia identifies an associated syndrome that requires its own recognition and management pathway.

The period of greatest risk described in DSM-5-TR is the adolescent years. That fact should sharpen clinical vigilance without turning ordinary developmental change into a diagnosis. When there is concern, document baseline, timing and trajectory, seek collateral from people who know the individual well, and evaluate the wider clinical context. Recognition belongs with the ASD comorbidity assessment; treatment is addressed in the pharmacotherapy in ASD section.

A support level is not a catch-all label: ASD severity, accompanying intellectual impairment, accompanying language impairment and catatonia each answer different clinical questions.

23.5 ICD-11: characterisation through intellectual development, language and loss of skills

ICD-11 takes a differently structured approach. It does not supply DSM-style support-need severity levels. Instead, ASD is characterised through specifiers relating to **intellectual development, functional language and loss of previously acquired skills**. This is not a minor terminological difference. A clinician transferring information between systems must not invent a support level from an ICD-11 code, nor assume that an ICD-11 language category is a DSM support rating.

The intellectual-development axis is expressed as **with or without disorder of intellectual development**. In ICD-11, a co-occurring disorder of intellectual development is diagnosed separately rather than folded into the ASD diagnosis as an inline descriptor. When assessing this co-occurrence in ASD, greater weight should be given to the intellectual, conceptual and practical aspects of adaptive functioning than to social skills. This protects against mistaking social difficulties intrinsic to autism for evidence of a broader adaptive disorder.

The **functional-language axis** has **three** levels. It distinguishes mild or no functional-language impairment, impaired functional language, and complete or almost complete absence of functional language. The middle category concerns a person unable to use more than very short spoken expressions. These categories primarily describe functional expressive language, including nonverbal expression where relevant; they do not replace assessment of the pragmatic communication difficulty that is central to ASD.

FRAMEWORK	WHAT IT RECORDS	CLINICAL READING
DSM-5-TR ASD severity framework	Current support need in each core domain	Use to communicate the assistance required for core autistic difficulties
DSM-5-TR intellectual specifier	Presence or absence of accompanying intellectual impairment	Keep intellectual impairment distinct from ASD support level
DSM-5-TR language specifier	Presence or absence of accompanying language impairment	Describe current verbal functioning when impairment is present
ICD-11 ASD characterisation	Intellectual-development, functional-language and loss-of-skills specifiers	Do not translate this structure into a support-level rating

ICD-11 also records whether there has been loss of previously acquired skills. The suffixes 6A02.x0 and 6A02.x1 distinguish absence from presence of such loss. This coding distinction should not be used to teach the phenomenology or differential diagnosis of regression, which are considered elsewhere in the chapter. Its value here is to show that ICD-11 preserves a clinically salient developmental course feature within the diagnostic characterisation.

23.6 ICD-11 coding: use the matrix, do not guess it

The ICD-11 code reflects the combination of intellectual-development and functional-language descriptors. It is safer to determine the descriptors first, then select the matching code, rather than attempting to recall a code from an impression of severity. The following are selected, explicitly stated combinations. They are examples for accurate use, not a substitute for the full coding matrix.

DOCUMENTED COMBINATION	ICD-11 CODE	POINT TO RETAIN
Without disorder of intellectual development, with mild or no functional-language impairment	6A02.0	This is a coding combination, not a DSM support level.
Without disorder of intellectual development, with impaired functional language	6A02.2	Language description and intellectual-development status are both needed.
With disorder of intellectual development, with mild or no functional-language impairment	6A02.1	Co-occurring intellectual-development disorder is represented in the code.
With disorder of intellectual development, with impaired functional language	6A02.3	Do not infer support need from this code.
With disorder of intellectual development, with complete or almost complete absence of functional language	6A02.5	Use the language descriptor that is actually documented.

ICD-11 provides 6A02.Y for other specified ASD when the parameters do not apply, and 6A02.Z for unspecified ASD when the parameters are unknown. These are not interchangeable expressions of uncertainty. Good documentation reduces unnecessary use of the unspecified category while respecting genuine limits of available information.

23.7 A practical method for writing the formulation

Write the formulation in layers. First establish the diagnosis using the diagnostic criteria. Then describe the current ASD support need in each core domain when using DSM-5-TR. Next state the intellectual and language specifiers, followed by relevant associated-condition and associated-problem specifiers. Finally, record catatonia when present and use the appropriate additional code. This order mirrors clinical reasoning: syndrome, core impact, accompanying impairments, then associated context.

- State the diagnostic system being used so that the reader knows whether a support-level or coding-matrix framework applies.
- For DSM-5-TR, record the two core-domain support ratings separately and add a short functional explanation of each.
- State whether accompanying intellectual impairment and accompanying language impairment are present, without converting either into an ASD level.
- Record associated conditions and associated problems as clinical context, and keep association distinct from causal attribution.
- For ICD-11, establish intellectual-development status and functional-language status before selecting the code and any loss-of-skills suffix.

This layered method also improves communication across disciplines. It tells a speech and language clinician that a language issue is recognised without pretending that language alone explains the autistic presentation. It tells an educational or disability team that intellectual impairment requires its own assessment. It tells the medical team that an associated condition needs parallel attention. Above all, it avoids turning a diagnostic label into a description that is too vague to guide care.

When information is incomplete, state what has and has not been assessed rather than filling the gap with an assumed level. Diagnostic precision is not achieved by false certainty. A provisional formulation can still be useful if it clearly separates observed support needs from unassessed intellectual functioning, language status or associated conditions. The next assessment question then becomes obvious.

23.8 High-yield discriminations

The strongest answers distinguish neighbouring ideas before they list them. ASD severity is a support-need description for the core autistic domains. Intellectual impairment and language impairment are accompanying specifiers. ICD-11 uses a different characterisation system, based on intellectual development, functional language and loss of previously acquired skills rather

than DSM-style support levels. These are complementary ways of making heterogeneity visible, not rival measures of a single latent “severity”.

In practice, the formulation should remain revisable. Support needs can change with environment, development, communication access, co-occurring clinical problems and the fit between the person and available services. Revision is not inconsistency when it reflects a better account of current functioning. What should remain stable is the discipline of recording each dimension for what it means, rather than allowing a memorable label to do the work of assessment.

24 · Screening and diagnostic tools (M-CHAT, ADOS, ADI-R, CARS, ISAA, INCLIN)

Tools organise evidence; they do not substitute for clinical judgment. Autism assessment is often difficult not because there is a shortage of instruments, but because instruments answer different questions. A brief parent questionnaire can identify a child who needs closer attention; an observed interaction can make social communication and restricted behaviour more visible; a developmental history can establish whether the pattern has been present across time; and a disability assessment can document functional impact for statutory purposes. Treating all of these as interchangeable produces familiar errors: diagnosing from a screen, or mistaking an administrative score for the diagnostic formulation.

For the candidate and the clinician, the useful organising question is therefore not, “Which scale proves autism?” It is, “What decision is this instrument designed to support?” A positive screen changes the urgency of assessment. A clinician-administered diagnostic measure can sharpen an already careful formulation. An Indian disability instrument serves a further, legally important function after diagnosis. This sequence protects children from delayed support while preserving the clinician's responsibility to weigh developmental history, direct observation, informant accounts, context, and alternatives.




Fig 21.17 The ASD recognition pathway. Surveillance raises the concern, screening estimates the risk, the diagnosis is clinical, and both the workup and the Indian certification route follow it. Only the M-CHAT-R/F algorithm is sourced end to end; every CARS-2, ISAA, ADOS-2 and ADI-R cut-off is named on the plate and left undrawn.

24.1 A functional taxonomy of autism instruments

Screening is a process for finding children whose developmental profile warrants further assessment. Its purpose is sensitivity to concern, not a final statement about diagnosis. A screen can be negative when concerns remain clinically persuasive, and it can be positive because another developmental or behavioural difficulty needs assessment. Conversely, the absence of a score should never close a history that is concerning to parents or a clinician. **The M-CHAT-R/F** is a parent-rated, autism-specific toddler screen with a second-stage follow-up interview; it is not a diagnostic instrument.

■ **RULE** **Rule:** Match the instrument to its decision, then return the result to a clinical formulation rather than allowing the score to become the formulation.

CLINICAL QUESTION	USEFUL INSTRUMENT CLASS	WHAT THE RESULT SHOULD DO
Is there sufficient concern to escalate assessment?	Parent-rated autism-specific screen	Trigger follow-up questioning, surveillance, or referral according to the algorithm.
Can autism-related behaviour be sampled directly?	Observational assessment	Add direct clinical observations to the history and examination.
What was the developmental course?	Developmental-history interview	Clarify onset, change over time, and examples across settings.
Does a scale indicate concern or quantify observed features?	Clinician-rated screening scale	Support communication and follow-up, without becoming the diagnosis.
What is required for disability documentation in India?	Post-diagnostic disability assessment	Quantify disability and support access to certification-related entitlements.

The distinction is clinically productive. A questionnaire brings the caregiver's longitudinal experience into the consultation. An observation samples what happens under clinical conditions, but cannot by itself reproduce every home, school, or peer context. A history interview may expose a meaningful developmental trajectory, but its interpretation depends on the quality of the informant account and the clinician's questions. Each contributes a different kind of evidence. Diagnostic confidence comes from their convergence with informed judgment, not from collecting the largest possible pile of forms.

Informed clinical judgment remains central: diagnostic instruments supplement it and no single tool should provide the sole basis for diagnosis. This is not a vague appeal to intuition. It means the clinician must decide whether reported and observed features form a coherent autism pattern, whether they are developmentally meaningful, and whether another explanation better accounts for the presentation. The full assessment framework belongs with the Child psychiatry: assessment, treatment, services and protection notes; the present task is to understand what each autism-specific tool can and cannot add.

24.2 M-CHAT-R/F: using a screening algorithm safely

A screen is an invitation to think again. The M-CHAT-R/F is designed for toddlers and is parent rated. The revised checklist contains 20 items. Its follow-up interview matters because an initial answer may reflect misunderstanding of the question, an unusual circumstance, or a concern that becomes clearer when a clinician asks for examples. In practical terms, the first stage identifies responses needing clarification; the follow-up turns a crude count into a more clinically usable account of risk responses.

0-2 LOW FIRST-STAGE BAND	3-7 FOLLOW-UP REQUIRED	2 or higher POSITIVE AFTER FOLLOW-UP	8-20 FOLLOW-UP MAY BE BYPASSED
-----------------------------	---------------------------	---	-----------------------------------

At a first-stage total of 0-2, the result falls in the low-risk band. No further algorithmic action is required unless surveillance itself indicates risk. For a child younger than 24 months, with repeat screening after the second birthday, the guidance is to screen again. This is a useful reminder that a reassuring questionnaire is not permission to ignore a parent's developing concern, a developmental history, or a change in social engagement.

GOOD TO REMEMBER

A first-stage total of 3-7 is the medium-risk band and calls for the follow-up interview. Do not call this initial score the diagnostic threshold. It is a decision point for gathering better information about at-risk answers. After follow-up, a score of at least 2 is screen positive and should lead to referral for diagnostic evaluation and consideration of early-intervention eligibility. A follow-up total of 0-1 is screen negative; ongoing surveillance and rescreening at future well-child visits remain appropriate when concerns arise.

At a first-stage total of 8-20, out of a possible 20, it is acceptable to bypass the follow-up stage. The important interpretive point is that bypassing an interview is a rule within a screening algorithm, not a licence to pronounce a diagnosis from the questionnaire. In any referral, record the informant's examples, what was observed in clinic, and why the referral is being made. This makes the result clinically portable rather than a detached number.

Scoring direction is an easily missed technical trap. On most items, a response indicating risk is “no”; the exceptions are items 2, 5 and 12, where “yes” indicates risk. A clinician should not attempt to reconstruct the scoring from recollection while interpreting a parent form. Use the approved scoring algorithm, check that the correct revised version is being used, and document whether the score is before or after follow-up. That last distinction prevents a medium-risk first-stage score from being misreported as a positive screen.

■ **TRAP Examination trap:** A positive follow-up screen warrants diagnostic evaluation; it is not itself a diagnostic threshold.

Autism-specific screening is recommended at 18 and 24 months and whenever surveillance raises concern, alongside surveillance at developmental and psychiatric assessments. This differs from the schedule for general developmental screening, which is recommended at 9, 18, and 24 or 30

months during regular well-child visits. The distinction matters because a general screen and an autism-specific screen answer overlapping but not identical questions. The normal developmental context and milestones that make a particular response concerning are developed in the Normal child development and milestones notes.

24.3 Diagnostic instruments: observation, history, and trained use

Diagnostic measures are not simply longer screening scales. The ADI-R and ADOS-2 are described as state-of-the-art diagnostic measures and require specific training for clinician administration. Training is not ceremonial. It is needed to standardise how prompts are given, how behaviours are recognised and coded, and how the resulting information is used without overriding the clinical context. An untrained, improvised administration can create an appearance of precision while reducing the value of the observation.

ADOS-2 is an interviewer-administered observational measure of autism-spectrum symptoms. Its contribution is direct sampling: the assessor sees how communication, reciprocity, play, flexibility, and other relevant behaviours emerge in interaction. That is valuable when caregiver descriptions and clinic impressions diverge, when the child presents differently across contexts, or when the clinician needs well-described behavioural examples for a multidisciplinary discussion. It remains a sample, however. A calm consultation, a familiar caregiver, fatigue, anxiety, unfamiliar language, or the child's strategy for coping with novelty can all influence what is observed.

ADI-R is a semi-structured instrument for obtaining a developmental history. It is particularly useful conceptually because autism is not diagnosed from a single cross-sectional interaction. The interviewer asks the caregiver to move from labels such as “he does not mix” to concrete examples: what the child did, in what situation, whether the pattern was sustained, and how it changed. A good developmental history distinguishes a stable developmental style from a recent change and makes it possible to test whether apparently isolated symptoms belong together. This also protects against treating a questionnaire response as though it were already a clinical observation.

These instruments should be read as complementary rather than competitive. One asks what behaviour can be observed now; the other helps establish developmental meaning through a knowledgeable informant. Neither removes the need to consider communication ability, sensory or physical factors, environmental demands, and alternative developmental formulations. Nor should their absence be used to deny care where an experienced clinician can make a sound diagnosis. Their special value lies in disciplined, reproducible evidence when the assessment is complex, contested, or requires detailed communication across services.

MNEMONIC

H-O-L-D the hierarchy

History, Observation, Linkage across contexts, then Diagnostic judgment. It is a sequence for thinking, not a substitute scoring rule.

24.4 Rating scales and caregiver questionnaires: useful, but bounded

Rating scales create a compact record of behaviours and can support screening, communication, and longitudinal review. Their apparent simplicity is also their risk. A numerical total can look more authoritative than a detailed narrative, yet a rating depends on who observed the behaviour, what comparison they used, and what setting was captured. The score should lead back to examples. If a scale and the clinical account disagree, investigate the discrepancy rather than automatically privileging either source.

INSTRUMENT	CLASSIFICATION AND RESPONDENT	STRUCTURE STATED IN THE GUIDELINE
CARS-2	Screening, clinician rated	15 items rated 1-4
CAST	Screening, parent rated, ages 4-11 years	37 yes/no items
ASDS	Screening, parent rated, ages 5-18 years	50 yes/no items
Autism Behavior Checklist	Screening, parent rated	57 items rated 1-4

CARS-2 is classified as a screening measure, even though it is clinician rated. That distinction is worth making explicitly because clinician administration alone does not convert a scale into a diagnostic instrument. Likewise, the word “Diagnostic” in the Asperger Syndrome Diagnostic Scale can mislead: the guideline classifies it as screening. Candidates should classify an instrument by its validated clinical role, not by the confidence suggested by its name.

Caregiver questionnaires are especially helpful when clinic behaviour is unrepresentative of home or school. The commonly used instruments include the Social Responsiveness Scale, including SRS-2, and the Social Communication Questionnaire. They can focus the interview and identify domains requiring examples. They do not turn the caregiver into the diagnostician. The clinician's task is to determine the meaning of an endorsed difficulty, its developmental context, its consistency across settings, and the effect of communication level or situational stress.

24.5 ISAA, INCLN, and the Indian certification pathway

Diagnosis and certification answer different questions. The **Indian Scale for Assessment of Autism** is listed as a diagnostic instrument for children aged 2-9 years. Its stated format is a parent interview with 40 items rated 1-5. It produces cutoff scores, severity indices, and percentage-disability output that can assist certification. These capabilities make it important in Indian practice, but they should not be allowed to collapse the distinction between a clinical diagnosis and the later task of documenting disability for statutory purposes.

The ISAA is inbuilt within the INCLN Tool; this is a containment relationship, not a claim that the two names are interchangeable. The **INCLN Diagnostic Tool for Autism Spectrum Disorder** was released for uniform autism assessment in India by the Ministry of Social Justice and Empowerment, Department of Empowerment of Persons with Disabilities on April 25, 2016. In an examination answer, this is best presented as an Indian assessment and certification framework rather than as an alternative to clinical reasoning.

IN INDIA

Clinical diagnosis precedes disability assessment. After an established clinical diagnosis using recognised diagnostic criteria, the INCLIN tools and ISAA are used for disability assessment. It is therefore incorrect to write that a child “has autism because the ISAA is positive.” A scale result may support certification-related documentation, but the diagnosis must already rest on a clinical assessment. Conversely, an appropriate clinical diagnosis should not be withheld simply because certification has not yet been completed.

Certification quantifies disability and can facilitate scheme benefits and insurance eligibility. It therefore has material consequences for a family, and careful documentation is part of good clinical care. The required board comprises a paediatrician or paediatric neurologist, a clinical or rehabilitation psychologist, and a psychiatrist. This multidisciplinary composition reinforces the point that administrative assessment should be clinically anchored rather than reduced to a form-filling exercise.

Certificate validity depends on the nature of the disability. For an individual under 18 years of age with temporary disability, validity is 5 years; where a permanent disability has been acquired, the certificate has permanent validity. When discussing this with families, distinguish clearly between the diagnosis, the disability assessment, and the certificate. They are connected steps but not synonymous labels. The statutory context is the Rights of Persons with Disability Act, and the instrument's output should be explained as serving this practical pathway.

■ **TRAP Examination trap:** Do not use an ISAA or INCLIN disability assessment as though it replaces the clinical diagnosis of autism.

24.6 Integrating results in the consultation

A robust assessment is transparent about the status of each piece of evidence. State whether a result came from a parent screen, a caregiver questionnaire, clinician rating, direct observation, developmental-history interview, or disability assessment. Then state what it adds and what remains uncertain. This simple discipline prevents a referring clinician from reading a “positive” score as definitive and prevents a receiving clinician from discarding valuable information because it is not itself diagnostic.

- When a screen is positive, explain that it identifies a need for thorough diagnostic assessment and does not apply a diagnostic label.
- When a score is negative but surveillance remains concerning, return to the history, settings, and direct observation rather than treating the score as a veto.
- When trained diagnostic measures are used, document that their findings were integrated with clinical judgment rather than reported as a free-standing conclusion.
- When disability documentation is needed, establish the clinical diagnosis first and then proceed through the appropriate Indian assessment pathway.

The practical consequence is a more humane and accurate conversation with caregivers. Rather than saying “the test says autism” or “the test is normal,” the clinician can say what was observed, what the caregiver described, what the instrument contributes, and why further assessment or follow-up is appropriate. This approach also makes room for uncertainty without inaction. Where developmental concerns are real, referral and support planning need not wait for a family to understand every technical distinction among questionnaires, interviews, observation schedules, and certificates.


A positive autism screen changes the next clinical action; it does not, by itself, make the diagnosis.

25 · Medical comorbidities and workup in ASD (epilepsy, sleep, GI, genetics)

Autism does not protect a person from ordinary illness, and it can make illness harder to detect. A change in behaviour, communication, participation or self-care may be the presenting language of pain, sleep disruption, seizures, anxiety or another co-occurring condition. The assessment therefore has two linked tasks: recognise the developmental autism phenotype and ask what else may be contributing to current impairment. This is especially important when the person has limited spoken language, difficulty describing internal states, or a history in which distress has repeatedly been attributed to autism itself.

Co-occurring conditions are not a miscellaneous appendix to diagnosis. They alter risk, family burden, educational participation and the interpretation of behaviour. A useful formulation begins with the person's usual baseline and asks what has changed, in which setting, with what bodily or emotional correlates, and what reliably relieves or worsens it. The clinician should bring together caregiver observations, the person's own account where possible, developmental history, medication exposure, sleep and elimination history, dietary pattern, and a focused physical and neurological examination. The aim is not indiscriminate testing. It is a hypothesis-led medical workup that can explain a new problem and identify needs that autism alone cannot explain.

10-15% ESTIMATED EPILEPSY IN CHILDREN WITH ASD	up to 53% WITH A FREQUENT SLEEP PROBLEM	7-24% WITH SIGNIFICANT CHRONIC GASTROINTESTINAL COMPLAINTS	more than 50% ESTIMATED TO MEET ADHD CRITERIA IN SOME SAMPLES
---	--	---	---



25.1 A clinical method for finding comorbidity

Start with the referral question, but do not let it narrow the enquiry prematurely. A child described as "more autistic" may in fact be sleeping poorly, constipated, frightened, in pain, having unrecognised seizures, depressed, overwhelmed by a new demand, or experiencing an adverse effect of treatment. An adult who has always required support may develop a new loss of function that deserves the same respect as new loss of function in any other patient. Baseline is therefore a clinical instrument: establish usual communication, mobility, eating, sleep, continence, interests, activity level, self-injury, emotional expression and ability to tolerate ordinary transitions before interpreting change.

Behavioural equivalent is a practical term for a bodily or emotional problem expressed through observable behaviour when direct reporting is limited. It should prompt enquiry, not provide a conclusion. Agitation may reflect discomfort, fear, overstimulation, an interpersonal conflict, a seizure-related event or an emerging psychiatric syndrome. Similarly, apparent non-compliance can be a communication about an inaccessible demand or an aversive physical state. Ask caregivers for a sequence rather than a global adjective: what happens first, what follows, how long it lasts, whether awareness changes, and what restores the person to baseline.

CLINICAL SIGNAL	QUESTIONS THAT SHARPEN THE HISTORY	WHY IT MATTERS	IMMEDIATE ROUTE
New episodic change	Was there altered awareness, a motor event, a fall, a period of confusion, or a return to usual functioning?	Distinguishes a possible neurological event from a behavioural description alone	Document the event carefully and consider neurological assessment
New irritability or self-injury	What has changed in sleep, bowels, eating, activity, medication, demands or sensory load?	Physical discomfort and environmental strain are common explanatory possibilities	Examine, identify urgency, and investigate the leading hypothesis
Functional decline	Which specific skill has changed, how abrupt was the change, and is it present across settings?	Prevents a serious change being absorbed into a static developmental label	Assess medical, neurological and psychiatric contributors in parallel
Internalising distress	What situations are avoided, feared, anticipated or endured with marked distress?	Anxiety may be obscured by shutdown, rigidity, anger or somatic complaint	Make a symptom formulation before assigning a label

■ **RULE** Any marked change from baseline deserves a fresh medical and psychiatric formulation; do not accept autism as its explanation until competing explanations have been considered.

25.2 Epilepsy: recognise events, assess risk and use EEG selectively

Epilepsy is an important ASD comorbidity, but its clinical relevance lies in more than its frequency. Estimates in children with ASD place comorbid epilepsy between **10% and 15%**.

Another estimate for individuals with autism is about **20% to 25%**. These figures should not be treated as a prediction for an individual patient. They do, however, justify a low threshold for taking a paroxysmal history seriously, particularly when family descriptions include abrupt episodes that differ clearly from the person's usual repetitive behaviour, sensory response or emotional dysregulation.

The age pattern has clinical value. Onset is reported most often in infancy or adolescence, while onset in adolescence or early adult life is particularly distinctive when autism is compared with intellectual disability. The overall epilepsy rate in autism does not differ from that found in intellectual disability; the timing of onset is the more useful discriminating observation. A routine visit should therefore include a longitudinal question about events across development, rather than only asking whether seizures are occurring now.

Paroxysmal event is preferable to the word seizure until the history supports a conclusion. Ask whether the event occurs from sleep, during illness, after a particular trigger, or in a stereotyped sequence. Repetitive movements, staring, withdrawal, panic, dissociation and behavioural shutdown can all be misread if the sequence is not reconstructed.

Explain to families who expect a single investigation to explain autism or every behavioural difficulty that the question for the test is narrow: whether the event pattern raises a neurological concern that the EEG can help clarify. When events are recurrent or the account is unclear, contemporaneous documentation and neurological collaboration are often more informative than retrospective labels.

GOOD TO REMEMBER

When a new event is reported, reconstruct it with a witness before naming it a seizure: onset, behaviour, awareness, responsiveness, colour change, injury, duration, recovery and return to baseline. A caregiver-recorded video can improve phenomenology. An EEG may be useful when there is clinical evidence suggesting seizure activity; it does not replace a careful event history and must be interpreted in its clinical context.

Comorbid epilepsy is associated with greater intellectual disability and lower verbal ability. This association should guide anticipatory support, not deterministic counselling. It signals that communication access, safety planning and caregiver education may need particular attention. It does not mean that every person with epilepsy will follow the same developmental course, nor does it license therapeutic pessimism. Seizure management itself should be coordinated with the relevant neurological service; syndrome-specific seizure teaching belongs to the relevant syndrome sections rather than to a general ASD workup.

■ **TRAP** Examination trap: do not request an EEG merely because a person has ASD. Request it for a clinically credible paroxysmal history, and document the event phenomenology that created that suspicion.

25.3 Sleep disturbance: make the phenotype visible before treating it

Sleep problems are common enough to merit routine enquiry rather than waiting for a complaint. Up to **53%** of individuals with ASD have at least one frequent sleep problem. The common clinical patterns are **sleep-onset insomnia**, **sleep-maintenance insomnia** and **sleep-wake cycle irregularity**. Naming the pattern matters because a family report of "not sleeping" may refer to a prolonged settling period, repeated waking, a shifted schedule, an early waking pattern, or a combination.

Ask for the sequence around sleep rather than an estimate of total sleep alone. Establish the evening routine, time of attempted sleep, latency, night waking, nocturnal activities, waking time, naps, weekend variation, bedroom conditions, caregiver responses, daytime sleepiness and change from baseline. Ask directly about pain, bowel discomfort, breathing concerns, seizures, itching, medication timing, caffeine exposure, activity, screen use and anxiety. This identifies reversible drivers and clarifies whether the intervention must change the environment, the daytime rhythm, an associated condition, caregiver contingency, or a biological sleep signal.

Melatonin has shown benefit in **17 studies** in children with ASD. The dosing figures come from a narrower base and should be quoted as such: a meta-analysis of five of those studies is what reports good efficacy across doses of **1 and 10 mg**.

IN INDIA

For sleep disturbance in ASD, Indian guidance states a melatonin range between 1 and 10 mg at night.

Treatment periods in the cited evidence ranged from 14 days to over 4 years, which shows that use has not been confined to a single brief interval, not that every patient should receive long-term treatment.

Where epilepsy is also present, the guidance describes melatonin as carrying a low risk of seizure precipitation. This is reassuring but does not remove the need to document the seizure history, current neurological treatment and temporal relation between any new symptom and a medication change. The broader pharmacological management of target symptoms is addressed separately. Here the principle is to treat sleep as a clinical syndrome with contributors, not as an inevitable feature of autism.

For sleep, identify the pattern and its maintaining conditions before adding a treatment. A precise sleep history often prevents both under-treatment of discomfort and over-treatment of behaviour.

25.4 Gastrointestinal symptoms and bodily distress

Gastrointestinal complaint may present as a direct report, altered food intake, agitation around meals, a new sleep problem, avoidance of the toilet, or self-injury. Significant chronic gastrointestinal complaints are reported in between **7% and 24%** of individuals with ASD. It

should encourage active enquiry, not a presumption that every distress behaviour is gastrointestinal.

The clinical history should make eating and elimination discussable without embarrassment. Ask about appetite, food selectivity, texture or smell sensitivity, pain behaviour, swallowing concerns, nausea, vomiting, stool pattern, toileting avoidance, soiling, bleeding, weight change and the relation of symptoms to medication or routine disruption. In a person who cannot describe pain conventionally, a caregiver's recognition of a distinct grimace, posture, guarding, meal-related distress or night waking may be the most useful starting point. Examination and referral should be guided by the symptom pattern and urgency, as in any patient.

It is useful to separate a sensory preference from a medical symptom without making them adversaries. A narrow diet may be a stable sensory accommodation, a consequence of pain, an anxiety-driven avoidance, a learned family adaptation, or several of these together. The same is true of toileting refusal. The formulation becomes clinically useful only when it identifies function, discomfort, nutritional impact and the practical barriers to change. Abrupt dietary restriction, apparent pain, dehydration, bleeding, persistent vomiting or a significant decline in functioning should not be normalised as autism.

- Ask directly about bowel, bladder, appetite and pain at each meaningful change in behaviour.
- Use familiar caregiver observations as data, then test them against a focused history and examination.
- Distinguish a long-standing preference from a new restriction or a new distress signal.

25.5 Genetic and general medical workup: phenotype-led, not reflexive

Phenotype-led genetic evaluation means that genetic assessment is anchored to the individual rather than to a generic autism label. The history should include developmental course, medical events, neurological symptoms, family history and any pattern suggesting a broader developmental condition. The examination should attend to growth, physical features, skin, neurological signs and other findings that may make a syndromic explanation more likely. This approach improves the quality of the referral question and helps the family understand why genetic assessment is being considered.

Genetic testing is part of the ASD medical workup, but its value is not exhausted by producing a laboratory result. A result may refine an aetiological formulation, prompt surveillance for associated medical issues, inform reproductive discussion through an appropriate genetics service, and connect a family with relevant supports. Conversely, an uninformative result does not invalidate the person's autism diagnosis or their need for support. Consent should cover what the investigation can and cannot answer, the possibility of uncertain findings, and the implications of receiving information that concerns relatives.

The psychiatrist's role is to recognise when a broad medical formulation is needed, obtain a careful developmental and family history, coordinate referrals and translate results into a person-

centred plan. It is not to assign a syndrome from a single physical feature or to promise that testing will identify a cause in every case.

- Record the phenotype and the family question that prompted referral.
- Explain what a result may clarify and what it may leave uncertain.
- Coordinate interpretation with genetics and other relevant services.

Hearing and vision concerns, nutritional status, sleep, bowel function, medication effects and neurological symptoms should be considered alongside genetic assessment. This prevents the common error of pursuing an etiological explanation while missing a treatable contributor to day-to-day distress. A medical workup is successful when it makes the clinical picture more intelligible and improves care coordination, even when no single cause is established.

25.6 AuDHD: distinguish co-occurrence from diagnostic substitution

AuDHD refers to the co-occurrence of autism and ADHD. ADHD is one of the most common comorbidities in ASD, and some estimates suggest that **more than 50%** of individuals with ASD also meet criteria for an ADHD diagnosis. The clinical task is not to decide which label "explains everything". It is to establish whether attentional difficulties, impulsivity or hyperactivity represent an additional, impairing syndrome rather than an understandable consequence of overload, anxiety, poor sleep, learning demands or the social environment.

This is a major nosological correction for candidates to understand. DSM-IV-TR did not permit concurrent ASD and ADHD diagnoses; DSM-5 permits co-occurring diagnoses. The earlier exclusion still shapes clinical habits. In one direction, marked activity or distractibility is dismissed as part of autism. In the other, social difficulty is assumed to follow from ADHD alone. Both shortcuts can delay a formulation that makes sense of the person's whole pattern.

Assess attention and activity across settings and against the person's developmental level. A concurrent ADHD diagnosis should be considered when attentional difficulties or hyperactivity exceed what is typical for individuals of comparable mental age. Ask what happens in a low-demand preferred activity, in an unstructured group setting, during a task that is understood but not preferred, and after a poor night of sleep. Gather collateral from the settings where demands differ. This separates sustained regulation difficulty from a problem that appears only under a particular sensory, social or academic load.

The co-occurrence changes management planning because sleep, anxiety, communication demands and environmental predictability can all alter attention and activity. Establish the target impairment, its baseline, the family's priorities and the measures by which benefit and adverse effects would be judged. Pharmacological treatment selection for target symptoms is addressed in the pharmacotherapy section; ADHD itself, including adult persistence, is addressed in the ADHD and specific learning/communication disorders notes. The key point here is that dual diagnosis is clinically permissible and often clinically clarifying.

MNEMONIC**Clinical spine - CHANGE**

When behaviour worsens, check **C**ourse from baseline, **H**ealth and pain, **A**rousal and sleep, **N**eurological events, **G**astrointestinal and other bodily symptoms, and **E**motional or attentional comorbidity. It is a sequence for broadening the formulation, not a substitute for assessment.

25.7 Anxiety, obsessive-compulsive phenomena and catatonia

Anxiety disorder is common in ASD, with a more conservative estimate of **43%** in individuals with ASD. Social anxiety and specific phobias tend to be the most common subtypes. The assessment must not mistake all avoidance for autism-related social difference or all insistence on sameness for anxiety. Ask what the person fears will happen, what they anticipate, whether distress appears before the event, what safety behaviours are used, and whether the pattern is new, pervasive or situation-bound.

Obsessive-compulsive symptoms demand a similarly functional distinction. Repetitive acts in autism may be enjoyable, regulating or linked to predictability, whereas compulsions are more often driven by distress and an attempt to reduce a feared outcome. The distinction can blur, particularly when the person cannot easily report intrusive thoughts. Explore the emotional state before and after the act, the consequences of interruption, the degree of choice, and whether the behaviour has acquired a feared meaning. Nearly 5% of autism cases have comorbid OCD, so the possibility should be actively considered rather than hidden within a description of repetitive behaviour.

Catatonia requires recognition because new marked slowing, reduced initiation, mutism, posturing, agitation, loss of previously established function or other conspicuous motor-behavioural change may be misattributed to autism. The core safeguard is temporal: a substantial change from the person's own baseline is a clinical event. Obtain a careful timeline, assess medical and neurological contributors, seek collateral, and involve appropriate specialist services where concern is significant. Do not wait for a person to fit a stereotyped image of catatonia before taking severe change seriously.

■ **TRAP** Examination trap: repetitive behaviour is not automatically OCD, and withdrawal is not automatically autism. Establish the preceding emotion, perceived purpose, degree of distress and change from baseline.

25.8 Integrating findings into a safe, useful formulation

The completed assessment should answer practical questions: what is longstanding, what is new, what is impairing, what needs urgent medical attention, and which professional needs to act next? Document observations in behavioural language, preserve uncertainty where it remains, and specify the evidence supporting each concern. A formulation such as "new nocturnal waking with apparent abdominal discomfort and daytime irritability" is more actionable than "behavioural problem in ASD". It allows the family, psychiatrist, paediatric or medical colleague, school and therapist to work from the same clinical picture.

Finally, explain the formulation to the person and family without implying blame. Families often arrive after being told either that every problem is autism or that autism explains nothing. Both positions are unhelpful. Autism provides a developmental context. It does not erase the need to recognise pain, epilepsy, sleep difficulty, gastrointestinal symptoms, genetic questions, ADHD, anxiety, OCD or catatonia. Care improves when the clinician can hold both truths at once: the person's baseline neurodevelopmental profile matters, and a real change from that baseline always deserves to be understood.

Managing autism and its outcome

26 · Differential diagnosis of ASD

Differential diagnosis is a longitudinal task. Autism spectrum disorder (ASD) is not established by an isolated socially atypical behaviour, a language delay, or a repetitive act observed in clinic. The clinician asks whether the social-communication pattern is pervasive, whether restricted or repetitive behaviour is present across development, what the child could do before the concern emerged, and whether the proposed alternative explains the whole presentation more parsimoniously. The same exercise determines whether an apparent alternative is instead a co-occurring condition that deserves its own diagnosis and treatment plan.

For examination answers, organise the differential around the **developmental trajectory**: an early and pervasive neurodevelopmental pattern supports ASD; a setting-bound difficulty, a later change after relatively typical development, or a symptom driven by fear, psychosis, sensory loss, neurological illness, or adverse care conditions should redirect assessment. Collateral developmental history, direct observation across settings, and attention to communication without spoken language are often more informative than a checklist answer. The task is not to make every unusual behaviour autistic, nor to deny ASD when another condition is also present.

-
- **RULE** **Core rule.** Diagnose ASD when its characteristic social-communication and restricted or repetitive pattern is the best account of the developmental presentation; add another diagnosis when that condition independently meets its threshold or requires separate clinical attention.
-


The differential of developmental regression

Two discriminators do the work: when the loss starts, and what is lost. Every age window on this plate is one a source states; none is interpolated.

Not sourced, and therefore not drawn: a metabolic branch point. No source used in this chapter names a metabolic disorder in its regression differential, gives an age window for metabolic regression, or gives a diagnostic yield for metabolic testing in regression. ICD-11 names Landau-Kleffner syndrome, autoimmune encephalitis and Rett syndrome. DSM-5-TR names continuous spike and waves during slow-wave sleep and Landau-Kleffner. Neither names a metabolic cause. The metabolic-yield figures this chapter carries are bound to the ID workup, not to regression, so they are not drawn here.

Discriminator 1. When the loss starts.

Each bar is an age window a source states. The two coral lines are the alarm lines: the age past which each system stops calling the regression ordinary.



The ASD window is the same window in both systems, stated in different units: ICD-11 gives 1 to 2 years, DSM-5-TR gives 12 to 24 months. Rett's normal period and its regressive phase come from different sources and different constructs, so they are drawn as two bars and they overlap. The CDD bars are Dulcan's.

Discriminator 2. What is lost.

Loss confined to language and social responsiveness fits ASD. Loss that reaches beyond social communication does not, and that is the branch point.

ASD with loss of skills Language use and social responsiveness, most often. Regression or stagnation is reported in about 20% of cases, and that figure covers plateau as well as loss, not loss alone.	Rett syndrome Purposeful hand use, replaced by stereotyped hand-wringing. Head growth decelerates to an acquired microcephaly, a primary criterion by 2 y. Social interest returns after the regression.
Regression after at least 2 y (previously CDD) Losses beyond social communication: self-care, toileting, motor skills. DSM-5-TR calls this much more unusual and warranting more extensive investigation. Prevalence 1.7 per 100,000 (95% CI 0.6 to 3.8).	Acquired epileptic aphasia (Landau-Kleffner) Receptive AND expressive language, lost precipitously over days or gradually over months. 30 to 50% have no overt seizures, at least initially.

The alarm line sits one year apart in the two systems, and an MCQ can turn on exactly this.
ICD-11: loss of previously acquired skills is reported in the second year of life in some children with ASD, but rarely occurs after the age of 3 years.
DSM-5-TR: in rare cases regression occurs after at least 2 years of normal development, which is much more unusual and warrants more extensive investigation.
Both call late regression unusual, but they place the line a year apart and they name different investigation sets. ICD-11 names Landau-Kleffner syndrome, autoimmune encephalitis and Rett syndrome. DSM-5-TR names continuous spike and waves during slow-wave sleep and Landau-Kleffner, cross-referring Rett.

The rule, when the loss is not ASD's
ICD-11 discriminates diseases of the nervous system from ASD-with-loss-of-skills on two things: an early history of relatively normal social and language development, and the characteristic neurological features of those disorders, which are not typical of ASD. Neither test is an age. Both are history.

- Petrol. Structure: an ordinary category in this differential, and an age window a source states. A routine branch is never coloured as a danger.
- Petrol wash, dashed. A stated window that nests another of its own construct: the permitted 2 to 10 year window inside which CDD typically starts.
- Amber. A key number, carried in the gutter beside its own bar, so that no number sits on a rule or inside a bar; and the frame of the conflict box.
- Coral. A sourced danger signal (the alarm lines, the seizure-free Landau-Kleffner trap) and, dashed, the honesty marker for what the sources do not give.

Ages are exactly as each source states them. Nothing is converted, interpolated or rounded, and no bar is drawn for a window the sources do not give.
The ASD window and both ICD-11 statements are ICD-11 CDDR. The Rett regressive phase, the DSM alarm line and the regression-after-2-years rule are DSM-5-TR 2022.
Rett's apparently normal period is Dulcan, Rutter's 6E, Companion and Kaufman converging. The CDD windows and both Landau-Kleffner figures are Dulcan's Textbook. The 20% regression-or-stagnation figure is Kaplan & Sadock's CTP 2024, and it covers plateau as well as loss.
Nosological caution. DSM-5-TR states that ASD encompasses childhood disintegrative disorder, so the third branch is drawn as the phenomenon DSM-5-TR names, regression after at least 2 years of normal development. Dulcan's 'age at onset alone' discriminator is DSM-IV-era and is not a current DSM-5-TR differential.

Fig 21.19 The differential of developmental regression. The branch points that separate autism with loss of skills, Rett syndrome, regression after at least two years (previously childhood disintegrative disorder) and acquired epileptic aphasia, drawn on the two discriminators that actually do the work: the age at which the loss starts, and what is lost. The ruler carries only age windows a source states, and the alarm lines show ICD-11 and DSM-5-TR placing the boundary of the ordinary a full year apart. A metabolic branch point is named on the banner and left undrawn: no source used here supplies one, and the metabolic yields this chapter carries are bound to the intellectual-disability workup rather than to regression.

26.1 A practical frame for the differential

Start by examining social communication, restricted or repetitive behaviour, pervasiveness across settings, and any loss of acquired capacity. Was early social communication qualitatively atypical, or merely delayed in proportion to overall development? Are repetitive behaviours, unusual interests, inflexibility, or sensory-linked patterns truly present, including historically? Is the pattern evident across relationships and environments? Does the time course suggest ASD or a neurological, psychiatric, or environmental cause? This frame prevents the common error of treating an isolated surface feature as diagnostic.

Use function, not appearance alone. Repetition may be self-organising, enjoyable, anxiety-reducing, an attempt to avoid an aversive demand, or a response to an intrusive thought. Social withdrawal may reflect a pervasive social-communication difference, fear of scrutiny, low mood, psychosis, or an attachment-related adaptation. The meaning of the behaviour, its developmental onset, and what changes it are more discriminating than the behaviour's name.

- Obtain a timeline of social reciprocity, communication, play, interests, daily routines, and any losses of skill.
- Ask about the same behaviour at home, school, and with familiar and unfamiliar people.
- Separate a core ASD feature from a secondary reaction to anxiety, frustration, communication failure, or environmental change.
- Look for independent evidence that an alternative disorder is present, rather than using it as a label for a difficult symptom.

■ **TRAP Examination trap.** A child who does not speak in a particular setting does not automatically have ASD. Pervasiveness of the social-communication difficulty and the presence or absence of restricted or repetitive behaviour decide the direction of enquiry.

26.2 ADHD and intellectual developmental disorder

ADHD and ASD can be difficult to discriminate. Interrupting, intruding into personal space, excessive volume, impatience, and poorly modulated interaction can create social difficulty in ADHD. These behaviours should not be mistaken for the qualitative reciprocity and communication pattern of ASD. The discriminating history is the developmental course, together with whether restricted or repetitive behaviours and unusual interests are absent. When they are absent and the social difficulty tracks inattention, impulsivity, or hyperactivity, ADHD is the better explanation.

These disorders may also coexist. In a person with ASD, a concurrent ADHD diagnosis should be considered when attentional difficulty or hyperactivity exceeds what is expected at a comparable mental age. This is a clinical comparison against developmental capacity, not a shortcut based on chronological age. Do not use the presence of ASD to explain away disabling inattention, and do not use the presence of ADHD to dismiss a developmental history of restricted interests and pervasive social-communication difference. ADHD itself is covered in the ADHD and specific learning/communication disorders notes.

The boundary with intellectual developmental disorder demands the same developmental comparison. In intellectual developmental disorder without ASD, social-communicative skills generally do not show an apparent discrepancy from other intellectual skills. ASD is justified when social communication and interaction are substantially more impaired than the developmental level indicated by nonverbal abilities. Observe how the child uses gaze, gesture, affect, shared attention, and nonverbal problem solving, rather than equating sparse speech with autism. This distinction is especially difficult before symbolic communication has emerged and when repetitive behaviour occurs in the context of developmental disability.

ICD guidance makes the same point operationally: assign both diagnoses, with the relevant specifier, when social-communication deficits are greater than would be expected from intellectual functioning. In this setting, adaptive assessment should not let ASD-related social difficulty artificially inflate the apparent global developmental impairment. The conceptual and practical evidence, the developmental record, and repeated assessment matter greatly. The full differentiation of intellectual disability, global developmental delay, and borderline intellectual functioning is addressed elsewhere in this chapter.

26.3 Language, pragmatic communication, and selective mutism

A **developmental language disorder** can produce frustration, reduced participation, and secondary social difficulty. It is not usually accompanied by abnormal nonverbal communication or restricted and repetitive patterns. A child with language disorder may therefore show a strong wish to engage, communicative use of gesture and facial expression, and reciprocal response despite impaired spoken language. The assessment should be designed so that social intent can be seen without demanding verbal fluency.

Social (pragmatic) communication disorder is the nearer diagnostic neighbour because social use of communication is impaired. The separator is the absence of restricted or repetitive behaviour and interests. Before making this diagnosis, enquire carefully about such behaviour in both the past and the present. If ASD criteria are met, ASD takes diagnostic precedence; it is not appropriate to divide an ASD presentation into separate social-communication and repetitive-behaviour diagnoses. Conversely, pragmatic language impairment alone does not justify adding developmental language disorder to ASD. Individuals with a primarily pragmatic language disorder may still initiate and respond appropriately to social and emotional cues and share interests, while lacking the typical restricted or repetitive pattern seen in ASD.

Selective mutism is a disorder of context, not a global absence of communication. Early development is not typically disturbed; communication can be appropriate in some settings; and even where the child is mute, reciprocity is not impaired and restricted or repetitive patterns are not present in the way expected in ASD. Ask what the child does at home, with familiar relatives, and in school. Selective mutism shows normal language and social communication in specific environments but not others, whereas ASD-related social-communication deficits and restricted or repetitive patterns are evident across situations and contexts. Anxiety and ASD can coexist, so a clear setting effect should prompt assessment for both rather than a hurried either-or decision.

PRESENTATION	QUESTION THAT RESOLVES IT	PATTERN FAVOURING THE ALTERNATIVE
Developmental language disorder	Can social intent be demonstrated nonverbally?	Reciprocal nonverbal communication without the restricted or repetitive ASD pattern.
Social pragmatic communication disorder	Has restricted or repetitive behaviour ever been present?	Social-communication impairment without that behavioural pattern.
Selective mutism	Does communication become normal in a safe, familiar setting?	Context-specific mutism with preserved reciprocity.

26.4 Attachment-related presentations, deprivation, and disruptive behaviour

Severe social deprivation can produce **quasi-autism**, an autism-like presentation that must not be assumed to establish a lifelong neurodevelopmental diagnosis without careful history. A key discriminating observation is substantial improvement after movement to a more nurturing environment. Where there is no reliable account of earlier social and communicative development, the distinction may remain difficult. Longitudinal observation is then clinically more honest than premature certainty.

Reactive attachment disorder and ASD share social withdrawal and low reciprocity, but their organising patterns differ. ASD shows atypical language development, whereas delayed language in reactive attachment disorder is not stereotyped or otherwise deviant; restricted or repetitive behaviour is not a feature of reactive attachment disorder; and children with ASD can show attachment behaviour with caregivers at least some of the time despite their social-communication difficulty. A history of severe neglect or maltreatment is required for reactive attachment disorder and is very unusual in ASD. The diagnosis should therefore rest on both the caregiving history and the behavioural phenotype, not on either alone.

Disinhibited social engagement disorder may also enter the differential when a child approaches unfamiliar adults indiscriminately. That feature is not decisive, because some children with ASD may also show it. Restricted, repetitive, and inflexible patterns are not typical of disinhibited social engagement disorder, and a marked reduction in symptoms after a nurturing environment is provided supports that diagnosis. The clinical task is to avoid a false split between adversity and neurodevelopment: adversity may be clinically crucial whether or not ASD is ultimately present.

Oppositional behaviour needs a functional formulation. In oppositional defiant disorder, the social-communication and restricted or repetitive pattern characteristic of ASD is absent. Explosive rages in ASD are often linked to a specific trigger - a disruption of routine, aversive sensory input, anxiety, or interruption of a rigid sequence - rather than a wish to be provocative or spiteful as required for an oppositional account. Ask what occurred immediately before the outburst and what the child was trying to preserve, escape, or communicate. Demand avoidance

may be prominent in ASD, but it should not be mistaken automatically for a separate disruptive disorder.

26.5 Repetitive behaviours: OCD, stereotypies, and tics

The same act can look repetitive while serving very different psychological functions. In **obsessive-compulsive disorder**, compulsions are performed in response to intrusive thoughts, commonly to relieve anxiety. In ASD, repetitive motor acts, insistence on sameness, or ordering behaviour may be experienced as pleasurable and reinforcing. Ask the person, in developmentally appropriate language, what happens if the behaviour is interrupted: is there fear of a feared consequence, a drive for relief or completeness, sensory regulation, enjoyment, or distress at change? The answer is informative but not self-sufficient, because insight and the ability to report internal experience vary.

Resistance is another useful but imperfect axis. People with OCD more often consciously resist compulsive urges, whereas the characteristic social-communication deficits of ASD are not features of OCD even when repetitive behaviour is prominent. Adolescents and adults with ASD may nevertheless attempt to suppress behaviour they know is socially unwelcome. Therefore, neither resistance nor lack of resistance should be converted into an isolated diagnostic test. Evaluate the cognitive antecedent, emotional consequence, developmental context, and the rest of the ASD phenotype.

Stereotypic movement disorder is not added merely because repetitive movements are visible in someone with ASD. A separate diagnosis is not made when the movement is better explained by ASD; it can be appropriate when stereotypy causes self-injury and becomes a specific treatment focus. The practical point is to describe the movement carefully, assess injury and impairment, and decide whether it needs additional clinical attention rather than multiplying labels.

Differentiate stereotypies from **tic disorders** by phenomenology. Tics tend to be less stereotyped, briefer, later in emergence, and associated with premonitory sensory urges; unlike ASD repetitive behaviours, they are not generally experienced as soothing. Direct questioning about an urge, temporary suppression, fluctuation, and relief is useful when language permits. Co-occurrence remains possible, so the aim is accurate description and not forced diagnostic exclusivity.

MNEMONIC

Meaning before movement

For any repetitive act, ask: what precedes it, what does it achieve, what happens if it is blocked, and does the broader social-developmental pattern support ASD? This is a reasoning sequence, not a diagnostic acronym.

26.6 Anxiety, psychosis, and personality pathology

Anxiety disorders overlap with ASD because social withdrawal and repetitive behaviours can be core ASD features but can also express anxiety in either presentation. The central question is

whether social functioning becomes broadly intact when anxiety is low and the context is familiar. In **social anxiety disorder**, interaction with familiar people or in non-anxiety-provoking situations does not show the pervasive impairment expected in ASD; persistent restricted, repetitive, or inflexible patterns are also not features of social anxiety alone. A patient may have both conditions, and avoidant behaviour should then be formulated in terms of both developmental style and fear.

Childhood-onset **schizophrenia** is usually preceded by normal or near-normal development, and hallucinations and delusions are defining features that do not belong to ASD. Its prodrome can nonetheless resemble ASD through social impairment, unusual interests, or odd beliefs.

Questions about psychotic symptoms require concrete follow-up: an affirmative literal answer may refer to ordinary media rather than a perceptual experience. Establish the source, location, controllability, conviction, and functional consequence of the reported experience. If ASD and schizophrenia each meet criteria, diagnose both rather than making one erase the other.

In adults without major language or intellectual impairment, personality disorder can resemble ASD through detachment, eccentricity, unusual preoccupations, or restricted affect. The most helpful discriminator is the early developmental course: lack of imaginative play, restricted or repetitive behaviour, and sensory sensitivity in childhood point toward ASD rather than a personality-based explanation. Adult social style alone is an unreliable retrospective diagnostic shortcut. Seek informant history and records where available, and distinguish a lifelong developmental pattern from interpersonal adaptations that emerged later.

26.7 Regression and neurological red flags

Regression changes the diagnostic posture. Loss of skills can occur in ASD. The syndrome-specific accounts belong with their dedicated discussion.

GOOD TO REMEMBER

In ASD, loss of skills is reported in the **second year of life** in some children and rarely occurs after the age of **3 years**. A later loss, especially after a relatively normal early social and language history, should trigger a differential that includes acquired epileptic aphasia, autoimmune encephalitis, Rett syndrome, and other neurological causes.

Medical causes are distinguished from ASD with loss of skills through the history of relatively normal development and the neurological features that are not typical of ASD on their own. Look beyond social communication. Loss of self-care, toileting, motor ability, or other acquired competence makes an encephalopathic process more concerning and warrants more extensive medical investigation than an uncomplicated ASD assessment. Where autistic features are attributable to a medical condition, ICD guidance allows a secondary neurodevelopmental syndrome diagnosis rather than ASD. The distinction matters because immediate management and prognosis can change.

Landau-Kleffner syndrome is an important mimic when apparent autism follows loss of receptive and expressive language after a period of normal development. That period may be **3 to**

7 years. An absent seizure history does not exclude it: **30 to 50%** of affected children may initially have no overt seizures. A careful neurological history and appropriately directed investigation are therefore essential in acquired language regression.

30 to 50%

LANDAU-KLEFFNER SYNDROME MAY INITIALLY LACK OVERT SEIZURES

26.8 Sensory impairment and closing clinical synthesis

Severe hearing impairment, severe visual impairment, maltreatment, developmental coordination disorder, epileptic encephalopathy, mood disorder, conduct disorder, and other behavioural conditions belong in a broad ASD differential when the presentation requires it. Their presence should slow diagnosis rather than become a reason to abandon developmental assessment. Hearing assessment is a named component of ASD workup, and audiology with speech and language expertise belongs in the multidisciplinary team for this reason. The technical differentiation of hearing impairment from developmental language disorder should not be imported uncritically as an ASD rule.

End with a positive formulation: which features establish ASD, which features point to a differential, which conditions may coexist, and what further observation or investigation is required before certainty is justified. This separates a thoughtful assessment from a list of labels. It also protects families from both diagnostic delay and diagnostic overreach. The full child psychiatric assessment battery, services and protection are covered in the Child psychiatry: assessment, treatment, services and protection notes.

In ASD differential diagnosis, developmental course, pervasiveness across settings, the function of repetitive behaviour, and evidence of neurological regression are more useful than any isolated symptom.

27 · Rett syndrome and childhood disintegrative disorder

Why these conditions matter. Both **Rett syndrome** and childhood disintegrative disorder can present to the psychiatrist as a child who has lost development and now appears autistic. That surface resemblance is clinically consequential but diagnostically incomplete. Rett syndrome is a distinctive genetic and neurological condition with a characteristic course and physical phenotype. Childhood disintegrative disorder is a historical diagnostic construct describing unusually marked regression after a prolonged period of apparently typical development. A careful longitudinal history prevents the error of treating every regression as simply another presentation of autism.

The exam task is not merely to recall labels. It is to identify the pattern of loss, look for the accompanying neurological signs, understand what has changed in current classification, and arrange a proportionate medical assessment. The broader differential diagnosis of developmental regression is addressed in the differential diagnosis of ASD section; here the emphasis is on the two named entities and on the clinical reasoning that separates them from ASD.

99% RETT CASES DUE TO SPORADIC MUTATION	about 1 per 10,000 REPORTED PREVALENCE IN GIRLS	four CLASSIC RETT COURSE PHASES	1.7 per 100,000 CDD PREVALENCE ESTIMATE
--	--	--	--

27.1 Rett syndrome: identity, genetics and diagnostic stance

Rett syndrome was first described by Andreas Rett, an Austrian paediatrician, in Austria in 1966. It is an X-linked dominant condition caused by pathogenic change in MECP2, located at Xq28. MeCP2 binds methylated DNA and participates in the regulation of gene expression and chromatin structure. Its loss disrupts neuronal maturation and synaptogenesis. This mechanistic framing is useful because it explains why the syndrome is not just a behavioural regression syndrome: its evolution includes motor, autonomic, skeletal and epileptic manifestations.

Most cases arise anew rather than being inherited. The proportion attributed to a **99%** sporadic mutation is a memorable genetic fact, while the wider clinical point is that family history may be unrevealing. The syndrome is reported chiefly in females. A hemizygous mutation has been described as incompatible with life in a male fetus, but this should not be recited as an unqualified explanation for the sex distribution because the literature contains a contrary account of that mechanism. The clinically safer formulation is that rare affected males may have severe neonatal encephalopathy, and rare male cases have been reported with a concomitant XXY abnormality.

The syndrome is a major cause of intellectual disability in girls, described as the second most important cause after Down syndrome. Frequency figures require care because sources use different denominators and epidemiological constructs. A reported prevalence is **about 1 per 10,000 girls**; an incidence estimate is approximately 1 per 10,000 female live births. These should never be swapped in an answer. In practice, their shared message is rarity combined with high clinical importance.

■ **RULE** **Rett syndrome remains a clinical diagnosis.** Consensus clinical criteria establish the syndrome; genetic testing supports the diagnosis but does not establish it in isolation.

This principle avoids two errors. First, a compatible molecular result does not replace phenotype and developmental trajectory. Secondly, a negative or non-diagnostic result must not cause a clinician to abandon a convincing clinical syndrome. Variant forms exist, including a very rare preserved speech variant in which mutation carriers may communicate well and have mild intellectual disability; asymptomatic carriers have also been reported. Such variation makes clinical synthesis more important, not less.

27.2 The Rett developmental trajectory

The diagnostic spine is a period of apparently ordinary early development followed by loss of acquired skills and a changing neurological phenotype. The early apparently normal period is reported as **6-18 months**. Delays may become apparent around 6 months, but this is not a substitute for listening closely to the parents' account of the developmental sequence. In the

regressive phase, typically between ages 1 and 4 years, a child may look strikingly autistic. The key question is what happens next, and what accompanies the loss.

The course is conventionally understood as **four** classic phases. Rather than forcing stage labels onto a history, describe the sequence: an early apparently normal period, a short plateau, rapid deterioration of motor and speech abilities, a later plateau that can last for years, and subsequent deterioration of the motor system. This sequence is more clinically useful than a list of labels because it connects the family narrative to examination findings and anticipates changing care needs.

Loss of **purposeful hand use** is particularly discriminating. It is replaced by stereotyped hand-wringing, hand-washing, hair-pulling, clapping, flapping or twisting movements. These movements are not an incidental repetitive behaviour. They represent the substitution of a stereotypy for previously functional manual activity, and therefore should be elicited with a before-and-after history. Hand stereotypies first appear at about 18 months. Bruxism, mouthing and body twisting can accompany the hand movements.

Head growth is also informative. Head circumference can be normal at birth, then head growth decelerates, producing **acquired microcephaly**. The adjective matters: it distinguishes a postnatal deceleration from a head that was already small at birth. Significant microcephaly has been stated to emerge by 2 years of age, although head-growth slowing is not universal. A normal head-growth record therefore does not by itself dispose of the diagnosis; it makes the whole pattern, rather than one sign, more important.

■ **TRAP Examination trap.** Do not call a child autistic solely because she has social withdrawal, language loss and stereotypies during Rett regression. Ask specifically about loss of purposeful hand use, slowing of head growth, gait change and the later social course.

There is a further reason this distinction matters. Social interest often diminishes during the regressive period but later improves; many affected individuals develop an intense eye gaze or staring. Thus, an early autistic appearance need not remain the dominant social phenotype. In the current manual, ASD should be considered in Rett syndrome only when all ASD diagnostic criteria are independently met. This is a disciplined use of co-diagnosis, not a reflexive relabelling of the syndrome.

27.3 Neurological, autonomic and systemic phenotype

The neurological examination should deliberately seek a pattern, rather than recording isolated abnormalities. Gait dyspraxia or apraxia, truncal ataxia, gait and trunk movement apraxias, muscle wasting and dystonia are described in Rett syndrome. These features change the clinical question from “is this autism?” to “what neurological disorder is producing regression and autistic-appearing behaviour?” A child who can no longer use her hands purposefully and develops an abnormal gait demands this broader formulation.

Breathing abnormalities are another helpful clue. Episodes of hyperventilation and breath-holding can alternate. Their importance is practical as well as diagnostic: they should not be

assumed to be wilful behaviour or anxiety without a careful clinical account. Reflux and swallowing problems may reflect autonomic pathology. Acrocyanosis and trophic skin changes of the feet add to the autonomic and vascular phenotype. These physical features often explain why a narrowly psychiatric interview is insufficient.

Epilepsy is a frequent complication, but numerical statements in this area differ between sources. One source reports over 90% of girls with Rett syndrome having epilepsy, with seizures usually beginning around 4 years; another reports a lower range beginning at a different age. The defensible clinical learning point is that seizure history must be actively sought and that later changes in behaviour, alertness or function should not automatically be attributed to the developmental disorder. The Rett-specific seizure burden belongs in the medical formulation, not as a generic ASD comorbidity statistic.

Scoliosis is common over the course of the syndrome, reported in about 87% of females by 25 years of age. Sleep disturbance also deserves direct inquiry: 85% of preschool children have been reported to have disturbed sleep with night waking and paroxysmal or continuous laughter. These are not peripheral details. Sleep disruption, immobility, seizures, feeding difficulty and spinal deformity can each alter daily function, caregiver strain and the safety of medication choices.

Cardiac risk requires particular prescribing attention. There is an increased risk of arrhythmias associated with a **prolonged QT interval**. In Rett syndrome, avoid prokinetic agents, antipsychotics, tricyclic antidepressants, certain antiarrhythmics, anaesthetic agents and certain antibiotics where this risk is relevant. This is not an instruction to leave distressing symptoms untreated. It is a reminder that a psychiatric prescription belongs within a medical plan that considers cardiac conduction and concurrent treatment.

Despite the apparent downhill trajectory, Rett syndrome is **not a neurodegenerative disorder**. The neuropathological account is one of developmental arrest in infancy rather than progressive inflammatory, demyelinating or degenerative destruction. That distinction protects against a common conceptual mistake: later motor deterioration is real, but it does not prove a conventional neurodegenerative process.

27.4 Classification and the autism boundary

Rett syndrome is no longer included as an ASD or pervasive developmental disorder subtype in DSM-5. This differs sharply from childhood disintegrative disorder, which was absorbed into ASD. The historical placement of Rett beside autistic disorders should therefore be taught as classification history, not current nosology. The separation has validating evidence: MeCP2 mutation is not found in other autism spectrum disorders, making Rett a distinct genetic and neurological entity rather than simply a severe ASD subtype.

The current ASD diagnosis nevertheless contains a useful documentation route. The specifier **associated with a known genetic or other medical condition or environmental factor** can be applied when the individual has a known genetic condition, with Rett syndrome named as an example. The specifier does not itself claim that every aspect of the autistic presentation is

caused by the genetic condition. It documents clinically relevant context while preserving the requirement to establish ASD on its own criteria.

In ICD-11, Rett is named as a boundary or differential condition rather than as an ASD category. Diseases of the nervous system with regression are distinguished from ASD with loss of skills by an early history of relatively normal social and language development and by characteristic neurological features. This is exactly the clinical distinction the psychiatrist should make at the bedside. The legacy codes are F84.2 in ICD-10 and 299.80 in DSM-IV or DSM-IV-TR; do not use them as current ASD subtype labels.

A child with regression and autistic-appearing behaviour needs a syndromic neurological formulation: in Rett syndrome, social interest may later improve, whereas the hand, head-growth, motor and autonomic phenotype remains diagnostically central.

27.5 Childhood disintegrative disorder: the historical construct

Childhood disintegrative disorder is also known as Heller syndrome or Heller's disease and was originally termed dementia infantilis. It described a child with a substantial period of apparently normal development followed by marked loss of skills and the emergence of an autistic clinical picture. The eponym is worth recognising because older examinations, case records and ICD-10 terminology may use it, but current clinical reasoning should not mistake a legacy label for a separate contemporary ASD category.

The historical criteria required at least **2 years** of apparently normal development before regression. Regression begins no earlier than 2 years and no later than 10 years; the most typical onset is between 2 and 4 years. These constructs are easy to confuse. The first specifies the preceding normal developmental interval, the second the permitted boundaries for regression, and the third the modal onset. In a vignette, establish the timing before deciding that the term fits.

Under the legacy DSM-IV-TR criteria, loss had to occur in two or more domains. The domains were expressive or receptive language; social skills or adaptive behaviour; bowel or bladder control; play; and motor skills. This breadth of loss is clinically important. A history that includes toileting and motor loss cannot be reduced to a narrow social-communication account. It should trigger medical evaluation alongside developmental and psychiatric assessment.

- Ask what the child could actually do before decline, using concrete examples rather than global parental labels.
- Establish whether language, social or adaptive abilities, continence, play and motor skills were all affected, and in what sequence.
- Seek neurological and systemic symptoms alongside the behavioural change, then pursue the regression differential described in the differential diagnosis of ASD section.

Following the regression, **autistic symptomatology** emerges, with impairments in socialisation and communication together with restricted, repetitive patterns of behaviour. The clinical

features are otherwise quite similar to autistic disorder, and age at onset is the principal distinguishing feature. This is why a prolonged normal developmental period makes the distinction easier, while a shorter apparently normal period makes it less secure.

27.6 Rett syndrome and childhood disintegrative disorder compared

CLINICAL AXIS	RETT SYNDROME	CHILDHOOD DISINTEGRATIVE DISORDER
Core framing	MECP2-related neurological syndrome	Historical syndrome of regression after apparently normal development
Typical developmental timing	Apparently normal early period of 6-18 months	Regression most typically between 2 and 4 years
High-yield clues	Loss of purposeful hand use with hand stereotypies; acquired microcephaly and motor signs	Loss across language, social or adaptive, continence, play and motor domains
Current DSM position	Not an ASD subtype	Subsumed within ASD
Expected outcome	Long-term neurological disability with changing motor and medical needs	Generally more severe intellectual disability and poorer outcomes than autism

The table should be used as a reasoning aid, not as a substitute for a history. Rett syndrome has a biologically distinctive phenotype, while childhood disintegrative disorder is defined by its temporal pattern of broad regression. In both cases, apparent autism is a clinical presentation that must be interpreted in context. The most hazardous mistake is diagnostic closure after observing social withdrawal or repetitive behaviour.

27.7 Current status, intervention and outcome in childhood disintegrative disorder

Childhood disintegrative disorder is not a separate diagnosis in DSM-5 or DSM-5-TR; it is encompassed within ASD and referred to as a disorder previously described as childhood disintegrative disorder. There is no CDD category in ICD-11 CDDR. Its legacy codes are F84.3 in ICD-10 and 299.10 in DSM-IV or DSM-IV-TR. These codes are useful for interpreting older records, but they should not be presented as current parallel diagnostic categories.

DSM-5 absorption should not make the regression clinically bland. A child with the pattern formerly called CDD requires extensive medical investigation, because regression warrants a search for neurological and medical explanations. The detailed regression differential belongs in the differential diagnosis of ASD section, and the broader ASD medical workup is addressed in the medical comorbidities and workup section. The psychiatrist's immediate contribution is to preserve the chronology, document the breadth of lost skills and ensure that an autistic formulation does not truncate investigation.

Intervention methods for CDD are identical to those used for autism. This statement does not mean the clinical task is identical: marked functional loss often changes the family's practical needs and the level of support required. The named behavioural programmes, speech and

occupational interventions, and family support are dealt with in their respective sections of this chapter. Here, the key point is that a historical CDD label does not exclude access to ASD-focused intervention.

Outcome is generally more severe than in autism. CDD has been described as having a much worse prognosis, with more severe intellectual disability and poorer outcomes than autism. This is why it is important to be honest with families without becoming deterministic: outcome language should describe present support needs, continuing medical assessment and the need to revise the formulation if the course changes. It should never be used to imply that rehabilitation, communication support or family partnership are futile.

27.8 Practical clinical synthesis

GOOD TO REMEMBER

When a child with possible ASD has lost skills, establish the developmental baseline and order of loss. Ask about purposeful hand use, continence, play and motor ability; then examine for acquired microcephaly, stereotypies, gait or trunk abnormalities, breathing changes and other neurological signs. Decide whether this is ASD with loss of skills, a known syndrome such as Rett syndrome, or a regressive disorder requiring a broader medical differential.

- For Rett syndrome, preserve the distinction between a clinical diagnosis and a supportive molecular result.
- For any psychiatric prescription in Rett syndrome, incorporate the arrhythmia and QT-risk context into the medical review.
- For historical CDD, document the prolonged normal period and the domains of loss before applying a legacy label; in either presentation, communicate that regression deserves explanation, not merely classification.

These two conditions are therefore important not because they add obscure eponyms to ASD, but because they teach a general clinical discipline. Developmental regression is a temporal event with neurological and functional meaning. Recognising its pattern protects the child from premature diagnostic closure and gives the family a more coherent account of why assessment, treatment and support must extend beyond a behavioural description.

28 · Behavioural interventions (ABA, TEACCH, PECS, ESDM)

Behavioural intervention is a clinical method, not a branded package. In autism spectrum disorder, the psychiatrist's task is to translate a broad concern - communication, participation, learning, distress around change, or a behaviour that has become difficult to manage - into an observable target and a coherent programme. The useful question is not which acronym is fashionable. It is which learning opportunity, environmental support, communication route and review process are most likely to help this person participate more effectively in ordinary life.

Behavioural intervention means planned alteration of the conditions in which a skill is taught or a behaviour occurs, with observation of whether the planned change helps. It is not

synonymous with making a child compliant, and it must never be framed as a contest of wills. Good work begins with strengths, interests, communication needs and the burden placed on the person and family. It then turns a vague wish such as “behave better” into a target that can be observed, practised and reviewed. This fragment concerns named behavioural and educational programmes; the service model of early intervention belongs in the management of intellectual disability, while speech and occupational therapy and family support are addressed separately.


The autism intervention map: the named programmes by target, by age, and by what the evidence will actually license

Four things are mapped: the programme, what it targets, the age its source gives it, and the rating the ledger permits. The fourth is the hard one.

The IPS CPG's Table 7 is mangled in the parse this chapter was built from: its rows and its evidence column desynchronise. Four of its ratings survived corroboration. Three of those four name a behavioural programme, and those three are the only IPS ratings drawn here: TEACCH Low, PECS Moderate, ESDM Moderate or insufficient. No rating is drawn for ABA-based EIBI itself, for SCERTS, for Floortime or DIR, or for parent-delivered home ESDM: those bindings are positional, held at partial, and an exam artefact should not spend its error budget on them. Not sourced, and therefore not drawn: the Lovaas 40 h per week figure; a month-level age band for ESDM; the phase structure of PECS; an evidence grade for speech and language therapy or for occupational therapy; an effect size, sample size or follow-up for PACT; the second NICE-recommended intervention; and TEACCH framework component 2, lost to the same parse.


A. Mapped by age. Two age numbers exist in this chapter's intervention sources, and these are both of them.

A bar is drawn only where a source states a band. Where a row says "no age band", that is the finding, not a gap in the drawing.



B. Mapped by target. What the programme is trying to change, and the only strength the ledger will license for it.

"Drawn" means the rating survived corroboration and appears on this plate. "Withheld" means the ledger holds it at partial, so it is named and not drawn.



How much: three sources, three doses, and the ledger adjudicates none of them
IPS CPG, ABA narrative: ideally more than 20 h per week, for a child under the age of 4.
Tasman: every intensive early intervention programme should encompass a minimum of 20 to 25 h per week.
IPS CPG Table 7, the ABA-based row: intensive teaching for 20 to 40 h per week, for 1 to 4 years, at a 1:1 adult-to-child ratio.
The three agree only at their floor of 20 and diverge above it. Quote the source with the number. The bare "ABA is 40 hours a week" belongs to none of them.

Two traps, both sourced, both examinable
Trap 1. Two rating scales, not one. IPS Table 7 rates evidence for effectiveness (Low, Moderate, Not established). Rutter's Table 36.1 tabulates the Rogers and Vismara 2008 tiers (Well-established, Probably efficacious). TEACCH's Low and the Lovaas model's Well-established are not two points on one scale and must never be ranked against each other. In that review, exactly one autism treatment was designated well-established, and it is the Lovaas model.
Trap 2. The IPS CPG misprints ESDM as "Early Stage Denver Model" in all three of its occurrences. The correct expansion is "Early Start Denver Model" (Dawson and Rogers, New Oxford). Know the guideline's error and do not cite the Indian guideline as the authority for the expansion.

What the Indian guideline actually recommends, from a table the parse hazard does not touch
IPS CPG Table 10, its overview of recommendations, is structurally independent of Table 7, which makes it the safest IPS binding in this chapter's ledger. Its nonpharmacological column holds exactly three entries: ABA, PECS and social stories. A programme can be expounded at length by this guideline and still not appear in that column. Described is not recommended, and this plate keeps the two apart.

Legend:
An age band the source states in years. The bar's ends are the sourced numbers, and they are the only two on the plate.
A band the source states in words, not years. The bar spans the axis because the cell names life stages; its ends are not boundaries.
A sourced number: an intensity, a duration, a ratio, an age bound. Every one of them traces to a single claim row.
A sourced warning: a parse hazard, a guideline misprint, or a scale that must not be ranked across.

Every value here is a Day-21 claim row. Age: ASD-MGMT-20 (under 5 y, the ABA-based row), ASD-MGMT-02 (under 4 y, the ABA narrative), ASD-MGMT-14 (TEACCH), ASD-MGMT-17 (PECS, a population and not an age). Intensity: ASD-MGMT-01, 03, 04, 05, 06. Ratings drawn: ASD-MGMT-13 (TEACCH), 16 (PECS), 19 (ESDM), 08 (Lovaas EIBI), 10 (PRT), 09 (exactly one well-established). Counts: 53-10 (six modalities, the latter three), ASD-MGMT-24 (the NICE two, held at partial because the source never names the second). Names and expansions: ASD-MGMT-07, 11, 15, 18, 23, 25. Table 10: claim 53-05. Rutter's counter-readings: claims 53-08 and 53-14.
The parse hazard is declared by the ledger itself, not inferred here: IPS Table 7 was split into two half-tables by the parser and its rating column desynchronises down the page.
No pharmacological agent, dose or target symptom appears on this plate and none is missing: the pharmacotherapy note owns all of it, and this plate stops at the programmes.

Fig 21.20 The autism intervention map. The named behavioural programmes placed by age and by target, carrying only the evidence ratings that survived corroboration: the IPS guideline's Table 7 is mangled in the parse behind this chapter, so TEACCH Low, PECS Moderate and ESDM Moderate-or-insufficient are drawn while the ratings for ABA-based EIBI itself, SCERTS, Floortime and parent-delivered home ESDM are named and withheld.

28.1 Clinical logic before programme selection

A programme label does not replace formulation. The clinician should first establish what is happening, where it happens, what tends to precede it, what follows it, and what the person may be communicating through it. The same outward behaviour can call for quite different changes. This is why copying a technique from another child's plan is poor practice.

Functional formulation links a target behaviour to its context and probable purpose. It makes the intervention humane and testable. Behavioural reasoning should reduce avoidable distress, not explain it away.

GOOD TO REMEMBER

When a child leaves a task, ask what the behaviour may be communicating before choosing a response. It may reflect an incomprehensible demand, a need for a break, an attempt to obtain an item, sensory overload or distress needing medical or psychiatric assessment. The plan may need a communication response, a better-matched task, clearer cues, a smaller step or a predictable way to ask for help.

■ RULE Treat the function of a behaviour and the person's communication need, not merely its visible form.

When the target and its context are explicit, family members, teachers and therapists can work towards the same practical outcome. When they are not explicit, adults often respond inconsistently and the plan becomes a series of reactions rather than treatment.

Targets should be meaningful to the person's daily life. They may concern requesting, tolerating a transition, joining a routine, sharing attention, completing a manageable activity, or reducing a barrier to participation. Progress must be interpreted in context: a newly learned response that occurs only in a training room has not yet become a dependable everyday skill. The clinician should ask where a skill is expected to occur, who can prompt it, what support makes it possible, and how the support will be reduced or adapted when appropriate. This keeps the programme connected to participation rather than to worksheet performance.

- State the target in observable language and identify why it matters in daily life.
- Record the setting, likely antecedents, consequences and communication context.
- Choose a teaching route that matches the person's current access to communication and learning.
- Review benefit, burden, generalisation and acceptability with the family and educational team.

28.2 Applied behavioural analysis and intensive behavioural teaching

Applied behavioural analysis is a framework for using behavioural principles to define targets, arrange teaching opportunities, record response and alter the plan in the light of observed learning. Its strength is not a rigid script. It is disciplined attention to what is being taught, what cues are available, what response is expected, what consequence makes a response worth repeating, and whether the skill transfers outside the teaching situation. A plan can be systematic without being mechanical, and it can be individualised without becoming unmeasurable.

Teaching may be organised into brief, clearly cued learning opportunities or embedded in meaningful activity and routine. The clinical distinction is useful: a more structured format can help when a new discrimination needs clarity, whereas opportunities in ordinary activity are important when a response needs to become useful beyond a formal session. In either case, prompting should help the person succeed rather than repeatedly expose failure. The team should watch for prompt dependence, because an apparently learned response may actually be a response to an adult’s familiar cue.

Early intensive behavioural intervention is an ABA-based programme organised around a broad developmental teaching plan rather than an isolated target. The model is associated with **Ivar Lovaas**. A historical evidence review reported that only one autism treatment met its well-established category, namely that EIBI model, classified as well-established. That classification is useful for understanding the evidence history; it should not be turned into a promise about any individual child or into a reason to disregard individual goals and feasibility.

Intensity is often discussed as though it alone determines quality. It does not. The relevant question is whether there is enough repeated, purposeful teaching for a child to learn and use skills, while preserving play, rest, family life, education and responsiveness to distress.

IN INDIA

The IPS guidance recommends, ideally, more than **20** hours per week of ABA for children below 4 years. A broader intensive-programming statement sets a minimum of **20-25** hours per week. These are programme-intensity statements, not a prescription to fill every waking hour with adult-led drills.

20-40 ABA-BASED EIBI TEACHING, WEEKLY HOURS	1-4 ABA-BASED EIBI PROGRAMME, YEARS	1:1 ADULT-TO-CHILD RATIO STATED FOR THE EIBI ROW	<5 USUAL TARGET AGE IN THE EARLY-INTERVENTION ROW, YEARS
---	--	---	---

More specifically, the ABA-based EIBI row describes teaching between **20-40** hours per week over between **1-4** years, with an exactly 1:1 adult-to-child ratio for that EIBI row. The same early-intervention row is directed at young children, usually below 5 years. These details describe an intensive programme architecture. They do not mean that every component, or every teaching moment, must be delivered in an identical format.

The psychiatrist should make the architecture clinically workable. Ask who delivers each part, how parents understand the aim, whether the educational setting can reinforce the same communication route, and what data will trigger a change in plan. A programme can be intensive on paper but ineffective if targets are poorly chosen, staff turnover is high, the child is distressed, or skills never leave the treatment room. Conversely, a lower-resource setting can still use a disciplined behavioural formulation when it selects priorities carefully and trains the adults who share the person’s routine.

-
- **TRAP Examination trap.** Do not equate ABA with a single teaching technique, or assume that a stated adult-to-child ratio applies only to one teaching format. The cited ratio belongs to the ABA-based EIBI row as a whole.
-

28.3 Structured TEACCHing

TEACCH expands to **Treatment and Education of Autistic and related Communication-handicapped Children**. Its clinical contribution is the deliberate use of structure so that expectations are legible. Rather than assuming that verbal explanation will organise behaviour, the environment, task presentation and sequence are made easier to interpret. This can reduce ambiguity for a person who finds open-ended social or practical demands difficult to process.

Structured TEACCHing is best understood as a way of making the environment do some of the organising work. The framework is described as having exactly four components. In clinical use, the central question is whether the arrangement helps the person see what is happening now, what is expected next, how much is involved, and when an activity is complete. A visual or physical cue is not therapeutic merely because it has been produced; it is useful only if it reduces uncertainty and supports a meaningful response.

Visual structure can be incorporated into home, school or adult settings without reducing the person to a timetable. It may clarify transitions, make materials easier to locate, distinguish work from leisure, or show the endpoint of an activity. The clinician should avoid turning structure into inflexibility. If a plan is imposed without attending to preference and communication, it can become another demand. The better aim is predictable access to activity and choice, with room for the person to signal refusal, need for help, or a wish to change course.

The evidence-for-effectiveness rating listed for TEACCH is low. This should prompt honest discussion rather than dismissal. A low rating does not prove that a structured environment is useless for an individual; it says that confidence in an effect from the cited evidence base is limited. TEACCH is listed for children, adolescents, and adults, which is clinically important because environmental clarity can remain relevant across settings and stages of life. The decision should be based on a specific participation problem, a clearly described support, and observable review.

28.4 PECS and functional communication

PECS means **Picture Exchange Communication System**. It is a communication intervention in which pictures or symbols can provide an accessible route for an individual to initiate an exchange. Its relevance is not that pictures are inherently superior to speech. Its relevance is that a person who cannot reliably use speech may still have a clear reason to communicate, and should have a practical means of doing so.

Functional communication is the clinical aim: a response that changes the person's access to something meaningful in a socially understandable way. When an individual can request, refuse, seek help or indicate a preference, adults need not infer every need from distress or behaviour. This is why communication planning belongs in behavioural formulation. It may reduce conflict

by replacing an ineffective route with a more reliable one, but it should never treat communication as a device for suppressing autonomy.

PECS is listed for nonverbal individuals, with a moderate evidence-for-effectiveness rating. In assessment, establish whether the person notices the visual representation, can physically manage the exchange, understands what will follow, and has communication partners who will respond consistently. The picture, book or display is not the intervention by itself. The intervention is the repeated, successful experience that an intentional act reaches another person and has a meaningful consequence.

Clinical review should include whether the communication route is available where it is needed, not just during planned teaching. A system that stays in a cupboard or depends on one trained adult can produce a misleading appearance of non-response. Equally, the clinician should not make a symbolic system a compulsory endpoint. The right question is whether it expands the person’s agency and participation. The separate discussion of speech and language therapy addresses broader communication assessment and therapy; the present focus is the behavioural logic of an accessible communicative exchange.

MNEMONIC

MAP the communication plan

Meaningful need, Accessible response, Partner who responds. If any part is missing, the plan risks becoming a display rather than communication.

28.5 ESDM within early behavioural programmes

ESDM is the **Early Start Denver Model**. In a clinical plan it should be named accurately and then linked to a specified developmental and behavioural target, rather than invoked as a generic synonym for good early care. It appears in the same ABA-based early-intervention row as EIBI and other named programmes. That placement helps the candidate remember that programme labels are not competing diagnoses: they are organised approaches whose usefulness depends on the individual’s needs, the quality of delivery and the fit with family and educational contexts.

The evidence rating listed for ESDM is **moderate or insufficient**. This wording requires restraint. It should not be simplified into a claim of established benefit for every outcome, nor used to imply absence of any possible benefit. When discussing it with a family, be precise about the proposed targets, the opportunity cost, the support required of caregivers and the plan for reviewing whether the programme is helping. In a fragmented service system, a recognisable model may be less important than a coherent, respectful plan that can actually be delivered and monitored.

The early-intervention row is directed to young children usually below 5 years. This is a description of the target age for that programme row, not a rule that support ceases after that threshold. Across the lifespan, the psychiatrist still identifies barriers to participation and coordinates a plan appropriate to the person’s current setting. The form of teaching, the

communication route and the environmental adaptation may change, but the principle of setting meaningful goals and reviewing their effect remains constant.

28.6 Comparing programmes without false equivalence

Named programmes overlap in their concern with learning and participation, but they do not answer precisely the same clinical question. The accompanying table is a way to hold the distinction in mind. It is not a menu from which a family should be asked to choose unaided. The psychiatrist's role is to identify the main barrier, explain what a proposed approach is intended to change, and ensure that claims about evidence are communicated with their stated limits.

APPROACH	CLINICAL ORGANISING IDEA	EVIDENCE OR SCOPE STATED IN THE SOURCE	QUESTION FOR REVIEW
ABA-based EIBI	Planned behavioural teaching across meaningful targets	Between 20-40 hours per week over between 1-4 years	Are skills becoming useful outside teaching?
TEACCH	Environmental and task structure that makes expectations clearer	Low evidence rating; listed for children, adolescents, and adults	Does the support reduce ambiguity and improve participation?
PECS	Accessible symbolic exchange for communication	Moderate evidence rating; listed for nonverbal individuals	Can the person initiate a meaningful exchange in daily settings?
ESDM	Named early behavioural programme linked to individual goals	Moderate or insufficient evidence rating	Are proposed goals, burden and observed benefit explicit?

The comparison also guards against a familiar clinical error: treating an evidence rating as a property of the child. It is not. Ratings describe a body of evidence for an intervention as studied and categorised. The individual question remains whether a carefully specified, feasible intervention is improving a valued outcome. A programme with a stronger evidence classification may still be poorly delivered or poorly matched; a supportive adaptation with limited evidence may be reasonable when it has a clear rationale, low burden and observable benefit. Neither statement licenses therapeutic drift. Every plan needs a target, a responsible team, a way to observe change and a point at which it will be reconsidered.

28.7 Monitoring, ethics and the psychiatrist's coordinating role

Treatment fidelity means that the agreed intervention is actually being delivered in a recognisable and consistent way. It is an ethical as well as a methodological issue. Families are often asked to commit time, effort and money. If the target is vague, adults use incompatible prompts, or records are collected without changing the plan, the family carries the burden while nobody can say what is being tested. Clear documentation protects both the person and the team.

Monitoring should be practical. It can include the frequency or context of a target response, the level of assistance needed, the settings in which it occurs, the person's affect and signs of distress, and the views of family and staff. The record is not an end in itself. It should answer whether the

intervention is helping, whether the goal remains meaningful, and whether a different explanation of the difficulty is now more plausible. A plateau is not a moral failure by the child or caregiver. It is a prompt to re-examine the formulation, access to learning opportunities and competing sources of distress.


The psychiatrist also has a boundary-setting role. Behavioural work must preserve dignity, assent where it can be expressed, choice, and access to communication. It should not make the suppression of harmless difference its central aim. Concerns about pain, sleep, seizures, anxiety, mood change, trauma, bullying, family stress or adverse effects of medication require appropriate assessment rather than escalation of a behavioural plan. The wider assessment and care system is covered in the Child psychiatry: assessment, treatment, services and protection notes.

What the candidate should be able to defend. A sound answer does more than expand acronyms. It explains why a target was chosen, how the proposed method fits the observed barrier, what the evidence statement does and does not justify, and how progress will be judged in ordinary settings. The best intervention plan is not the most elaborate one. It is the one that gives the autistic person a more understandable environment, a more usable route to communication and participation, and a team capable of learning from the result.

Remember: Select ABA, TEACCH, PECS or ESDM by the clinical problem and delivery context, then judge the programme by meaningful everyday change rather than by its acronym.

29 · Pharmacotherapy in ASD (target symptoms, risperidone, aripiprazole)

Medication in autism is symptom-directed. It is not a treatment for the autistic condition as a whole. No medicine is established to alleviate the **core features of autism spectrum disorder**. This distinction is the conceptual anchor for a safe prescription: the clinician is treating a disabling, observable target and its consequences, not attempting to medicate autistic identity or to substitute a drug for developmental, environmental, or family work.



- Define the target
- Measure baseline
- Review benefit and harm
- Check context
- Time-limited trial
- Never for autism itself

Fig 21.21 Medication in autism is directed at a severe, observable target behaviour and measurable impairment, not at autism itself. Context is assessed first, outcomes are defined before prescribing, and benefit and harm are reviewed as a time-limited clinical trial.

The marks in this area come from clinical precision. A good answer begins by naming the target behaviour, judging its severity and impact, and stating why medication is justified now. It then selects a medicine whose evidence fits that target, starts conservatively, monitors benefit and harm with the same discipline, and plans review from the outset. The prescription is therefore a time-limited clinical experiment with explicit outcomes, rather than an automatic response to a diagnosis.

29.1 The target-symptom framework

Irritability in autism is a clinical cluster, not merely a description of a child who is difficult to manage. In the regulatory construct for risperidone, it includes aggression towards others, deliberate self-injury, tantrums, and rapidly changing moods. These behaviours may cause injury, exclude the person from ordinary settings, exhaust carers, or make education and communication

impossible. They deserve active treatment, but the treatment target must be stated in behavioural language that can later be reviewed.

-
- **RULE** **Prescribe for a defined target behaviour and measurable impairment, never for autism itself.** A medication trial is defensible when the target is severe, persistent, and associated with meaningful risk or loss of function, after the immediate context has been understood. It is not defensible simply because a person is autistic, has repetitive behaviour, or causes concern to others.
-

Before calling a presentation psychiatric, reconstruct its function. Consider what precedes the behaviour, what follows it, whether pain, fear, sensory overload, communication failure, a demand that exceeds capacity, or a change in routine is maintaining it, and whether the behaviour has become a reliable route out of an intolerable situation. This is not an argument against medication. It is what prevents a medicine from being used to silence the only available signal of distress. The wider assessment of co-occurring medical and psychiatric conditions belongs with the medical-comorbidity discussion; here the practical point is that an unexplained new target should be assessed before it is medicated.

Agree in advance what improvement would look like: fewer episodes, less intensity, less injury, better participation, or greater ability to use supports. Also agree what would make the trial unsuccessful: no clinically meaningful change, unacceptable adverse effects, or a target that turns out to have been incorrectly formulated. Carer observations are valuable, but must be joined to direct clinical enquiry and the person's own communication wherever possible.

-
- **TRAP** **Examination trap.** Do not write that antipsychotics treat the social-communication differences or restricted interests of autism. Their established role in this section is management of severe irritability and associated dangerous behaviour.
-

29.2 Deciding whether medicine is needed

A target-symptom formulation protects against diagnostic overshadowing. The question is not whether the behaviour appears dramatic; it is whether there is a sufficiently clear syndrome of irritability, a serious consequence, and a realistic opportunity to judge change. It also protects the person from prolonged exposure to a drug when its intended benefit was never specified.

- Describe the target in observable terms and identify the harm, distress, or loss of access that makes it clinically important.
- Seek a working explanation for the behaviour, including communication and environmental contributors, before interpreting it as a medication-responsive syndrome.
- Record a baseline and choose a small set of outcomes that matter to the person and carer.
- Discuss the expected trade-off openly: reduction of dangerous behaviour may be valuable, but sedation, appetite change, movement effects, endocrine effects, and weight change may alter quality of life.

This approach also clarifies the place of non-drug intervention. Environmental adaptation, communication support, behavioural work, and carer support remain necessary even when

medication is appropriate. A drug may lower the intensity of a crisis enough for those measures to become usable; it does not replace them. The detailed programmes and therapies are addressed elsewhere in this chapter.

Shared decision-making is particularly important where the observable benefit is reported mainly by adults around the child. Ask what has changed for the person: comfort, participation, sleepiness, appetite, mobility, alertness, and ability to engage. A quiet child is not necessarily a better child. The therapeutic objective is improved safety and access to life, not convenience for a system that has failed to adapt.

29.3 Evidence base and place of risperidone

Risperidone and aripiprazole are the **two FDA-approved atypical antipsychotics** for irritability in children and adolescents with autism. For risperidone, the approved wording is irritability associated with autistic disorder. The listed age range is **5-16 years**. These regulatory facts should not be converted into an indiscriminate rule that every distressed autistic young person requires an antipsychotic. They identify a well-studied target, not a substitute for formulation.

Risperidone has the clearest short-term evidence when severe irritability is the dominant target. In a controlled trial, **69%** of participants receiving risperidone showed substantial reduction in irritability, compared with 12% receiving placebo. Across additional trials, short-term response generally lay between 57% and 72%. The clinical reading is not that response is guaranteed. It is that a clearly observed trial can be worthwhile where the presenting problem is dangerous, severe, and recognisably within this target cluster.

69% SUBSTANTIAL IRRITABILITY REDUCTION WITH RISPERIDONE	12% CORRESPONDING PLACEBO RESPONSE	2.7 kg MEAN WEIGHT GAIN IN SHORT-TERM LARGE TRIALS	5.6 kg MEAN WEIGHT GAIN AT FOLLOW-UP IN THE CITED STUDY
---	---	--	---

Evidence must be interpreted alongside burden. In the discontinuation phase of a risperidone study, 62.5% relapsed when switched to placebo. This makes abrupt, reflexive withdrawal after a response unwise. It does not mean that treatment must continue indefinitely. It means that the earlier formulation and the conditions that supported improvement should be revisited before tapering, and that the family needs a plan for recognising recurrence.

A meta-analysis of 46 randomised controlled trials found risperidone and aripiprazole the most effective agents for irritability, with moderate to large effects relative to placebo. This supports their priority over speculative polypharmacy. It also explains why an answer that simply lists psychotropics loses the clinical point: the evidence concerns a particular target, not a general drug treatment of autism.

29.4 Risperidone: practical dosing and titration

Start low and make each change interpretable. Weight-sensitive paediatric dosing is designed to reduce avoidable adverse effects while allowing time to judge the direction of change. The smaller-weight schedule begins at **0.25 mg/day** and then moves to 0.5 mg/day. Subsequent

increases are 0.25 mg at intervals of 2 weeks or more, within a licensed range of 0.5-3 mg/day whose upper end is not a target for a small child: therapeutic benefit plateaus at **1 mg/day**, and only patients over **45 kg** may require more. Read the range as the outer bound of the licence and the plateau as the clinical ceiling. That plateau applies to both weight schedules rather than to the lighter one alone; escalation beyond a useful response should therefore not be routine.

For the higher-weight schedule, the initial dose is 0.5 mg/day, followed by 1.0 mg/day. Further increases are 0.5 mg at intervals of 2 weeks or more, with a total range of 1.0-3 mg/day. The 1 mg/day plateau applies to this schedule too. Only those weighing more than **45 kg** may require higher doses, and it is in that subgroup alone that the effect plateaus at 3 mg/day. Reading the higher plateau as licence for every child on this schedule is the error the weight bands exist to prevent. Below **15 kg** there is no paediatric dosing data at all, and the source's instruction is caution for that reason rather than caution about data that exist. The distinction is the whole point: an absent evidence base is not a cautious one.

CLINICAL SITUATION	STARTING AND EARLY DOSE	LATER ADJUSTMENT	INTERPRETIVE POINT
Lower body-weight schedule	0.25 mg/day, then 0.5 mg/day	Increase by 0.25 mg at intervals of 2 weeks or more	Licence 0.5-3 mg/day, but benefit plateaus at 1 mg/day: the top of the range is not a target at this weight
Higher body-weight schedule	0.5 mg/day, then 1.0 mg/day	Increase by 0.5 mg at intervals of 2 weeks or more	Licence 1.0-3 mg/day, but benefit plateaus at 1 mg/day: the 3 mg/day ceiling belongs to patients over 45 kg alone, so a lighter child on this schedule is not titrated toward it

GOOD TO REMEMBER

The table is a dosing aid, not permission to chase the top of a range. At each step, ask whether the pre-agreed target has changed enough to justify further exposure. A partial response may be clinically important if it prevents injury or restores access to school and family life. Conversely, an apparently calmer presentation accompanied by marked sedation may represent harm rather than success. Avoid simultaneous medication changes where possible, obtain information from the settings in which the target occurs, and resist adjunctive psychotropics before the initial trial has become interpretable.

29.5 Risperidone: adverse effects, monitoring, and review

Weight gain is the adverse effect that most often changes the risk-benefit balance over time. In the larger short-term trials, mean gain was 2.7 kg. In a follow-up study it was 5.6 kg, an increase of 16.7% from baseline body weight. A paediatric estimate over a year was 7.5 kg, compared with expected annual gain of 3-3.5 kg/year. These figures make routine measurement clinically necessary rather than an administrative formality.

Prolactin elevation is another important adverse effect. Reported rates in paediatric risperidone users range from 49% to 87%, compared with 2% to 7% on placebo. Galactorrhoea has been reported in 0.8%, and gynaecomastia in 2.3%, of paediatric users. These are not simply laboratory concerns: enquire sensitively about symptoms, pubertal concerns, and distress, and interpret any abnormality in the whole clinical context.

Movement effects require active questioning and observation, especially when altered posture, restlessness, or behavioural change might otherwise be attributed to autism. Tardive dyskinesia was reported in 0.1% of paediatric participants in industry-sponsored risperidone trials, while an annual estimate for young people receiving antipsychotics was 0.3% per year. Transient somnolence after initiation had a median duration of 16 days. The numbers are low enough to tempt omission and important enough to make that unsafe: movement phenomena and sedation may be difficult to elicit from a person with limited speech.

Monitoring must therefore connect directly to the indication. At every review, revisit the original behaviour, weight trajectory, appetite, alertness, movement symptoms, endocrine symptoms, and the person's capacity to participate. If benefit is marginal, adverse effects should carry more weight in the decision. If benefit is substantial, monitoring is still not optional, because an initially acceptable trade-off can become unacceptable as exposure continues.

29.6 Aripiprazole: alternative evidence and dosing

Aripiprazole is the other approved antipsychotic for autism-related irritability in children aged 6-17 years; the FDA approval was granted in 2009. It is an alternative when the same target syndrome is present and risperidone is unsuitable, not a drug for core autism. Its initial dose is 2 mg/day. The usual range is 5-15 mg/day, the maximum is 15 mg/day, and titration proceeds by 5 mg/day at intervals of no less than 1 week.

Across the pivotal trials, active-arm responder rates were 52.2% and 55.8%, while placebo rates were 34.7% and 14.2%. The trial-to-value pairing is not reliable and should not be asserted. The defensible conclusion is that the medicine has evidence of benefit for irritability, but the size of individual benefit and the placebo contribution need careful reading rather than slogan-level certainty.

At short-term follow-up, aripiprazole-associated weight gain was 1.3-2 kg, somewhat lower than that reported with risperidone in the cited trials. It was not associated with prolactin elevation. These differences may matter when choosing between agents, especially where weight or endocrine effects have already shaped the discussion. They do not make aripiprazole benign: treatment-emergent extrapyramidal symptoms and vomiting were reported more often than in the risperidone studies. Moreover, aripiprazole may be ineffective for self-injurious behaviours. The target behaviour, rather than an imagined universally preferable drug, should drive selection.

The comparative choice is therefore a conversation about the individual target and anticipated trade-offs. Risperidone offers a strong evidence base with substantial concerns about weight and prolactin. Aripiprazole offers a distinct adverse-effect pattern, including movement effects and vomiting, and may have limitations where self-injury is the central target. Both require the same

disciplined baseline, follow-up, and willingness to stop if the promised benefit does not materialise.

29.7 Restricted and repetitive behaviour and the limits of extrapolation

Restricted and repetitive behaviours and interests are often distressing, but they should not be treated as a single undifferentiated drug target. A Cochrane review found no evidence of effect and emerging evidence of harm for selective serotonin reuptake inhibitors in children with autism. This is an important corrective to the tempting, but unsound, analogy that repetitive behaviour in autism should automatically be treated as though it were an obsessive-compulsive symptom.

The negative citalopram trial illustrates the point: responder rates were 33% with citalopram versus 34% with placebo. Evidence should be allowed to overrule pharmacological intuition. If an SSRI is being considered for another clearly formulated indication, that decision requires its own assessment; it should not be presented as treatment of core repetitive behaviour in a child with autism.

Where fluoxetine is used in the paediatric regimen cited for repetitive behaviours, the starting schedule is 2.5 mg/day for 1 week and the maximum is 0.8 mg/kg/day. These dosing details do not overturn the negative child evidence. In an examination answer, that distinction earns more credit than listing a drug without stating the weakness of its indication.

29.8 Duration, discontinuation, and the longitudinal plan

Planned discontinuation belongs in the initial prescription, not only after adverse effects emerge. For irritability in autism, antipsychotic treatment appears beneficial for **6-12 months**, after which discontinuation should be considered. This is a prompt to review, not a rigid deadline. The clinician should ask whether the original syndrome persists, whether the environment and communication supports have changed, whether adverse effects have accumulated, and whether a carefully observed reduction is now feasible.

The success of medication in autism is not a quieter person. It is a safer, less distressed person with greater access to ordinary life, at an acceptable cost in adverse effects.

When a person has improved, explain that withdrawal is also a clinical test. A gradual, monitored reduction allows the team to distinguish continuing need from inertia. If the target returns, reconsider the formulation rather than simply restoring medication without thought: relapse may identify a genuine medication-responsive syndrome, an environmental trigger, loss of support, or an insufficiently prepared taper. If the target does not return, the successful outcome is not merely drug cessation but a person spared unnecessary exposure.

For the MD or DNB candidate, the mature answer is consequently structured rather than encyclopaedic: state that core autism is not medicated, define the target, justify either risperidone or aripiprazole for severe irritability, give cautious dosing, name the adverse-effect trade-off, and

make review and discontinuation part of treatment. That sequence shows pharmacological knowledge and, more importantly, clinical judgement.

30 · Speech and occupational therapy and family support in ASD

The practical aim is participation, not performance. Speech and language work, occupational therapy and family support translate an autism formulation into a plan that can be used during ordinary communication, play, self-care, learning and community life. They are not interchangeable therapies, and they should not be presented to families as a menu of equally established options. The useful clinical question is: which barrier is limiting this person's participation now, and what change can the child, family and communication partners realistically carry across settings?

A good plan makes communication more available, adapts demands and environments, and gives carers a role that is sustainable. It also makes uncertainty explicit. This matters especially where an intervention has an appealing explanatory story or a memorable label: plausibility, popularity and individual preference do not by themselves establish effectiveness. The psychiatrist's contribution is to hold the formulation together, protect the family from exaggerated promises, and ensure that therapy goals remain linked to daily function rather than to a fashionable technique.

30.1 Speech and language therapy: from language form to functional communication

Speech and language therapy begins with an account of how the person currently communicates, what communication partners understand, and where breakdown occurs. The assessment is broader than spoken output. It asks what the person can initiate, how they show choice or refusal, whether they can repair a misunderstanding, and which routines make communication easier or harder. The resulting targets should be concrete enough for a caregiver and teacher to recognise in ordinary life. A target that cannot be seen outside the therapy room is difficult to generalise and difficult to judge honestly.

■ **RULE** **Treat communication as access to agency.** The person needs a reliable way to request, refuse, share attention and participate, whether or not speech is the route.

This framing prevents a common drift towards making speech itself the sole outcome. Spoken language may be an important goal for some people, but clinical work should not postpone access to communication while waiting for it. The therapist therefore works with the person's existing signals and with the adults who respond to them. The psychiatrist should ask whether the proposed therapy has identified a meaningful communication function, a communication partner, and a setting in which the new skill will be practised.

A **communication partner** is not merely a helper standing nearby. The partner notices an attempt, allows time for a response, interprets it cautiously, and creates repeated opportunities for the person to communicate. This is why family and classroom work cannot be an afterthought to individual sessions. A technically correct target can fail if the adults around the person repeatedly anticipate needs, change the routine without warning, or respond inconsistently.

Conversely, a modest target can become powerful when it is recognised in the home, school and clinic.

Clinical review should distinguish a communication goal from a behavioural label. For example, a difficult moment may represent an unmet need, an inaccessible instruction, an intolerable demand, or an ineffective means of protest. The response is not to assume intent from the behaviour alone, but to test whether a clearer route for communication and a changed interaction pattern reduce the impasse. This keeps speech and language work tied to formulation without turning every difficulty into a speech problem.

- Specify the everyday situations in which communication fails.
- Agree on the response expected from communication partners.
- Choose a visible marker of change that families can recognise without specialist scoring.
- Review whether the skill appears outside the session, not only during prompted practice.

30.2 Augmentative and alternative communication in the therapy plan

Augmentative and alternative communication covers every route that supplements or replaces speech: gesture, signing, objects of reference, picture and symbol systems, and speech-generating devices. The named manualised picture-exchange programme is described with the other manualised interventions in the behavioural-interventions section of this chapter; the question here is the one the speech and language therapist actually answers, which is what the person needs a communicative route *for*, and how that route is chosen, taught and kept available.

The clinical decision is not which system is best in the abstract. It is which communicative acts are currently impossible for this person, and what would make them possible. A route is worth introducing when there is a real need to request, refuse, seek help, comment or indicate a preference; when the person can physically and perceptually manage it; and when the people around them will respond to it consistently. Assessment therefore examines comprehension as carefully as expression, because a system chosen for output alone will be abandoned by a person who cannot follow what it is being used to negotiate.

The practical discipline is to decide what communicative act the route is meant to make possible and whether the family can keep it available in the settings that matter. A picture system that remains in a folder at school is not a functional communication system at home. Equally, the introduction of a symbol system should not lead adults to stop attending to gesture, vocalisation, movement toward an item, or any other communicative attempt. The method should widen the person's options, not narrow adults' attention.

■ **TRAP Examination trap.** Do not present augmentative communication as a last resort that follows failed speech therapy, and do not describe it as a substitute for therapeutic formulation. It is offered alongside speech and language work from the outset, because an accessible route now does not foreclose speech later; the evidence does not show that a symbol route suppresses talking.

Review asks whether the selected communicative functions have become more available in daily routines, whether the person initiates them with multiple partners, and whether the route itself has become a barrier through poor access, excessive prompting or inconsistent adult response. A system that only works in the therapy room has not yet done its job.

30.3 Occupational therapy: participation, adaptation and sensory questions

Occupational therapy addresses participation in the occupations of ordinary life. Its clinical value lies in observing the fit between a person, a task and an environment, then changing the part of that fit that can be changed. This may involve simplifying a routine, preparing a transition, altering materials, building a practicable self-care sequence, or helping adults recognise what makes an activity inaccessible. The emphasis is not on a universal profile of autism; it is on the particular activity that has become difficult for this person and family.

Sensory integration therapy, associated with Jean Ayres, requires sharper evidence literacy. The IPS narrative frames sensory integration as challenge within a purpose-designed sensory environment, intended to support adaptation in daily activity. It assumes that a child may be receiving too much or too little environmental stimulation. That description explains why the intervention can sound intuitively compelling to families: it connects a visible response to a tangible environment. It is a rationale, however, not an outcome demonstration.

CLINICAL QUESTION	CONSTRUCTIVE RESPONSE	BOUNDARY THAT PROTECTS THE FAMILY
What activity is difficult?	Describe the task, setting, demand and response in functional terms.	Do not infer a single cause from the label “sensory”.
What can change now?	Adapt the routine, materials or adult response and observe participation.	Keep goals visible outside the therapy room.
Is sensory integration proposed?	Discuss its rationale and agree a time-limited, functional review.	Explain the uncertain evidence before commitment.
Are sound-based treatments offered?	Ask what is delivered and what safety safeguards are in place.	Do not treat a sensory rationale as evidence of benefit.

The IPS evidence rating for sensory integration is **Not established, but potentially effective**; its overview also places sensory integration outside the guideline’s recommended nonpharmacological group. This is not a contradiction to be concealed from parents. It is a reason to state the uncertainty plainly, define a functional outcome in advance, and avoid allowing an expensive or time-consuming programme to displace communication, education, family support or medical review.

30.4 Sensory integration and auditory integration: explaining contested evidence safely

A sympathetic explanation is not an endorsement. Sensory distress can be real and can interfere substantially with participation. The clinical response should be respectful observation, environmental problem-solving and a careful search for what the person is communicating

through their response. It should not convert an observation into a certainty that a particular branded intervention will correct an assumed underlying deficit.

Systematic reviews find the evidence base for sensory integration inadequate for recommendation as an evidence-based ASD treatment, with poor study design a central concern. A different textbook account reports **an absence of controlled efficacy studies**. Rather than pretending these summaries are identical, the clinician should communicate their common practical implication: controlled evidence does not establish sensory integration as a treatment for autism. The source-level tension reinforces, rather than weakens, the need for transparent uncertainty.

Auditory integration training needs an even firmer boundary. Reviews do not support a conclusion that it is effective in ASD, and they identify a potential hearing risk from loud sound exposure. The distinction between occupational adaptation and a sound-based treatment matters. A clinician who hears “sensory therapy” should ask what is actually being proposed, what outcome is promised, and whether a potential harm is being minimised in the sales language.

For sensory interventions, validate the difficulty, adapt the environment, define a functional outcome, and state the evidence limit before recommending a programme.

30.5 Family support and parent-mediated work

Family support is clinical work, not a polite adjunct after referral. Families need a coherent formulation, clear goals, practical help in negotiating settings, and space to describe what a proposed programme costs them. A plan that depends on constant availability, specialised materials or flawless delivery may be technically elegant but clinically unsustainable. The psychiatrist should make the burden visible early, especially when multiple services each prescribe home tasks without coordination.

GOOD TO REMEMBER

Before adding a parent-mediated task, ask whether it can be carried across the family’s routines. Engaging parents as **co-therapists** can help maintain treatment effects and generalise learned responses across settings, but the same therapeutic tasks can increase parental stress when they become an added burden. Protect capacity: the parent is a partner whose knowledge of routines is indispensable, not an untrained therapist to be judged for poor adherence.

Preschool parent-mediated social-communication interventions teach parents, carers or teachers strategies that increase **joint attention** and reciprocal communication, often through interactive play and action routines. Their evidence should be read with precision. Improvements are usually found in proximal measures closely related to what was trained or in broader developmental measures; autism behaviours are rarely assessed and longer-term outcomes remain unstudied. This does not make the work pointless. It tells the clinician what result may reasonably be expected and what result should not be promised.

JASPER expands to Joint Attention, Symbolic Play, Engagement and Regulation. It is one of **two** evidence-based ASD interventions recommended by NICE in the cited account. The remaining intervention is not named in that source, so it should not be guessed. **PACT** means Preschool Autism Communication Trial and illustrates the same broad clinical principle: the adult-child interaction is a treatment context that needs skilled support, realistic goals and review of burden.

2

NICE-RECOMMENDED EVIDENCE-BASED INTERVENTIONS IN THE CITED ACCOUNT

NICE found too little evidence to recommend a programme for parental stress, mental health or quality of life, and directs practical intervention towards behaviours that challenge. This is a statement about the evidence base, not permission to ignore carers. Clinicians should still listen to strain, coordinate demands, make plans understandable, and reassess whether the family can live with the regimen. Where family distress requires assessment or treatment in its own right, that work belongs within the broader child psychiatric care framework, including the Child psychiatry: assessment, treatment, services and protection notes.

30.6 Matching expectations to developmental stage and setting

- **RULE Match the intervention to the functional target.** The target must survive in the setting where it actually matters, not only in the therapy room.

Home, clinic and educational settings each reveal different demands. The point is not to prescribe identical practice everywhere, but to make sure that the adults in those settings understand the same goal and respond in compatible ways. This is particularly important when a child is taught a new communication act in a setting but encounters a different routine, material or response elsewhere.

Social-communication programmes can have a role where social isolation is the therapeutic target. Their evidence does not justify comparable claims for restricted, repetitive behaviours or sensory sensitivities. Candidates should avoid claiming that a communication programme treats every domain of autism. It may improve an interactional pathway while leaving another source of difficulty unchanged. That is not a failure of the child or family; it is an instruction to keep the formulation plural and to revise the plan.

For older school-aged children and young people, the evidence for social-skills and peer-mediated work is methodologically limited. A possible signal in favour of more intensive and longer work is not enough for firm conclusions. In clinical practice, this calls for proportionate claims, collaboratively chosen goals and regular review of whether a group or peer-based format is producing meaningful participation rather than merely attendance.

- Write a shared functional aim in language that family and school can use.
- Identify the adult responses that will support that aim in each setting.
- Ask what new burden the plan creates before adding another home task.
- Review benefit, feasibility and unintended effects together.

30.7 The psychiatrist's synthesis and review

The psychiatrist need not deliver every therapy to lead the treatment plan. The task is to ensure that referrals answer a clinical question, that different professionals are not pursuing incompatible goals, and that the family is not asked to carry an impossible programme. At review, ask what has changed in the specified activity, who has observed the change, where it is seen, and what has become harder. A vague report that therapy is “going well” should be translated into observable participation and weighed against burden.

This approach also guards against false choices. It is not speech therapy versus occupational therapy versus family work. Communication access, environmental fit and caregiver capacity are linked, but each needs a distinct assessment and an honest account of evidence. When an intervention is uncertain, the ethical response is neither contempt nor uncritical referral: explain the rationale, disclose the limitation, agree a meaningful review point and retain interventions with clearer functional value. Document the family's own priorities and the practical barriers that may prevent follow-through. At the next review, revise the plan rather than interpreting an unworkable intervention as a failure of motivation.

31 · Prognosis and outcome in ASD

Prognosis is not a single endpoint. In autism spectrum disorder (ASD), outcome has to be described across the abilities and circumstances that make adult life workable: communication, adaptive functioning, education and employment, relationships, autonomy, mental health, safety, and the fit between a person's needs and the support actually available. A child may make striking developmental gains yet continue to need help in daily life; another may meet fewer diagnostic features over time but remain vulnerable at transitions. The useful clinical question is therefore not whether autism will “go away”, but what trajectory is emerging, what risks can be anticipated, and what support will allow the person to participate as fully and safely as possible.

For examination answers, separate the course of autistic features from functional outcome and from service outcome. They overlap but are not interchangeable. A reduction in visible autistic symptoms does not itself prove independence, and a constrained adult outcome may reflect a poor transition environment as much as the person's intrinsic abilities. Prognosis should be communicated as an individual formulation: it recognises uncertainty while identifying practical levers for surveillance, preparation and support.

31.1 The longitudinal frame

Autism is developmental, not a progressive neurological decline. ASD is **not a degenerative disorder**. Learning, adaptation and compensation can continue across the life course, which is why adult outcome cannot be inferred simply from the severity of a child's presentation at one assessment. Development may be uneven: a person can gain language, practical routines or self-awareness while continuing to have substantial difficulty with reciprocity, flexibility, sensory demands or social judgement. Clinical review must therefore look for new demands as well as changes in core features.

At the population level, many people show a **gradual reduction in autistic symptoms** alongside improved adaptive abilities. The most important qualification is the breadth of individual variation. The same diagnostic label includes people whose adult lives are relatively autonomous and people who require continuing intensive support. That variation is not a reason to abandon prognosis. It is a reason to make prognosis descriptive, revisable and grounded in function rather than in a label alone.

■ **RULE** State prognosis as a trajectory of function and support needs, not as a promise of cure or a verdict based on diagnosis alone.

A minority have a more difficult behavioural course in adolescence, whereas improvement is the more usual direction overall across the long run. This distinction is clinically useful. It prevents the erroneous reassurance that developmental progress eliminates later risk, while avoiding the equally damaging assumption that autism inevitably worsens. Behavioural deterioration requires assessment of change in context, physical health, seizures, psychiatric symptoms, environmental demands and the adequacy of support. It should not be dismissed as an unavoidable feature of ASD.

31.2 Diagnostic stability and its meaning

Diagnostic stability is one strand of outcome research, not a complete outcome measure. A systematic review brought together 23 longitudinal studies of children diagnosed using earlier diagnostic categories. Such work asks whether a later profile still crosses a diagnostic threshold. It does not by itself answer whether the person is communicating well, participating in education, safe from exploitation, or living with a satisfying degree of autonomy. A resident should explicitly make this distinction in a long-answer question.

Within the earlier category of **DSM-IV autistic disorder**, the diagnosis was generally stable. At follow-up, however, approximately **15%** no longer met criteria for that earlier category. This should not be translated into a claim that a corresponding proportion of people with present-day ASD lose the diagnosis. The denominator, diagnostic system and question are different. It is better understood as evidence that developmental profiles can change sufficiently for some individuals to cross a categorical boundary.

Earlier labels of Asperger syndrome and pervasive developmental disorder not otherwise specified showed a **more variable diagnostic course**. Follow-up could move in either direction: some profiles no longer met criteria for a pervasive developmental disorder, whereas others became more consistent with autistic disorder. That finding illustrates a broader teaching point. Categories are useful clinical shorthand, but development is dimensional and context-sensitive. A later reassessment should be based on the current developmental history and functional picture, rather than treated as an administrative confirmation of a childhood label.

■ **TRAP** Do not equate a change in diagnostic threshold with normalisation. Diagnostic status, symptoms, adaptive functioning and participation must be reported separately.

31.3 What adult outcome studies actually show

Adult outcome studies are difficult to compare because they use different samples, eras of diagnosis, follow-up methods and definitions of a good outcome. Some use a common classification ranging from very good to very poor, while others emphasise schooling, work, living arrangement or social inclusion. These are related outcomes, but they are not synonyms. The evidence base also contains relatively few prospective studies that follow children from early childhood into adult life. That limitation matters when counselling families: precise-sounding estimates may conceal a very particular cohort and a very particular endpoint.

23 LONGITUDINAL STUDIES IN ONE STABILITY REVIEW	approximately 15% LATER BELOW AN EARLIER AUTISTIC- DISORDER THRESHOLD	between <5% and 25% VERY-GOOD RATING ACROSS PROSPECTIVE COHORTS
--	--	---

Using that common outcome classification, the proportion rated as having a **very good outcome** lay **between <5% and 25%** across prospective childhood-to-adulthood studies. This is not an employment rate, a school-attendance rate or a prevalence estimate. It is a rating on a particular outcome tool. The correct interpretation is that very favourable adult outcomes occur, but their frequency varies substantially by cohort and definition.

Another older, narrower set of estimates concerns children with **childhood autism**, not the contemporary whole ASD spectrum. Around **10% to 20%** improved from early childhood and subsequently attended an ordinary school and obtained work. A further 10% to 20% could live at home but could not work and required special schooling or a training setting. At least **60%** improved little and could not lead an independent life. These figures are worth knowing only with their boundaries intact: they are historical, category-specific outcome tiers, not a universal prediction for all persons now diagnosed with ASD.

OUTCOME QUESTION	WHAT IT CAPTURES	WHY IT MUST NOT BE SUBSTITUTED FOR ANOTHER OUTCOME
Diagnostic status	Whether current features meet a specified threshold	It does not establish independence or quality of life.
Symptom trajectory	Change in autistic features over time	Improvement can coexist with persistent support needs.
Functional outcome	Communication, practical autonomy, participation and work	It depends on both personal skills and opportunity.
Living arrangement	Where and with what support a person lives	Family resources and service availability influence it greatly.
Outcome classification	A study-defined global rating	It should not be relabelled as a single attainment figure.

Improvement is also not synonymous with the disappearance of autistic traits. People with childhood autism who improve may continue to have language problems, emotionally detached interpersonal style, and unusual behaviour. The clinical implication is respectful realism:

celebrate gains without withdrawing supports prematurely, and assess the person's own goals rather than judging success only against a conventional adult template.

31.4 Predictors: useful, but never deterministic

The strongest recurring predictors are markers, not destiny. Better long-term outcome is associated with **higher intellectual ability** and the presence of useful speech before **5 years of age**. A second account places useful speech at age 5 years. The slight difference in wording should not be forced into a false precision. Both formulations identify the early emergence of functional language as a favourable marker, while preserving the fact that people with similar apparent strengths can have very different adult outcomes.

Why are these predictors clinically plausible? Intellectual ability and functional speech can broaden the routes through which a child learns, signals distress, understands rules, uses education and later negotiates environments. They may also make it easier for carers and teachers to recognise strengths and tailor demands. Yet this is interpretation, not a reason to reduce prognosis to a score or a language milestone. Social understanding, emotional regulation, daily living skills, family circumstances, educational fit, co-occurring conditions and adult services can each alter what those strengths mean in real life.

There is also an **optimal outcome** literature describing a minority whose later presentation no longer meets diagnostic criteria. In observational comparisons, this group had four distinguishing features four:

- milder social impairment at baseline, despite comparable early communication and restricted or repetitive difficulties;
- earlier access to intervention;
- more intensive behavioural intervention services; and
- a greater likelihood of receiving applied behaviour analysis rather than pharmacological intervention.

These observations are important but do not prove that any one service caused the outcome. Baseline social differences may themselves influence both eventual outcome and access to an intervention. Service intensity may also be a proxy for family resources, local provision or a clinician's perception of need. This is the distinction a mature answer should make: observational association can guide hypotheses and service planning, but causal claims require a design able to separate treatment effects from pre-existing differences.

■ **RULE** Use prognostic markers to individualise support and follow-up, never to ration intervention or forecast an individual's ceiling.

That caution is especially necessary when discussing earlier identification. There has been **no randomised combined screening-and-treatment trial** demonstrating that screening and an earlier age at diagnosis itself lead to better long-term outcome. This does not make timely recognition unimportant. It means that the causal chain should be stated honestly: earlier

recognition may allow access to help sooner, but the long-term benefit of a screening-and-treatment pathway has not been established by that trial design. The detailed approach to early intervention belongs with the management sections of this chapter.

31.5 Adolescence: where trajectories can diverge

Adolescence increases both opportunities and strain. New educational settings, changing social expectations, bodily changes, growing self-management demands and the approach to adult roles can expose a mismatch between a person's coping strategies and their environment. A previously manageable pattern may become more visible when routines change or supports recede. It is therefore good practice to establish a functional baseline and to ask what has changed before assuming a deterioration reflects the autism itself.

Some studies describe a **deterioration in functioning during adolescence** that may coincide with the development of epilepsy or with the onset of psychiatric disorders. The occurrence of epilepsy in ASD, its assessment and medical management are covered in the medical-comorbidity section. In this prognosis section, its relevance is as a warning that a loss of functioning needs active medical review rather than passive observation.

Similarly, the onset of **co-occurring anxiety or mood disorder** may accompany adolescent deterioration. A changed presentation may be behavioural rather than verbal: withdrawal, irritability, sleep change, avoidance, increased rigidity, loss of skills in familiar routines or escalating distress can all be clinically meaningful. Do not presume that every change is a core autistic feature. The map of psychiatric and medical comorbidity, including how to detect it, is addressed in the medical-comorbidity section; pharmacological treatment of target symptoms is addressed separately.

Regression should also be handled carefully in prognostic discussion. A developmental history of regression may shape a family's fear about future loss, but it is not a licence to make a numerical prediction. When there is new loss of function, the clinician must revisit the differential diagnosis and investigate the current change. The differential of regression is considered in the differential-diagnosis section of this chapter.

31.6 Mortality and preventable risk

Mortality in ASD appears **slightly increased** relative to the general population, but the available evidence is limited and no numerical rate should be implied where one has not been established. Prognostic counselling should avoid alarmist language. The clinical value of raising this issue is not to frighten families; it is to focus attention on preventable hazards and on continuity of medical care.

One proposed pathway concerns medical events such as status epilepticus or accidental death associated with epilepsy. This makes seizure monitoring during adolescence a potentially preventable part of risk reduction when seizures emerge or change. The point is not that every deterioration is epileptic, nor that epilepsy explains all excess mortality. It is that a new seizure history, impaired recovery after an event, or a change in functioning deserves prompt clinical attention.

A change in function in an adolescent or adult with ASD is a clinical event: reassess health, mental state, environment and support before attributing it to autism.

31.7 Transition to adulthood is a prognostic intervention point

Services shape outcome. The exit from school can be profoundly stressful and may slow overall progress. The move into adulthood often reveals gaps in adult mental-health care, social care and developmental-disability provision. This is a service problem as well as a developmental challenge. A young person's competence can appear to fall when familiar structures disappear, travel demands increase, family carers become exhausted, or no one coordinates education, work, benefits, healthcare and daily support.

Transition planning should begin with an account of the person's present strengths, communication preferences, sensory and behavioural needs, health risks, decision-making support, family capacity and preferred adult roles. It should identify what the young person can increasingly manage independently and what assistance remains necessary.

GOOD TO REMEMBER

Make the transition plan concrete: who will notice deterioration, who will coordinate appointments, what happens when routine breaks down, and how meaningful daytime activity will be maintained. This protects dignity while avoiding a sudden and unrealistic withdrawal of care.

The evidence for specific transition-planning interventions remains limited, so clinical reasoning matters. Build continuity wherever possible, prepare the receiving service, and make the handover understandable to the person and family. The broader assessment battery, service organisation and safeguarding framework are set out in the Child psychiatry: assessment, treatment, services and protection notes. The detailed discussion of adult ADHD belongs in the ADHD and specific learning/communication disorders notes.

31.8 A practical way to communicate prognosis

A useful prognostic conversation starts with what is already known and names uncertainty without retreating into vagueness. Describe the person's trajectory in domains: communication, adaptive skills, learning, participation, mental health, physical health, safety and support network. Then identify favourable capacities, current barriers and foreseeable transition points. Families often need help distinguishing a realistic plan from a deterministic prediction. The clinician can say that development may continue, that outcomes vary markedly, and that progress is strengthened when demands and support are matched over time.

In a written formulation, avoid a bare phrase such as "poor prognosis". Instead state the active issues and the plan for each. This gives the team a framework for review and gives the family an intelligible account of why monitoring continues even when the person is doing well.

FORMULATION DOMAIN	QUESTION FOR FOLLOW-UP	CLINICAL USE
Developmental trajectory	What has improved, plateaued or changed?	Separates expected variation from a clinically significant loss of function.
Functional participation	What can the person do with the available support?	Prevents symptom counts from standing in for real-world outcome.
Communication and cognition	How are strengths being used in education and daily life?	Uses prognostic markers without turning them into ceilings.
Health and mental state	Has there been a new seizure, distress pattern or behavioural change?	Prompts review of potentially modifiable contributors.
Transition and supports	What will change when current structures end?	Turns a predictable risk point into a planned intervention point.

The final examination-quality conclusion is balanced. ASD usually permits ongoing learning and adaptation, but outcome is highly heterogeneous. Better intellectual ability and early useful speech are favourable markers, not guarantees. Co-occurring epilepsy or psychiatric illness, and failures at major transitions, can worsen functioning and require active review. The best clinical response is sustained, person-centred planning that measures meaningful function and keeps support available as needs change.

Discriminate, apply and consolidate

32 · Differentiation of intellectual disability from autism, GDD and borderline intellectual functioning

The discriminative task. A developmental presentation is not solved by attaching the most visible label to the child. Intellectual disability, autism spectrum disorder, global developmental delay and borderline intellectual functioning may all come to attention through slow learning, school difficulty, communication concerns or behaviour that unsettles adults. They answer different clinical questions. The useful formulation asks what the person can do in everyday life, how development has unfolded, whether social communication is qualitatively atypical for that developmental level, and whether the current assessment is a valid account of ability.

The marks are usually in the reasoning, not in reciting a checklist. Start by separating a broad developmental profile from a distinctive pattern of social-communication and behavioural difference. Then decide whether the profile has reached the threshold for intellectual disability or is better recorded as **borderline intellectual functioning**. This avoids opposite errors: calling every child with delayed language autistic, and using a reassuring test score to overlook everyday limitations that require support.

32.1 Begin with a developmental formulation, not a label

A **developmental formulation** is a map of strengths, difficulties and their context. It is more useful than an early diagnostic label because the same presenting complaint can arise from very different pathways. Slow acquisition of school skills may reflect a generally low level of functioning, a sensory or language barrier, missed learning opportunities, emotional distress, a neurodevelopmental condition, or an assessment that did not capture the child's usual capacity. The clinician therefore compares domains rather than treating a single complaint as the whole phenotype.

History supplies the frame. Establish the tempo of acquisition and whether abilities have continued to accrue, plateaued, or been lost. Ask how the person communicates needs, joins shared activity, copes with ordinary demands, learns from modelling, and manages changes in routine. Observe the same questions in the interview. A score is evidence, but not an explanation. It must be interpreted with language, sensory, motor, educational and behavioural context in view. This is especially important when the interview setting is unfamiliar or the person is anxious, overactive, shut down, oppositional, exhausted, or unable to understand the task demands.

■ **RULE** **Rule.** Diagnose the pattern of functioning and its real-world consequences; never let a conspicuous developmental symptom decide the diagnosis by itself.

The comparison must be made against **developmental level**, not merely chronological expectation. This prevents a common false positive for autism. A person with broadly delayed development may use simpler language, play, social approaches and coping strategies than peers. The pivotal question is whether social communication is more impaired, qualitatively different, or oddly uneven even when compared with that person's overall developmental level. Conversely, apparently sociable behaviour does not by itself settle the issue: the clinician asks whether contact is reciprocal, flexible, shared and meaningful, rather than simply frequent.

- Define the referral problem in ordinary functional terms before translating it into a diagnostic question.
- Seek convergent information from the person, family, school and direct observation.
- Describe strengths as carefully as limitations, since unevenness often carries the diagnostic clue.
- Record factors that could have distorted the assessment before treating its result as trait-like ability.

32.2 Comparison at the bedside

The table is a reasoning aid, not a substitute for assessment. Intellectual disability concerns the established diagnostic construct and its functional consequences. Autism spectrum disorder concerns a characteristic social-communication and behavioural pattern. Global developmental delay is a developmental description used when a stable intellectual-disability formulation cannot yet be made with confidence. Borderline intellectual functioning identifies a clinically relevant

zone near the boundary, where careful assessment and context matter more than an automatic diagnostic conclusion.

CLINICAL QUESTION	INTELLECTUAL DISABILITY	AUTISM SPECTRUM DISORDER	GLOBAL DEVELOPMENTAL DELAY	BORDERLINE INTELLECTUAL FUNCTIONING
Central task	Establish the intellectual and adaptive profile	Identify a characteristic social-communication and behavioural pattern	Describe developmental delay while the profile is still evolving	Judge a boundary presentation and its clinical relevance
Pattern of difficulty	Broad functional limitations are central	Qualitative difference may coexist with any intellectual level	Delay is described across development rather than presumed permanent	Difficulties may be real without establishing intellectual disability
Key comparison	Everyday function alongside intellectual assessment	Social reciprocity and behaviour relative to developmental level	Current abilities, opportunities and developmental trajectory	Intellectual and adaptive findings, including their mismatch
Common error	Equating a test result with the whole diagnosis	Calling global delay autism without qualitative evidence	Forcing a permanent label before adequate assessment	Using an obsolete numerical rule as a current diagnosis

Use the table dynamically. A person can occupy several categories in a clinically meaningful sense. Autism can occur with intellectual disability, and developmental delay can be the initial description before the later formulation becomes clearer. The task is not to make the categories mutually exclusive. It is to avoid using a category as an explanation for every feature and to state clearly what each diagnosis contributes to care planning.

The decisive comparison is qualitative pattern against developmental level, interpreted alongside everyday functioning, not the presence of delay alone.

32.3 Intellectual disability and autism spectrum disorder

When intellectual disability is being considered, autism must be established by a characteristic qualitative pattern beyond the effects of broad developmental limitation, rather than inferred from delayed performance alone.

GOOD TO REMEMBER

Observe how the person uses the abilities they do possess. Do they seek another person for comfort, enjoyment, help or shared attention? Can they respond to a familiar social bid when demands are low? Is there flexibility in the way people and objects are used? Interpret the answers in the context of communication ability and the environment, not as isolated signs.

In autism, the important finding is a disproportionate alteration in **relative social reciprocity** and in the behavioural pattern, rather than delayed performance alone. A person may have very limited speech yet show clear social interest and shared enjoyment that are consistent with their overall developmental level. Another may have fluent speech but persistently atypical reciprocity, difficulty using communication socially, inflexible interests or restricted patterns that cannot be explained by general developmental delay. The latter pattern warrants an autism assessment even when intellectual ability is not clearly impaired.

Do not set up a false contrast in which one diagnosis excludes the other. When autism and intellectual disability coexist, the formulation should state both and then show what observations support the autism diagnosis beyond global delay. That discipline protects against diagnostic overshadowing in either direction. It prevents clinicians from attributing all behaviour to intellectual disability, while also preventing an autism label from obscuring practical support needs, communication limitations or safeguarding concerns associated with intellectual disability.

■ **TRAP Examination trap.** Delayed language, poor eye contact in an unfamiliar interview, repetitive play at a low developmental level, or behaviour under stress is not by itself evidence of autism. The question is whether the social-communication and behavioural pattern is qualitatively atypical for that person's developmental level.

Clinical interviews should therefore include situations that lower performance pressure. Watch what happens when the examiner follows the person's lead, offers a familiar object, pauses for a response, or involves a caregiver. A narrow clinic observation can overread anxiety, sensory discomfort or communication mismatch as autism, especially when the person has intellectual disability. Equally, a warm greeting or brief eye contact cannot rule autism out. Longitudinal history and reports across settings are what convert an impression into a defensible formulation. Detailed diagnostic criteria and assessment methods for autism belong in the relevant sections of this chapter; here the practical point is to ask whether autism adds a distinctive explanatory pattern beyond general developmental difficulty.

32.4 Intellectual disability and global developmental delay

Global developmental delay is best understood as a developmental description, not as a shortcut to a lifelong conclusion. It communicates that delay is evident across the developmental profile while acknowledging that the available evidence may not yet support a stable intellectual-disability formulation. The distinction matters because a child's observed performance is shaped by opportunity, health, communication access and the feasibility of formal assessment. A label that is too final can close inquiry prematurely; a label that is too vague can fail to communicate support needs.

The key concept is **developmental trajectory**. Ask whether the profile is broadly delayed or strikingly uneven, whether new skills are being acquired, and whether the child's functioning is being assessed under conditions that permit a fair estimate. Repeat assessment may be more informative than forced precision when a child cannot engage with formal tasks, when communication access is poor, or when the environment has substantially limited experience.

The clinician should explain uncertainty plainly: uncertainty is not indecision when it accurately reflects the developmental state of the evidence.

This distinction also changes the conversation with families. The immediate work is to describe present strengths and needs, arrange appropriate developmental and functional assessment, and review change over time. It is not to promise a future intellectual level or to use a temporary descriptive term as a euphemism for a diagnosis that has not been established. The normal developmental framework should be read alongside the *the Normal child development and milestones notes*; assessment, services and protection are addressed in *the Child psychiatry: assessment, treatment, services and protection notes*.

- Use global developmental delay when it accurately describes the current developmental evidence and its limits.
- Do not assume that a broad early delay proves a fixed intellectual-disability formulation.
- Do not assume that later identification of intellectual disability was predictable from one early observation.
- State what must be clarified on follow-up: developmental course, everyday function, communication access and validity of assessment.

The distinction is therefore practical rather than semantic. It determines how confidently the clinician communicates prognosis, how the record represents uncertainty, and why follow-up must revisit the formulation rather than merely repeat a label. It also keeps the assessment open to alternative or coexisting explanations when the profile is not globally uniform.

32.5 Borderline intellectual functioning: category, boundary and limits

Borderline intellectual functioning is not a milder synonym for intellectual disability. In the DSM-5-TR, it is coded **R41.83** and appears in Section II among conditions that may warrant clinical attention. That location is clinically useful: it permits recognition of a relevant cognitive vulnerability without converting every such presentation into a formal mental disorder.

The nosological boundary must be stated accurately. Borderline intellectual functioning is a **condition**, rather than a formal DSM-5 disorder. It should not be treated as a clear-cut disability category or as proof of definite deficits across intellectual and adaptive domains. The category is appropriate when the cognitive profile is itself the focus of clinical attention, or when it materially shapes treatment or prognosis. In practical terms, it can alert the clinician to adjust explanations, consent processes, expectations of independent follow-through, and the format of psychological or pharmacological care, while continuing to assess the actual person rather than acting on a label.

A psychometric zone may be described as **about 70 to 85**, or as one to two standard deviations below the mean, but this is not a current definitional rule. The current DSM-5 approach does not use a numerical IQ criterion to define the category. The older DSM-IV-TR definition explicitly used **IQ 71-84**; importing that band into current diagnostic reasoning is a category error. A

numerical result can prompt a closer look. It cannot replace clinical assessment of function, context and discrepancy.

R41.83 CURRENT CODE	about 70 to 85 DESCRIBED PSYCHOMETRIC ZONE	71-84 OLDER DEFINITIONAL BAND	no numerical criterion CURRENT DEFINITION
-------------------------------	---	--	---

Why this boundary matters. A person can have modest cognitive efficiency yet function capably in familiar, well-supported settings. Another may appear competent until demands increase, language becomes technical, routines change, or treatment requires planning and self-monitoring. Neither observation alone settles whether the person has intellectual disability, borderline intellectual functioning, another condition affecting performance, or a mixture of these. The clinician must avoid both minimisation and overpathologising.

32.6 Separating borderline intellectual functioning from mild intellectual disability

This is the boundary at which disciplined assessment matters most. The current DSM-5 framing expressly compares borderline intellectual functioning with mild intellectual developmental disorder, rather than making a broad contrast with intellectual disability in general. The assessment requires careful appraisal of intellectual and **adaptive functions**, including the **discrepancies** between them. This is not a mechanical exercise. A discrepancy may signal uneven opportunity, a co-occurring disorder, an invalid test session, a skill that is context-bound, or a genuine pattern requiring further understanding.

Work in both directions. If a test result lies near the boundary but everyday functioning is more limited than expected, ask what accounts for the gap before deciding that it demonstrates intellectual disability. Consider communication barriers, educational experience, emotional state, psychiatric symptoms, physical illness, sensory difficulty, family scaffolding and the actual demands of the setting. If everyday functioning appears stronger than a test result, ask whether the person has learned routines exceptionally well, receives extensive unobtrusive support, or had a poor-quality testing experience. The aim is not to rescue a preferred diagnosis. It is to identify the most credible account of the data.

Test validity is central to this judgement. The DSM entry specifically cautions that co-occurring disorders can affect adherence to standardised testing, including schizophrenia and ADHD with severe impulsivity. A low score obtained when attention, motivation, comprehension or behavioural regulation has collapsed is not automatically evidence of stable intellectual ability. Nor should a clinician dismiss all low performance as psychiatric. Reassess the conditions of testing, use collateral evidence, and formulate whether the observed score is likely to represent the person’s usual functioning.

The practical record should make the reasoning visible. State the referral question, the data sources, their limitations, the relation between assessed ability and everyday functioning, and what alternative explanation has been considered. Then state why the conclusion is intellectual disability, borderline intellectual functioning, or a provisional formulation. This is kinder to families and more defensible clinically than an unsupported categorical assertion. Intelligence

testing itself and the instruments used for it are addressed in the assessment section of this chapter; do not confuse such testing with a complete functional evaluation.

32.7 Coding, language and the historical trap

The current label has a complicated historical trail, which explains why outdated terminology still appears in records and teaching material. In DSM-II, the 1968 entry used the label “borderline mental retardation” and the code 310. In 1973, the American Association on Mental Deficiency, now AAIDD, eliminated that name in favour of “borderline intellectual functioning”.

The coding location also shifted. DSM-III placed it on Axis I with V62.89 in 1980. DSM-III-R moved it to Axis II with V40.00 in 1987; DSM-IV later used Axis II V62.89 in 1994. Axes were removed in DSM-5 in 2013, which used R41.83.

This history should sharpen, not confuse, current practice. It shows why an old numerical band, an old axis placement, or stigmatising terminology is not a substitute for a contemporary formulation. Use language that describes the person’s current functioning and support needs. When older records are encountered, translate their meaning cautiously and verify what assessment was actually done before carrying a label forward.

Finally, “borderline” in borderline intellectual functioning has no relationship with borderline personality disorder. Similar wording is not shared aetiology, shared diagnosis or a reason to infer one condition from the other. Keeping this distinction explicit is particularly important when a patient has emotional dysregulation or interpersonal difficulty, because such features can tempt clinicians into semantic rather than phenomenological reasoning.

32.8 A defensible conclusion and its clinical use

Clinical attention is the right frame for borderline intellectual functioning. The category may be recorded when it changes how treatment is delivered or how prognosis is understood, not as a casual descriptor of anyone who struggles academically. Explain information plainly, check comprehension through the person’s own account, anticipate difficulties with abstract instructions, and involve supports appropriately while respecting autonomy. These are good clinical practices because they respond to observed needs, not because a code has replaced assessment.

In a written answer, a strong conclusion follows a consistent sequence. Identify whether the central issue is broad functional limitation, a distinctive autism pattern, an evolving developmental profile, or a boundary cognitive presentation. State the observations that support that judgement. Name any coexisting diagnosis separately. Finally, state what must be reassessed and how the formulation changes communication, support and follow-up. This makes the differential useful at the bedside rather than a set of mutually exclusive examination labels.

Diagnostic humility is not vagueness. It means using a provisional developmental description when the evidence is still developing, making an autism diagnosis only when its qualitative pattern is established, recognising intellectual disability through the established formulation rather than a score alone, and using borderline intellectual functioning accurately when it is

clinically relevant without overstating its status. For associated attention and learning questions, use the ADHD and specific learning/communication disorders notes.

33 · Four worked cases

Vignettes test clinical organisation, not label recognition. A short stem rarely gives a complete developmental history, a valid cognitive assessment, an adaptive profile and a collateral account. The candidate's task is to recognise what the available pattern supports, state what remains uncertain, and choose the next discriminating enquiry. The same discipline protects clinical practice from premature labelling.

Use each vignette as a **developmental formulation**: place the person's trajectory, everyday function, social-communication pattern, repetitive or restricted behaviour, medical context and environment alongside each other. A diagnostic label is the conclusion of that process, not a substitute for it.

Trajectory DEVELOPMENTAL COURSE	Function EVERYDAY INDEPENDENCE	Pattern SOCIAL COMMUNICATION AND BEHAVIOUR	Context MEDICAL, FAMILY AND EDUCATIONAL SETTING
---	--	---	--

33.1 Read the stem before naming the diagnosis

The useful first question is not “Which disorder is this?” but “What is impaired, in which settings, and since when?” Separate a person's learning and problem-solving capacity from **adaptive functioning**, and separate both from the quality of social reciprocity and the behavioural pattern. School difficulty, delayed language, disruptive behaviour and poor self-care are findings. None is diagnostic in isolation.

■ **RULE** **Rule.** Diagnose the pattern of impairment, then test whether the pattern is global, autism-specific, developmental but still evolving, or below expected ability without the pervasive functional deficits required for intellectual disability.

- **Trajectory:** establish whether development has been delayed, uneven, static or marked by loss of previously acquired abilities.
- **Current function:** ask how the person manages communication, self-direction, daily routines, safety and participation in ordinary settings.
- **Pattern and context:** seek social-communication differences, restricted or repetitive behaviour, sensory concerns, physical clues, medical symptoms and the accounts of people who know the person well.

33.2 The central discrimination

Intellectual disability remains an assessment question when a stem describes broad difficulty with learning and everyday function. An autism-spectrum formulation depends on its characteristic social-communication and behavioural pattern. These formulations may coexist. The presence of autism does not answer the question of intellectual functioning, and global learning difficulty does not by itself establish autism.

FINDING IN A STEM	WHAT IT SUGGESTS	WHAT PREVENTS AN OVERCALL
Broad difficulty learning and carrying out everyday tasks	Consider a global impairment formulation	Clarify adaptive functioning and the developmental course
Qualitative difference in reciprocal social communication with a restricted or repetitive pattern	Consider an autism-spectrum formulation	Do not infer intellectual disability from the autism label alone
Developmental concerns in an early developmental presentation without a stable assessment picture	Use global developmental delay as a working developmental formulation where appropriate	Plan review as developmental information becomes clearer
Academic or occupational struggle with relative preservation of everyday adaptive competence	Consider borderline intellectual functioning and competing explanations	Do not convert a low score, poor scholastic performance or social disadvantage directly into intellectual disability

■ **TRAP** **Exam trap.** “Delayed speech” is not a shortcut to autism, and “poor school performance” is not a shortcut to intellectual disability. The discriminating evidence is the whole developmental and functional pattern.

33.3 Vignette: broad developmental difficulty without an autistic pattern

A child is brought because progress across learning, communication and self-care has been slow. The family describes a steady developmental course and needs help across ordinary routines. In the consultation, the child seeks shared engagement in a manner that is proportionate to developmental level, with no clear history of a restricted or repetitive behavioural pattern.

The defensible formulation is that the stem raises concern for intellectual disability or, when developmental assessment remains incomplete, a developmental-delay formulation. The point is not to exclude autism because the child is friendly, nor to diagnose autism because language is delayed. Ask for formal assessment of ability and adaptive behaviour, collateral accounts across settings, developmental history, physical and neurological review, and an aetiological evaluation guided by the wider presentation.

Why the distinction matters. The intervention plan must address the actual pattern of support needs. If autism is added without its defining pattern, families may be diverted from the assessment and supports that explain the person's everyday difficulty.

33.4 Vignette: autism with an uneven profile

An adolescent has intense, narrow interests, inflexible routines and marked difficulty interpreting reciprocal social interaction. Academic achievement is strong in selected areas, while practical independence and social judgement are variable. The stem offers no valid basis to conclude that intellectual functioning is globally impaired.

The appropriate formulation is autism spectrum disorder with a separate assessment of intellectual and adaptive functioning. Uneven strengths do not cancel clinically important disability, but neither do social difficulty and inflexibility prove intellectual disability. Document

the areas of competence as carefully as the areas of difficulty, because support planning follows function rather than assumptions about a diagnostic label.

- Ask which difficulties are intrinsic to social communication and behavioural flexibility.
- Ask whether practical dependence reflects limited capacity, limited opportunity, anxiety, environmental mismatch or a combination.
- Describe co-occurring attention, mood, anxiety, sleep, neurological and medical concerns without attributing all distress to autism.

33.5 Vignette: the apparent boundary case

A young person has struggled academically for years and has been described as “slow.” At home and in the community, the person manages familiar routines, relationships and responsibilities with little extra help. An isolated test result is quoted in the referral note, but its validity, context and relation to everyday functioning are not described.

This stem is designed to expose **score determinism**. Consider borderline intellectual functioning, specific learning or communication disorders, attention problems, educational deprivation, sensory impairment, emotional disorder and contextual barriers. For the detailed assessment battery, see the Child psychiatry: assessment, treatment, services and protection notes; for attention and learning differentials, see the ADHD and specific learning/communication disorders notes.

GOOD TO REMEMBER

Do not diagnose from a number or a pejorative description. Interpret formal assessment alongside adaptive function and developmental history.

33.6 Vignette: change after an established developmental course

A caregiver reports that a child who had acquired skills is no longer using them in the same way. The current presentation includes communication difficulty and repetitive behaviour. The temptation is to treat the current surface pattern as sufficient for an autism diagnosis.

Here the clinical priority is a careful account of **regression**, its tempo, the affected skills, associated neurological or medical symptoms, and the possibility of an alternative developmental or neurological condition. Autism may remain part of the formulation, but loss of skills changes the workup and makes a broad differential essential. Do not reassure, diagnose or narrow the enquiry before establishing the trajectory.

MNEMONIC

MAPS

Map the developmental trajectory. Assess everyday adaptive function. Pattern-match social communication and restricted or repetitive behaviour. Search for syndromic, neurological, psychiatric and contextual signals. This is a retrieval map, not a substitute for assessment.

33.7 Turning a vignette into a clinical answer

A strong answer starts with a provisional formulation, gives the positive findings that support it, then names the information required to distinguish nearby alternatives. It states comorbidities and safety or medical concerns as possibilities to assess, rather than presenting them as conclusions not contained in the stem. A developmental diagnosis does not make new distress, behaviour change or physical symptoms self-explanatory: assess each reported change on its merits.

When asked for management, link recommendations to the person's identified functional needs and family priorities. Do not leap from a label to a named programme, medication or legal entitlement. The relevant chapters provide the detailed intervention, medical-workup, disability-certification and family-support pathways.

In every developmental vignette, the diagnosis must explain the whole pattern of trajectory, function and behaviour better than its nearest alternative.

34 · Consolidate: the master mnemonic, the retrieval prompts and the synthesis map

Consolidation is a clinical discipline, not a list of labels. Intellectual disability, autism spectrum disorder, developmental delay and borderline intellectual functioning can overlap in a referral, yet each answers a different clinical question. The task is to preserve that distinction while building a single account of the person’s abilities, difficulties, context and needs.

A high-quality answer therefore moves from the presenting concern to a **developmental formulation**. It does not let a test result, a screening result, a certificate or a diagnostic label substitute for the history, observation and functional account. The chapter’s component sections supply the detail; this closing section shows how to hold them together without flattening them.

MAP THE DEVELOPMENTAL COURSE	READ THE FUNCTIONAL PROFILE	CHECK CO-OCCURRENCE AND CONTEXT	PLAN MEANINGFUL SUPPORT
---	---------------------------------------	--	-----------------------------------




Fig 21.22 The chapter as one circuit. Intellectual disability and autism spectrum disorder set side by side across a shared spine of six questions, concept to cause to recognition to measurement to management to entitlement, asked of both conditions in the same order. The plate carries the shape of the chapter and no numbers at all, because every number in this chapter binds to the one construct it measures and lives with the fragment that owns that construct. The join band states the thing the side-by-side layout is there to teach, that the two columns are not alternatives and the same person can occupy both.

34.1 Start with the developmental question

The starting task is to establish the **developmental trajectory**: what was acquired, what is currently difficult, whether skills have been lost, and how the pattern affects everyday life. This makes the assessment developmental rather than merely behavioural. It also keeps current symptoms in relation to the person’s communication, cognitive and social capacities.

Developmental history should be interpreted alongside the account in the Normal child development and milestones notes. If the presentation includes regression, unusual episodic events, altered sleep, pain, sensory difficulty or a change from baseline, the formulation must remain open to medical and psychiatric contributors rather than assuming that all change belongs to the neurodevelopmental condition.

■ **RULE** **Rule.** Begin with the person’s developmental course and present functioning; use diagnostic categories to clarify the pattern, not to replace the formulation.

34.2 Separate constructs before combining them

Intellectual functioning, adaptive functioning, developmental level and the autism phenotype are related but not interchangeable. A cognitive assessment addresses an aspect of ability. Adaptive functioning asks what the person manages in ordinary settings. Autism assessment asks whether a characteristic pattern of **social communication** differences and **restricted and repetitive behaviour** is present. Each finding changes the meaning of the others, but none can be inferred from another alone.

CLINICAL TASK	QUESTION TO KEEP SEPARATE	WHERE TO RETURN FOR FULL TEACHING
Describe ability	What can the person understand, learn and do?	Intellectual disability definition, clinical assessment and IQ testing
Describe everyday function	What support is needed in usual environments?	Adaptive behaviour assessment and intellectual disability severity
Describe autism features	Is the social-communication and behavioural pattern characteristic of autism?	Autism diagnostic criteria, clinical features and diagnostic tools
Describe developmental concern	Is the available profile sufficient for a stable diagnostic conclusion?	Global developmental delay and differentiation of intellectual disability from autism, GDD and borderline intellectual functioning
Describe participation	What changes at home, in education, in work and in the community are required?	Management, family support, disability certification and benefits

■ **TRAP** **Examination trap.** Do not use an intelligence score as a synonym for adaptive functioning, or use either as proof for or against autism. They are different clinical constructs.

34.3 Read the profile, not only the label

A useful formulation gives a profile of strengths and vulnerabilities. It asks how communication is used, how social reciprocity is experienced, how routines and sensory demands affect participation, and how demands change behaviour. It records what helps the person engage, learn and regulate. This profile is often more useful for care planning than a diagnostic word alone.

- Describe the person's effective means of communication and the situations in which communication breaks down.
- Identify abilities that can be recruited for learning, self-care, relationships and participation.
- Notice environmental triggers, including mismatch between demands and capacity.
- Ask whether behaviour is communicating distress, pain, fear, overload, confusion or an unmet need.

This approach protects against **diagnostic overshadowing**: the error of attributing every new symptom or behaviour to an established developmental diagnosis. A change from baseline still deserves ordinary clinical curiosity. The full child psychiatric assessment battery, services and protection are set out in the Child psychiatry: assessment, treatment, services and protection notes.

34.4 Look actively for co-occurrence

Co-occurrence is clinically important because it can alter presentation, risk, family burden and the practical plan. Medical conditions, sensory problems, sleep disturbance, emotional symptoms, behavioural disturbance and psychiatric syndromes may make a familiar developmental profile look suddenly more difficult. The right response is not diagnostic proliferation; it is a disciplined search for treatable contributors and an account of how they interact.

When autism and intellectual disability coexist, describe each profile explicitly. When attentional symptoms or social difficulties raise the possibility of ADHD, use the ADHD and specific learning/communication disorders notes for the full assessment framework. Do not treat social difficulty as automatically explained by either condition.

MNEMONIC

FIND

is a practical closing mnemonic: **F**unction in daily life; **I**nteraction and communication profile; **N**eurodevelopmental course; **D**rivers of change, including co-occurring conditions and context.

34.5 Convert formulation into support

The endpoint of assessment is a plan that is intelligible to the person and family and usable across settings. State the priority difficulty, the strengths that can be used, the support required, the contributors that need treatment or further assessment, and the follow-up question. A plan framed around **support needs** is more clinically honest than one framed around a label alone.

- Link recommendations to a named functional difficulty rather than to diagnosis alone.

- Make family partnership part of the formulation, not an afterthought.
- Coordinate educational, communication, occupational, behavioural and mental-health input around shared goals.
- Review change over time, especially after a new stressor, treatment or transition.

Intervention models, speech and occupational support, family support, pharmacotherapy and prognosis are considered in their respective sections.

GOOD TO REMEMBER

Keep statutory processes separate from clinical formulation: certification may help access entitlements, but it does not replace an individualised account of function and care.

34.6 The final answer in practice and in examination

For a case discussion, write the problem representation before writing the diagnosis: developmental course, current profile, functional impact, possible co-occurrence, contextual modifiers and priority supports. This structure prevents the common drift from a complex clinical presentation into a label-led answer.

The must-remember synthesis is simple: diagnose carefully, describe function clearly, search for change and co-occurrence, then plan support around the person rather than the label.

Previously asked

Genuine past questions from this chapter's territory, with their board of origin. Only two examination questions in the catalogue fall inside intellectual disability and autism, and both are printed here. No question is reconstructed, paraphrased into a new one, or invented to pad the list.

QUESTION	TYPE	ASKED
Describe the etiology, clinical features and management of Fragile X syndrome	Long essay	MUHS 2013
Asperger's syndrome	Short note	MUHS 2012

- **TRAP** **Asperger's syndrome no longer exists as a diagnosis.** A question asked under the older nomenclature still expects you to know where the construct went and why it was absorbed, not to defend it as a current category. The historical topic in this chapter carries that story.

The quick review

One table, read down the left column, cover the right. Every line is taught in full somewhere above; nothing here is new. Where the sources disagree, the disagreement is the answer.

ASK	HOLD
WHAT MAKES INTELLECTUAL DISABILITY	All three criteria: intellectual deficits, adaptive deficits, onset in the developmental period. Not an IQ score on its own.
WHAT GRADES ITS SEVERITY	Adaptive functioning across the conceptual, social and practical domains. IQ is supporting evidence, not the grade.
WHAT GRADES AUTISM SEVERITY	Support needs, rated separately for the two domains. Never IQ. Grading autism by IQ is the classic error.
GLOBAL DEVELOPMENTAL DELAY	The under-fives placeholder for a child too young or too impaired for a valid standardised assessment. It is a holding position, not a diagnosis to keep.
THE FOUR INSTRUMENTS ROUTINELY CONFUSED	They measure four different constructs: intelligence, development, social maturity, adaptive behaviour. Being asked for adaptive scales and listing all four is the exam error this chapter exists to prevent.
THE STATUTORY INSTRUMENT FOR INTELLECTUAL DISABILITY IN INDIA	Not the mental-illness disability scale. The routes are separate, and the chapter shows which is which.
THE AUTISM DYAD	Social communication, all of its sub-criteria; restricted and repetitive behaviour, the stated minimum of its four. Language is a specifier, not a criterion.
WHERE ASPERGER'S WENT	Absorbed into the spectrum. A past question under the old name still expects you to know why.
THE COMMONEST PSYCHIATRIC CO-OCCURRENCE IN AUTISM	ADHD. DSM-IV forbade the joint diagnosis; DSM-5 permits it. The frequency is contested and the chapter gives both figures.
DIAGNOSTIC OVERSHADOWING	A new behaviour in intellectual disability gets blamed on the disability and the treatable illness is missed. It is a reasoning failure, not a knowledge failure.
WHAT NO DRUG DOES IN AUTISM	Treat the core features. The licensed agents target irritability, and the monitoring burden is the price.
WHAT THE SOURCES REFUSE TO TELL YOU	Several cut-offs, several band tables and several Indian figures are simply not in the literature consulted. The chapter says so where it happens rather than inventing them.

References

Every specific in this chapter is bound to one of the sources below, and the count is the number of claims each carried. Books and guidelines are cited author, title and year in the house convention; no PMID is asserted for a book. Where two sources disagreed, the chapter shows both rather than choosing, and those disagreements are named in the text where they fall.

SOURCE	CLAIMS CARRIED
IPS CPG CAP CHILD	165

SOURCE	CLAIMS CARRIED
Tasman Psychiatry	119
Kaplan & Saddock's CTP 2024	118
Companion to Psychiatric Studies	107
DSM-5-TR-2022	98
New Oxford Textbook of Psychiatry	85
Shorter Oxford Textbook of Psychiatry	69
RC Jiloha - Forensic Psychiatry - An Indian Perspective	59
ICD-11 CDDR	58
Rutters Child and Adolescent Psychiatry 6E	47
Lewis Child and Adolescent Psychiatry	46
Rutter's Child and Adolescent Psychiatry 6E	40
DSM-5-TR (2022)	39
Dulcan's Textbook of Child and Adolescent Psychiatry	33
Maudsley 15th edition	33
Shorter Oxford Textbook of Psychiatry, ch 17	28
Kaplan & Sadock's CTP 2024	26
Kaufman's Clinical Neurology for Psychiatrists	23
61:254-69)	17
Kaplan & Sadock's CTP 2024, intellectual disability chapter	17
Tasman Psychiatry, ch 42	17
Clinical Methods in Psychiatry 1st Ed 2024, AIIMS Publication	16
IPS CPG CAP CHILD - Kishore et al., Indian J Psychiatry 2019	15
Geriatric Psychiatry Study Guide	12
New Oxford Textbook of Psychiatry, Chapter 24	12
IPS CPG CAP CHILD (Kishore et al.)	12
Developmental Psychopathology (Cicchetti)	11
Dulcan's Textbook	11

On what is not here. This chapter declares its gaps rather than filling them. Where a value a reader might expect is absent, the absence is deliberate and stated at the point of use: the source consulted did not carry it, and nothing was

supplied from memory to cover the silence.

Index

Every entry jumps to its page. Numbers are the page of first mention.

#

6A00 172

A

a-not-b error 12
accommodation 13
accompanying intellectual impairment 296
Acquired epileptic aphasia 344
acquired microcephaly 344
adaptability 55
adaptation 13
adaptive functioning 55
ADI-R 328
adolescence 12
adolescent development 49
adolescent neurodevelopment 122
ADOS-2 329
adrenarche 118
adverse childhood experience 148
aetiological formulation 185
affective theory of mind 64
ainsworth 25
ainsworth's strange situation 25
Alcohol-related neurodevelopmental disorder 242
allostatic load 151
Alpha-fetoprotein 277
ambitendency 41
ambivalent 49
ambivalent or resistant attachment 156
anaclitic depression 145
anal stage 17
anal-expulsive 28
anal-retentive 28
animism 14
ann masten 150
anomaly 243
anti-D antibody 276
appearance 14
Applied behavioural analysis 359
aripiprazole 364
artificialism 14
ascertainment bias 306
Asperger's syndrome 289
assessment 15

assimilation 13
assisted performance 45
associated-condition specifier 289
associations 27
asymmetric tonic neck reflex 95
asymmetry 84
attachment 21
attachment behavioural system 49
attachment-pattern 21
AuDHD 336
Auditory integration training 374
authoritarian parenting 77
authoritative parenting 76
Autism (ISAA) 281
Autism Communication Trial 358
autism spectrum disorder 40
Autistic disturbances of affective contact 289
autistic personality disorder 289
autistic symptomatology 354
autoclitic 72
autonomy versus shame and doubt 17
autosomal dominant 249
avoidant 49

B

b. f. skinner 69
babinski reflex 112
Baseline exaggeration 262
basic trust versus mistrust 20
baumrind 43
baumrind parenting styles 43
Behavior Checklist 333
behaviour 13
behavioural control 77
behavioural equivalents 265
behavioural indicators 186
Behavioural intervention 272
behavioural phenotype 232
biopsychosocial 74
blood or urine 237
borderline intellectual functioning 172
borderline personality organisation 42
bowlby 49
breath-holding 260
broader autism phenotype 309
bucharest early intervention project 147

C

care	17
caregiver sensitivity	51
CARS-2	329
Case ascertainment	306
castration anxiety	28
catatonia	296
catatonia specifier	336
cell-free fetal DNA	277
centration	12
cephalocaudal	85
challenge	42
change-of-location task	65
Childhood disintegrative disorder	289
chomsky	43
chomsky's nativist theory	72
Chromosomal microarray	215
chromosome 14, designated t(14;21)	223
chronic gastrointestinal complaints	335
class-inclusion task	16
classification	12
Clinical assessment	60
Clinical judgment	182
clinically significant impairment	296
Co-occurrence	325
co-regulation	80
co-therapists	374
cognitive development	11
Cognitive disintegration	262
cognitive profile	67
cognitive theory of mind	64
cognitive-developmental	33
cold cognition	117
combinatorial reasoning	15
commitment	22
comparison	21
compensatory model	151
competence	16
Composite or index score	201
conceptual, social and practical domains	190
concrete operational	11
concrete operational stage	12
conservation	11
conservation task	11
contact comfort	146
context-sensitive risk-taking	116
contextual formulation	173
control of the sleep cycle	260
conventional level	33
cooperative play	104
Copy-number variants	309

Copy-number variation	311
core features of autism spectrum disorder	364
count	17
course	11
critical period	152
cumulative risk	147
cumulative-risk model	148

D

DASII	31
DDST-II	137
decentration	12
deficits in adaptive functioning	182
deficits in intellectual functions	182
demandingness	27
denominator	187
deprivation	50
deprivation-specific pattern	147
developing, maintaining, and understanding relationships	296
developmental arrest	353
developmental assessment	109
developmental assessment scales for indian infants	137
developmental cascades	151
Developmental course	184
developmental delay	26
developmental formulation	26
developmental history	17
Developmental language disorder	139
developmental level	45
developmental quotient	137
developmental red flag	31
developmental red flags	31
developmental screening	11
developmental screening test	137
developmental surveillance	11
developmental trajectory	116
developmental-theory alignment	91
deviance	135
deviation from normal	31
Diagnostic masking	262
Diagnostic measures	329
diagnostic overshadowing	148
diagnostic reductionism	174
Diagnostic stability	377
Diagnostic substitution	305
dialectical	45
diana baumrind	77
differential susceptibility	151
differentiation subphase	40
disappearance	28

discrepancies	321
disinhibited social engagement	50
disinhibited social engagement disorder	50
disorder of intellectual development	180
Disorders of intellectual development	181
disorganised	49
dissociation	135
distinction	35
domain-specific	16
domains of child development	17
domains-and-milestones framework	91
Down syndrome	221
dual diagnosis	256
dual-systems	25
dynamic assessment	30
dystonic	19

E

early adversity	141
early developmental period	172
early infantile autism	289
Early intensive behavioural intervention	358
Early intervention	141
Early Start Denver Model	358
echoic	72
echolalia	318
ecological validity	174
EEG	329
ego integrity	23
ego strength	19
egocentrism	12
electra complex	28
elicited	59
emmy werner	150
emotional refuelling	41
empathy	36
english and romanian adoptees study	147
environment	21
epigenetic principle	19
Epilepsy	232
equifinality	145
equilibration	13
era study	147
erik erikson	18
erikson	12
ethic of care	36
executive-function accounts	67
expected range	34
experience-expectant plasticity	148
expert clinical genetics consultation	215
exploration	15
expressive language	68

F

Facial angiofibromas	248
fading	46
false-belief	43
false-belief task	43
family systems theory	81
female germline	229
Fetal alcohol spectrum disorder	241
fetal alcohol syndrome	241
fidelity	18
fine motor	84
first maternal meiotic division	224
first-order false belief	65
fixation	26
Flynn effect	184
FMR1	229
FMRP	229
for social interaction	296
foreclosure	22
formal operational	11
formal operational stage	15
formal operational thought	30
Fragile X syndrome	228
fragile X-associated primary ovarian insufficiency	229
free trisomy	222
freud	12
freud's psychosexual theory	26
frontal release sign	115
function	22
functional profile	60
FXTAS	229

G

Galactosaemia	239
gender expression	22
gender identity	22
gender incongruence	121
gender role	80
gender-schema theory	131
generativity	18
generativity versus stagnation	20
Genetic architecture	217
genital stage	23
Gesellian model	207
global developmental delay	137
gonadarche	122
good fit	55
goodness of fit	54
goodness-of-fit	24
grading	182

gross motor	31
-------------------	----

H

Habilitation	47
hamartin	249
hand movements	352
Hans Asperger	289
hatching	12
heinz dilemma	33
Heritability	217
heterogeneity	214
higher IQ	222
higher mental functions	44
Homocystinuria	238
hope	18
hospitalism	145
hot cognition	117
Hoyme criteria	243
Hurler syndrome (MPS I)	239
hypermentalizing	68
hypomentalizing	68
hypothetico-deductive reasoning	12

I

identity achievement	23
identity diffusion	119
identity formation	22
identity statuses	22
identity versus role confusion	19
imaginary audience	118
impulsivity	117
inborn error of metabolism	236
INCLEN Tool	281
increase	46
independent performance	24
indicated	31
indicated prevention	275
individuation	12
industry versus inferiority	17
informed clinical judgment	329
initiative versus guilt	17
inner speech	45
institutionalization	141
integrity versus despair	20
intellectual assessment	187
Intellectual disability	68
Intellectual distortion	262
Intellectual functioning	47
Intelligence	33
interest	15
interests	29

internal working models	51
internalisation	28
intimacy	18
intimacy versus isolation	20
intraverbal	72
Iodine deficiency	276
irreversibility	12
Irritability	145
irritability associated with autistic disorder	367

J

james marcia	119
Jean Ayres	373
jean piaget	11
joint attention	63
Joint Attention, Symbolic Play, Engagement and Regulation	358
Jones and Smith	243

K

kohlberg	12
kohlberg's moral development	92

L

language acquisition	43
language acquisition device	72
language and sociocultural accounts	67
language and speech	84
language-impairment specifier	323
latency stage	28
lawrence kohlberg's theory of moral development	32
learning leads development	47
Leo Kanner	289
Lesch-Nyhan syndrome	239
lev vygotsky	44
lev vygotsky's sociocultural theory	44
libidinal object constancy	17
libido	17
limited guardianship	281
longitudinal history	173
love	18

M

M-CHAT-R/F	329
mahler	12
mand	15
margaret mahler's separation-individuation theory	37
maternal deprivation	50
maturational imbalance	25

maturational-imbalance model	117
measurement error	182
MECP2	351
median	12
mediation	37
Megalocephaly	312
mental dq	31
mentalizing	62
Metabolic screening	215
metacognition	15
michael rutter	150
milestones by age	86
modular or domain-specific accounts	67
monitoring	77
monotropy	50
moral development	17
moral judgement	17
moratorium	22
moro reflex	114
mosaicism	222
motor dq	31
mTOR hyperactivity	313
multifinality	145

N

nativist theory	72
neglect	14
neural tube defects	276
neurodevelopment of adolescence	101
Neurofibromas	253
neurofibromatosis type 1	248
neurofibromin	252
Neuroimaging	215
new york longitudinal study	55
Newborn screening	241
No behavioural phenotype is invariant.	256
noam chomsky	69
normal autistic phase	39
normal symbiotic phase	39
normalization	270
norman garmezy	150
not associated with prolactin elevation	369

O

object	12
object permanence	12
object-permanence task	16
obsessive-compulsive disorder	336
oedipus complex	28
onset during the developmental period	185
operant account	71

operation	11
Optimal outcome	379
or speech	262
oral stage	27
organisation	11
Outcome	13
overgeneralisation	72

P

palmar grasp	90
parallel play	104
Parent-mediated social-communication intervention	374
parenting style	43
pattern of loss	350
PECS	356
performance	13
performance test	204
Peripheral hyperserotonaemia	312
permissive parenting	76
persistence	21
person with benchmark disability	281
personal fable	118
perspective-taking task	16
pervasive developmental disorder	289
phallic stage	17
phenylalanine hydroxylase	236
piaget	11
Picture Exchange Communication System	358
Pivotal Response Treatment	358
placement	54
plantar grasp	112
plateau	90
pointing	64
poor fit	55
Positive behavioural support	272
positive reinforcement	71
postconventional level	35
poverty-of-the-stimulus argument	72
Practice effects	184
practising subphase	40
Pragmatic language	318
pragmatics	74
preconventional level	33
prefrontal cortex	117
Premutation	228
prenatal aetiology	214
Prenatal valproate exposure	313
preoperational	11
preoperational stage	12
presenting concern	173
prevalence	55

Primary prevention	274
primitive reflexes	90
private speech	30
privation	50
probabilistic association	144
productivity	71
prognostic marker	336
prolonged QT interval	353
promotive factor	150
propositional thought	12
protective factor	59
protective factors	128
protective or buffering model	151
proximodistal	85
proxy measure	186
psychological control	77
psychosexual development	22
psychosexual stages	17
psychosocial crisis	18
psychosocial deprivation	139
psychosocial development	122
Psychosocial masking	262
psychosocial moratorium	22
psychosocial stages	17
puberty	22
purpose	14
Pyridoxine	238

Q

quasi-autism	347
--------------------	-----

R

rapprochement crisis	17
rapprochement subphase	41
ratio IQ	197
re-emergence	112
reactive attachment disorder	50
reassessment	172
receptive language	87
red-flag	17
red-flag threshold	39
reflex table	90
refrigerator mother theory	289
regression	25
rené spitz	145
repeat expansions	215
repeat testing	188
resilience	143
resilience map	153
resistant	49
responsiveness	43

restricted and repetitive behaviour	319
Restricted and repetitive behaviours	370
Restricted and repetitive patterns	298
retrieval spine:	134
Rett syndrome	256
Rett syndrome remains a clinical diagnosis	351
risk factor	59
risk-taking	25
risperidone	364
rob(21;21) fusion	228
role confusion	12
rooting	112

S

sampling frame	303
Savant or splinter skills	315
scaffolding	17
scheme	13
score determinism	391
screening	11
second-order false belief	65
Secondary prevention	274
secure	13
secure base	25
Seguin Form Board	197
selective	49
Selective mutism	346
self-injurious behaviours	369
self-recognition	64
sensation seeking	127
sensitive period	122
sensitive periods	147
sensorimotor	11
sensorimotor stage	13
Sensory integration therapy	373
Sensory reactivity	319
separation	12
separation anxiety	85
separation-individuation	17
sequence	11
seriation	12
Severity classification	188
severity level	207
sex characteristics	127
sexual behaviour	22
sexual identity	130
sexual orientation	22
Sherman paradox	229
Shevell et al. (2008)	177
simulation account	67
skinner	43
sleep phenotype	336

slightly increased	380
slow-to-warm-up temperament	58
social (pragmatic) communication disorder	172
social and emotional	84
social and emotional development	84
social anxiety disorder	349
social communication	75
social quotient	206
Social reciprocity	84
social smile	85
social speech	46
social-cognitive theory	131
social-emotional reciprocity	296
social-interactionist	74
social-learning theory	131
sociocultural theory	12
specifier	191
speech	30
spotting deviation	31
statistical learning	74
steeling	151
stepping	112
stereotyped hand wringing	256
stereotypy	256
strange situation	25
strange situation procedure	52
stranger anxiety	85
strong	24
structured TEACCHing	361
subependymal giant cell astrocytomas	250
sublimation	28
Subtest scaled score	201
sucking	27
superego	28
supplementation	276
support needs	190
support plan	173
surrogate marker	176
symbiosis	38
symbolic play	21
syndrome	59
synthesis map	30
synthesis map:	102
syntonic	19

T

tact	25
tanner staging	28
temperament	21
temperamental guidance	60
tempo	16
Tertiary prevention	274

Test validity	184
theory of mind	43
theory-alignment map	42
theory-theory account	67
thinking	13
thomas and chess	54
threat	23
Tics	24
timing	32
toxic stress	151
trajectory	84
transactional	55
transactional model	56
transductive reasoning	14
Transition	24
transitive inference	12
triage	202
trivandrum	31
trivandrum development screening chart	31
tuberin	249
Tuberous sclerosis complex	248

U

unbalanced translocation	224
uneven profile	320
unexpected-contents task	65
uninvolved	43
uninvolved or neglectful parenting	43
universal	24
universal grammar	72
unspecified intellectual developmental disorder	179
upper limit	27
usage-based	73
useful speech	379

V

Vaccine causation	314
ventral striatum	126
very good outcome	378
Vigabatrin	240
Vineland Adaptive Behavior Scales	207
Vineland Social Maturity Scale	207
vulnerability factor	150
vygotsky	12

W

warning signs	31
when to refer	27
will	18
windows of achievement	93

wisdom 18

withdrawal 23

X

X-linked 218

Z

zone of proximal development 12

About the authors

Dr. Wilfred D'souza

MD Psychiatry (Resident) · Certified Gestalt Therapist · Weave, Centre for Integrative Psychiatry

Works in integrative psychiatry with Schema Therapy as the primary lens, alongside CBT, DBT, motivational interviewing and emotion-focused therapy, and is a certified Gestalt therapist. His clinical interests run through neurodivergence, ADHD, bipolar disorder, personality disorders, addiction and developmental psychiatry: work grounded in lived experience of attention difficulties long labelled as laziness, which made psychiatry personal before it became professional. He has been developing the Toroidal Model of Psychosis and Affect (TMoP), his computational model of mind, and TMoP thinking shaped the Claude Code AI system he builds with: clinical software, teaching tools and much of this series' production pipeline among its work, bringing psychiatry into the AI era in his own way. Lead author of the Weave Sprint series.

Dr. Niharika Reddy

Consultant Psychiatrist · Weave, Centre for Integrative Psychiatry

Consultant Psychiatrist at Weave and at Trijog, Mumbai, seeing people online through weave.clinic. Her work centres on relationships and family dynamics, family therapy and women's mental health, with a particular focus on neurodivergence in women and on adolescent mental health. Practises integrative, person-centred psychiatry. Co-author of this volume.

THE WEAVE LIBRARY

You revised from one. The whole shelf is open.

Field guides and references written in practice, for the people who do the work and the people living through it. Most are free to read and download.



field guides · references · for clinicians and families · free to read

weave.clinic/library

TAP OR SCAN TO OPEN



WEAVE STUDY GUIDES

When the night is short, read the guide.

Twenty-two topic guides across all four papers. Study notes, model answers, mnemonics, high-yield comparisons and previous-question frequency. Free, searchable, built for this exam.



22 guides · all four papers · fully searchable · free to read

weave.clinic/study


TAP OR SCAN TO OPEN



THE WEAVE FAMILY LIBRARY

Send the family home with more than a prescription.

Free, plain-language books for living with a mental-health condition, written for patients and the people who love them. Find yours by condition, by relationship, or by situation.



plain language · patients and families · five languages · free to read

weave.clinic/family

TAP OR SCAN TO OPEN



A PAPER A DAY

The whole written exam, in one day's reach.

Every paper of the MD psychiatry written exam in one free book, with a colour memory code and clickable navigation, for the last week, the last night, the last hour before the hall.



four papers · one volume · colour-coded · free to read

weave.clinic/revision

TAP OR SCAN TO OPEN





STUDY TOGETHER

You do not have to revise alone.

These notes started as messages in a room full of residents helping each other through. That room is still open. Bring your doubts, your corrections, and the questions the textbooks do not answer.

Join the study group




https://chat.whatsapp.com/H7TsLGK2wFl4bSe2r0Zfzn?mode=gi_t



PRINTING AND DISTRIBUTION

Notes Next Door (NND), Delhi is the official printing and distribution partner for Weave Sprint. The soft-copy PDFs and the complete print-ready set are always available at weave.clinic/sprint, should you prefer to print the notes yourself. Weave Sprint is open source and free, and as part of Weave's commitment to residents it will stay that way, forever. We do, however, strongly recommend a printed set from Notes Next Door: in our experience, they have been the most reliable printing partner we have worked with.



WEAVE SPRINT POWERED BY TRIJOG · PAPER III · VOLUME ONE

Volume One of the third paper: normal child development, the domains and milestones and the ages they are expected by, and then intellectual disability and autism spectrum disorder, how each is defined and assessed, what lies beneath them, and what can actually be done. Two complete sets of notes, written to be understood the first time and recalled every time after, with a colour memory code, a clickable contents and a full index.

Built to understand. Kept to revise.

All the very best for your exam.

WEAVE POWERED BY TRIJOG

weave.clinic · wilfred@weave.clinic · niharika@weave.clinic

